

## DAFTAR PUSTAKA

- Altenstadt, J.F.S., Hukkelhoven, C.W.P.M., Roosmalen, J., Bloemenkamp, K.W.M. 2013. Pre-Eclampsia Increases the Risk of Postpartum Haemorrhage: A Nationwide Cohort Study in The Netherlands. *PLOS ONE*, 12:8
- Altenstadt, J.F.S., Hukkelhoven, C.W.P.M., Roosmalen, J., Bloemenkamp, K.W.M. 2013. Pre-Eclampsia Increases the Risk of Postpartum Haemorrhage: A Nationwide Cohort Study in The Netherlands. *PLOS ONE*, 12:8
- Arnolds, D.E., Smith, A., Banayan, J.M., Holt, R., Scavone, B.M. (2018). National Partnership for Maternal Safety Recommended Maternal Early Warning Criteria Are Associated With Maternal Morbidity. *International Anesthesia Research Society*, 30 (30). <https://DOI:10.1213/ANE.0000000000003889>
- Aringer, M., Costenbader, K.H., Daikh, D.I., *et al.* (2020). 2019 EULAR/ACR Classification Criteria for Systemic Lupus Erythematosus. *Arthritis Rheumatol*, 71(9), p. 1400–1412
- Bartal, M.F., Lindheimer, M.D., Sibai, B.M. (2002). Proteinuria during pregnancy: definition, pathophysiology, methodology, and clinical significance. *Am J Obstet Gynecol*.
- Bircher, Charles. (2020). Trust Guideline for the use of the Modified Early Obstetric Warning Score (MEOWS) in Detecting the Seriously Ill and Deteriorating Woman. *NHS Foundation Trust*, 7
- Bozdog, Halenur, *et al.* (2015) . The frequency and fetomaternal outcomes of early- and late-onset preeclampsia: The experience of a single tertiary health center in the bustling metropolis of Turkey: Istanbul. *Medeniyet Medical Journal*, 30(4), p. 163-169
- Burch, V.C., Tarr, G., Morroni, C. (2008). Modified early warning score predicts the need for hospital admission and inhospital mortality. *Emerg Med J*, 25, p.674– 678
- Carle, C., Alexander, P., Columb, M. (2013). Design and internal validation of an obstetric early warning score : secondary analysis of the Intensive Care National

- Audit and Research Centre Case Mix Programme Database. *Anaesthesia*, 68, p.354-67.
- Cole, M.F. (2014). A Modified early obstetric warning system. *British Journal of Midwifery*, 22 (12)
- Cudmore, M., Ahmad, S., Al-Ani, B., Fujisawa, T., Coxall, H., *et al.* (2007). A: Negative regulation of soluble Flt-1 and soluble endoglin release by heme oxygenase-1. *Circulation* 115: 1789–1797
- Cunningham, F.G., Laveno, K.J., Bloom, S.L., Dashe, J.S. (2016). Hipertensive Disorder. *William Obstetric 25<sup>th</sup> ed*, p. 1608
- Cunningham, F.G., Laveno, K.J., Bloom, S.L., Dashe, J.S. (2016). Physiology of labor. *William Obstetric 25<sup>th</sup> ed*, p. 870
- Endradita, G. (2017). Panduan rawat inap rumah sakit. <https://galihendradita.wordpress.com/2017/04/23/panduan-rawat-inap-rumah-sakit/>. (Diakses 18 Desember 2021)
- Eskild, A., Vatten, L.J. (2009). Abnormal bleeding associated with preeclampsia: a population study of 315,085 pregnancies. *Acta Obstet Gynecol Scand* 88: 154–158.
- Espinoz, J., Vidaeff, A., Pettker, C.M., Simhan, Hyagriv. (2020). Gestational Hypertension and Preeclampsia. *ACOG Practice Bulletin*, 135, p.6
- Fong, A. *et al.* (2014). Pre-gestational versus gestational diabetes: A population based study on clinical and demographic differences. *J Diabetes Complications*, 28(1)
- Hannola, K., Hoppu, S., Mennander, S., Huhtala, S., Laivuori, H., Tihtonen, K. (2021). Obstetric early warning system to predict maternal morbidity of pre-eclampsia, postpartum hemorrhage and infection after birth in high-risk women: a prospective cohort study. *Midwifery*, 99
- Hoffman, M. (2020). Lung disease overview. <https://www.webmd.com/lung/lung-diseases-overview>. (Diakses 18 Desember, 2021)
- Jahan, E., Barua, T., Salma, U. (2014). An Overview On Heart Rate Monitoring And Pulse Oximeter System. *International Journal of Latest Research in Science and Technology*, 5: 148-152

- Kaduma, J., Seni, J., Chuma, C., Kirita, R., Mujuni, F., *et al.* (2019). Urinary Tract Infections and Preeclampsia among Pregnant Women Attending Two Hospitals in Mwanza City, Tanzania: A 1:2 Matched Case-Control Study. *BioMed Research International*, 1:8
- Kamus Besar Bahasa Indonesia. <https://kbbi.kemdikbud.go.id> (Diakses 18 Desember, 2021)
- Kaufmann, P., Black, S., Huppertz, B. (2003). Endovascular trophoblast invasion: Implications for the pathogenesis of intrauterine growth retardation and preeclampsia. *Biol Reprod* 69: 1–7.
- Kulandavelu, S., Whiteley, K.J., Qu D, Mu, J., Bainbridge, S.A., Adamson, S.L. (2012). Endothelial nitric oxide synthase deficiency reduces uterine blood flow, spiral artery elongation, and placental oxygenation in pregnant mice. *Hypertension* 60: 231–238.
- Lam, M.T.C., Dierking, E. (2017). Intensive Care Unit issues in eclampsia and HELLP syndrome. *Int J Crit Illn Inj Sci.* 7(3): 136–141.
- Levey, A.S., Levin, A., Kellum, J.A. (2013). Definition and Classification of Kidney Diseases. *Am J Kidney Dis*, 61(5), p.686-688
- Li, C., Gou, W., Liu, J., Liu, D., Li, X., Chen Q. (2021). Length of stay in the ward following a preeclamptic pregnancy. *Journal of Human Hypertension*. <https://doi.org/10.1038/s41371-021-00484-w>
- Liu, H., Allen J., Zheng, D., Chen, F. (2019). Recent development of respiratory rate measurement technologies. *Physiological Measurement*, 40: 7
- Lopez, E.O., Ballard, B.D., Jan, A. (2021). Cardiovascular Disease. <https://www.ncbi.nlm.nih.gov/books/NBK535419/>. (Diakses 18 Desember, 2021)
- Mangku, G., Senapathi, T.G.A. (2010). Tunjangan Homeostasis Perioperatif Dalam: Buku Ajar Ilmu Anestesia dan Reanimasi, hal 233-243.
- Mackintosh, N., Watson, K., Susanna, R., Jane, S. (2014). Value of a modified early obstetric warning system (MEOWS) in managing maternal complications in the peripartum period : an ethnographic study. *BMJ Qual Saf*; 23, p.26-34. <https://doi:10.1136/bmjqs-2012-001781>

- Martin, L.S. (2016). Superimposed Preeclampsia. *Adv Exp Med Biol*, 956, p.409-417
- Maynard, S.E., Min, J.Y., Merchan, J., Lim, K.H., Li, J., *et al.* (2003). Excess placental soluble fms-like tyrosine kinase 1 (sFlt1) may contribute to endothelial dysfunction, hypertension, and proteinuria in preeclampsia. *J Clin Invest* 111: 649–658.
- McCaig, D., Lyall, F. (2009). Inhibitors of heme oxygenase reduce invasion of human primary cytotrophoblast cells in vitro. *Placenta* 30: 536–538.
- Meazaw, *et al.* (2020). Systematic and meta-analysis of factors associated with preeclampsia and eclampsia in sub-Saharan Africa. PLOS ONE. <https://doi:10.1371/journal.pone.0237600>
- Mersha, Getnet, A., Abegaz, T.M., Seid, M.A. (2019). Maternal and perinatal outcomes of hypertensive disorders of pregnancy in Ethiopia: systematic review and metaanalysis. *BMC Pregnancy and Childbirth*, 19, p.458
- Mhyre, J.M., *et al.* (2014). The maternal early warning criteria : proposal from the national partnership for maternal safety. *Obstet Gynecol*, 124, p.782-6
- Namasivayam, P., Kalaivani. (2018). Implementation Of Meows In Predicting Maternal Morbidity And Using It To Improve Maternal Outcome. *Medical University Chennai*
- Paternina, A., *et al.* (2017). Performance of the Obstetric Early Warning Score in critically ill patients for the prediction of maternal death. *American Journal of Obstetrics & Gynecology*
- Phipps, E., Prasanna, D., Brima, W., Jim, B. (2016) Preeclampsia: Updates in Pathogenesis, Definitions, and Guidelines. *Clinical Journal of the American Society of Nephrology*
- Profil Kesehatan Provinsi DI Yogyakarta Tahun (2017). Tersedia di <http://www.kemendes.go.id> (diakses 13 September 2020)
- Quillon, A., Fromy, B., Debret, R. (2015). Endothelium microenvironment sensing leading to nitric oxide mediated vasodilation: A review of nervous and biomechanical signals. *Nitric Oxide* 45: 20–26.

- Rana, S., Lemoine, E., Granger, J., Karumanchi, SA. (2019). Preeclampsia Pathophysiology, Challenges, and Perspectives. *Circulation Research*, March 29. <https://DOI: 10.1161/CIRCRESAHA.118.313276>
- Ram, M., Anteby, M., Weiniger, C.F., Havakuk, O., Gilboa, I., *et al.* (2021). Acute pulmonary edema due to severe preeclampsia in advanced maternal age women. *Pregnancy Hypertension: An International Journal of Women's Cardiovascular Health* 25:150–155.
- RCOG. Multiple pregnancy: having more than one baby. <https://www.rcog.org.uk/globalassets/documents/patients/patient-information-leaflets/pregnancy/pi-multiple-pregnancy.pdf> (diakses 17 Desember, 2021).
- Ryan, H.M., Jones, M.A., Payne B.A., Sharma, S, Hutfield A.M., *et al.* (2017). Validating the Performance of the Modified Early Obstetric Warning System Multivariable Model to Predict Maternal Intensive Care Unit Admission. *J Obstet Gynaecol Can*; 39(9):728-733.
- Shields, L.E., Wiesner, S., Klein, C., Pelletreau, B., Hedriana, H.L. (2016). Use of Maternal Early Warning Trigger tool reduces maternal morbidity. *American Journal of Obstetric and Gynecology*, 214, p.527.e1-6.
- Singh, S., McGlennan, A., England, A., Simons, R. (2012). A validation study of the CEMACH recommended modified early obstetric warning system (MEOWS). *Anaesthesia*, 67, p.12-8. <https://doi:10.1111/j.1365-2044.2011.06896.x>.
- Sivakrishnan, S. (2018). Liver Diseases-An Overview. *World journal of pharmacy and pharmaceutical science*, 8(1), p.1385-1395
- Smith, V., Kenny, L.C., Sandall, J., Devane, D. (2019). Physiological track and trigger/ early warning systems for use in maternity care. *Cochrane Database of Systematic Reviews*, Issue 6. doi: [10.1002/14651858.CD013276](https://doi.org/10.1002/14651858.CD013276)
- Tamara, Y., Lutfi, M., Prawitasari, S., (2018). Hubungan Maternal Early Obstetrics Warning Score (MEOWS) dengan perawatan di Intensive Care Unit pada pasien Preeklamsia Berat di RSUP Dr. Sardjito. *Tesis*. FKMK UGM, Yogyakarta

- Thorpe, J.G., Love, N., Wrightson, J., Walsh, S., Keeling, N. (2006). Modified Early Warning Score (MEWS) in surgical in-patients: a prospective observational study. *Ann R Coll Surg Engl*; 88: 571–575 doi 10.1308/003588406X130615
- Too, G.T., Hill, J.B. (2013). Hypertensive crisis during pregnancy and postpartum period. *Seminars In Perinatology* : 37, 280–287
- Tuyishime, E., Ingabire, H., Mvukiyehe, J.P., Durieux, M., Twagirumugabe, T. (2020). Implementing the Risk Identification (RI) and Modified Early Obstetric Warning Signs (MEOWS) tool in district hospitals in Rwanda: a cross-sectional study. *BMC Pregnancy and Childbirth*. 20:568. <https://doi.org/10.1186/s12884-020-03187-1>
- Umar, A., Ameh, C.A., Muriithi, F., Mathai, M. (2019). Early warning systems in obstetrics: A systematic literature review. *PLoS ONE*, 14(5)
- Venkatesha, S., Toporsian, M., Lam, C., Hanai, J., Mammoto, T, *et al.* (2006). Soluble endoglin contributes to the pathogenesis of preeclampsia. *Nat Med* 12: 642–649.
- Wen, T., Yu, V.X., Wright, J.D., Goffman, D., Attenello, F., *et al.* (2018). Postpartum length of stay and risk for readmission among women with preeclampsia. *The Journal of Maternal-Fetal & Neonatal Medicine*. <https://doi:10.1080/14767058.2018.1514382>
- Wibowo, N *et al.* (2016). PNPk Diagnosis dan Tata Laksana Preeklamsia. POGI, HKFM
- Zaid, AA, *et al.* (2020). Advanced Maternal Age and the Frequency of Preeclampsia-A Single-Center Cross Sectional Study from Saudi Arabia. *J. Evolution Med. Dent. Sci Sept*, 14. <https://DOI: 10.14260/jemds/2020/592>