

DAFTAR PUSTAKA

- Abdelgawad, E.A., Abu-samra, M.F., Abdelhay, N.M., & Abdel-Azeem, H.M., 2020. B-mode ultrasound, color doppler, and sonoelastography in differentiation between benign and malignant cervical lymph nodes with special emphasis on sonoelastography. *Egypt. J. Radiol. Nucl. Med.* 51: 157. doi:10.1186/s43055-020-00273-4
- Adham, M., Kurniawan, A.N., Muhtadi, A.I., Roezin, A., Hermani, B., Gondhowiardjo, S., et al., 2012. Nasopharyngeal carcinoma in Indonesia: epidemiology, incidence, signs, and symptoms at presentation. *Chin. J. Cancer* 31: 185–196. doi:10.5732/cjc.011.10328
- Ahuja, A.T., 2008. Ultrasound of malignant cervical lymph nodes. *Cancer Imaging* 8: 48–56. doi:10.1102/1470-7330.2008.0006
- Ahuja, A.T., Ho, S.S.Y., Leung, S.F., Kew, J., & Metreweli, C., 1999. Metastatic adenopathy from nasopharyngeal carcinoma: successful response to radiation therapy assessed by color duplex sonography. *AJNR. Am. J. Neuroradiol.* 20: 151–6
- Ahuja, A.T., & Ying, M., 2004. Evaluation of cervical lymph node vascularity: a comparison of colour doppler, power doppler and 3-D power doppler sonography. *Ultrasound Med. Biol.* 30: 1557–1564. doi:10.1016/j.ultrasmedbio.2004.09.002
- Ai, Q.-Y.H., Hung, K.F., So, T.Y., Mo, F.K.F., Tsung Anthony Chin, W., Hui, E.P., et al., 2022. Prognostic value of cervical nodal necrosis on staging imaging of nasopharyngeal carcinoma in era of intensity-modulated radiotherapy: a systematic review and meta-analysis. *Cancer Imaging* 22: 24. doi:10.1186/s40644-022-00462-6
- Amanah, N.S., 2020. Faktor-faktor penyebab terjadinya karsinoma nasofaring (KNF). *WELLNESS Heal. Mag.* 2: 113–120
- Azizah, N., Hanriko, R., & Ramkita, N., 2017. Hubungan antara konsumsi ikan asin, ikan / daging asap, dan makanan kaleng dengan karsinoma Nasofaring di RSUD Abdul Moeloek periode tahun 2014-2016. *J. Kedokt. Univ. Lampung* 4: 260–263.
- Bialek, E.J., & Jakubowski, W., 2017. Mistakes in ultrasound diagnosis of superficial lymph nodes. *J. Ultrason.* 17: 59–65. doi:10.15557/JoU.2017.0008
- Burusapat, C., Jarungroongruangchai, W., & Charoenpitakchai, M., 2015. Prognostic factors of cervical node status in head and neck squamous cell carcinoma. *World J. Surg. Oncol.* 13: 51. doi:10.1186/s12957-015-0460-6
- Canreg, 2020. RKBR januari 2020. *Jogja Cancer Regist.* Available from: <https://canreg.fk.ugm.ac.id/laporan-data/registrasi-kanker-berbasis-rumah-sakit-dr-sardjito-fkkmk-ugm/januari-2020/>

- Chen, L., Qi, L., Zhang, J., Ma, Q., & Chai, X., 2021. Neutrophil–lymphocyte ratio as a prognostic factor for minute clear cell renal cell carcinoma diagnosed using multi-slice spiral CT. *Medicine (Baltimore)*. 100: e26292. doi:10.1097/MD.00000000000026292
- Dahlan, S., 2017. Statistik untuk kedokteran dan kesehatan, 6th ed. Salemba Medika, Jakarta.
- Dahlan, S., 2010. Besar sampel dan cara pengambilan sampel, Salemba Medika, Jakarta.
- Dhara, V., Hoda, N., Rajini, B., Sabitha, K., Vinitha, A., & Nathani, J., 2021. Significance of cervical node necrosis in preoperative MRI as a prognostic indicator: retrospective study of patients with SCC of tongue. *J. Oral Med. Oral Surg.* 27: 43. doi:10.1051/mbcb/2021021
- Feng, Y., Zhang, N., Wang, S., Zou, W., He, Y., Ma, J., et al., 2020. Systemic inflammation response index Is a predictor of poor survival in locally advanced nasopharyngeal carcinoma: a propensity score matching study. *Front. Oncol.* 10: 1–7. doi:10.3389/fonc.2020.575417
- Friborg, J.T., Yuan, J.-M., Wang, R., Koh, W., Lee, H., & Yu, M.C., 2007. A prospective study of tobacco and alcohol use as risk factors for pharyngeal carcinomas in Singapore Chinese. *Cancer* 109: 1183–1191. doi:10.1002/cncr.22501
- Hardianti A, 2016. Faktor risiko kanker nasofaring di Rumah Sakit Hasan Sadikin Bandung. *Fak. Kedokt. Univ. Padjadjaran* 1–9
- He, L., Xie, X., Xue, J., Xie, H., & Zhang, Y., 2022. Association of the systemic immune-inflammation index with all-cause mortality in patients with arteriosclerotic cardiovascular disease. *Front. Cardiovasc. Med.* 9: 1–10. doi:10.3389/fcvm.2022.952953
- Ho, F.C., Tham, I.W., Earnest, A., Lee, K.M., & Lu, J.J., 2012. Patterns of regional lymph node metastasis of nasopharyngeal carcinoma: a meta-analysis of clinical evidence. *BMC Cancer* 12: 98. doi:10.1186/1471-2407-12-98
- Hogan, L., 2021. What is lymph node metastasis learn about how cancer can affect the lymph nodes. Available from: <https://www.webmd.com/cancer/what-to-know-lymph-nodes>
- Hong, Y.T., Ngoc Minh, P.H., & Hong, K.H., 2016. An unusual metastasis of posterior neck and axillary lymph nodes from nasopharyngeal carcinoma. *Korean Soc. Head Neck Oncol.* 32: 23–27. doi:10.21593/kjhno/2016.32.2.23
- Islami, H., & Hariwiyanto, B., 2012. Alel gen HLA-A2402 sebagai faktor risiko terjadinya karsinoma nasofaring di RSUP Dr Sardjito. Available from: http://etd.repository.ugm.ac.id/home/detail_pencarian/53701

- Ji, Y., & Wang, H., 2020. Prognostic prediction of systemic immune-inflammation index for patients with gynecological and breast cancers: a meta-analysis. *World J. Surg. Oncol.* 18: 197. doi:10.1186/s12957-020-01974-w
- Jiang, W., Chen, Y., Huang, J., Xi, D., Chen, J., Shao, Y., et al., 2017. Systemic immune-inflammation index predicts the clinical outcome in patients with nasopharyngeal carcinoma: a propensity score-matched analysis. *Oncotarget* 8: 66075–66086. doi:10.18632/oncotarget.19796
- KEMENKES RI, 2017. Pedoman nasional pelayanan kedokteran. Kanker nasofaring. Kementerian Republik Indonesia, Jakarta
- Kenel, T., Garrel, R., Costes, V., Boisselier, P., Crampette, L., & Favier, V., 2019. Head and neck carcinoma of unknown primary. *Eur. Ann. Otorhinolaryngol. Head Neck Dis.* 136: 185–192. doi:10.1016/j.anorl.2019.04.002
- King, A.D., 2022. MR imaging of nasopharyngeal carcinoma. *Magn. Reson. Imaging Clin. N. Am.* 30: 19–33. doi:10.1016/j.mric.2021.06.015
- King, A.D., Tse, G.M.K., Ahuja, A.T., Yuen, E.H.Y., Vlantis, A.C., To, E.W.H., et al., 2004. Necrosis in metastatic neck nodes: diagnostic accuracy of CT, MR imaging, and US. *Radiology* 230: 720–726. doi:10.1148/radiol.2303030157
- Kölküçü, E., Atılgan, D., Kuyucu, Y.E., Özbek, L.M., & Unsal, V., 2021. Relationship between lymph node metastasis and lymph node density and preoperative neutrophil-lymphocyte ratio in patients undergoing radical cystectomy. *Bull. Urooncology* 20: 168–173. doi:10.4274/uob.galenos.2021.1809
- Lau, H.-Y., Leung, C.-M., Chan, Y.-H., Lee, A.W.-M., Kwong, D.L.-W., Lung, M.L. i., et al., 2013. Secular trends of salted fish consumption and nasopharyngeal carcinoma: a multi-jurisdiction ecological study in 8 regions from 3 continents. *BMC Cancer* 13: 298. doi:10.1186/1471-2407-13-298
- Li, Yongchao, Liu, M., Cui, Y., Zhu, Z., Chen, J., Zeng, F., et al., 2022. Increased risk of testosterone deficiency is associated with the systemic immune-inflammation index: a population-based cohort study. *Front. Endocrinol. (Lausanne)*. 13: 1–10. doi:10.3389/fendo.2022.974773
- Liang, C., Xu, Z., Shen, X., & Wu, K., 2022. Correlation between neutrophil-to-lymphocyte ratio and pretreatment magnetic resonance imaging and their predictive significance in cervical carcinoma patients referred for Radiotherapy. *J. Oncol.* 2022: 1–9. doi:10.1155/2022/3409487
- Lyshchik, A., Higashi, T., Asato, R., Tanaka, S., Ito, J., Hiraoka, M., et al., 2007. Cervical lymph node metastases: diagnosis at sonoelastography—initial experience. *Radiology* 243: 258–267. doi:10.1148/radiol.2431052032
- Mao, Y., Hedgire, S., & Harisinghani, M., 2014. Radiologic assessment of lymph nodes in oncologic patients. *Curr. Radiol. Rep.* 2: 36. doi:10.1007/s40134-013-0036-6

- Miura, Y., Mikada, M., Ouchi, T., Horie, S., Takeda, K., Yamaki, T., et al., 2016. Early Diagnosis of lymph node metastasis: importance of intranodal pressures. *Cancer Sci.* 107: 224–232. doi:10.1111/cas.12873
- Morimoto, Y., Kurokawa, H., Tanaka, T., Yamashita, Y., Kito, S., Okabe, S., et al., 2006. Correlation between the incidence of central nodal necrosis in cervical lymph node metastasis and the extent of differentiation in oral squamous cell carcinoma. *Dentomaxillofacial Radiol.* 35: 18–23. doi:10.1259/dmfr/24536918
- Mouawad, F., Rysman, B., Russ, G., Benoudiba, F., Garcia, G., Abgral, R., et al., 2019. Cystic form of cervical lymphadenopathy. Guidelines of the French Society of Otorhinolaryngology - Head and Neck Surgery (SFORL). Part 1: Diagnostic procedures for lymphadenopathy in case of cervical mass with cystic aspect. *Eur. Ann. Otorhinolaryngol. Head Neck Dis.* 136: 489–496. doi:10.1016/j.anorl.2019.05.015
- Moulanda, F., & Adham, M., 2021. Neck residue of nasopharyngeal carcinoma (NPC): Timing of diagnosis and treatment. *Int. J. Nasopharyngeal Carcinoma* 03: 17–19. doi:10.32734/ijnpc.v3i01.5585
- Nalbant, A., Demirci, T., Kaya, T., Aydın, A., Altındış, M., & Güçlü, E., 2021. Can prognostic nutritional index and systemic immune-Inflammatory index predict disease severity in COVID-19? *Int. J. Clin. Pract.* 75. doi:10.1111/ijcp.14544
- Nathanson, S.D., 2003. Insights into the mechanisms of lymph node metastasis. *Cancer* 98: 413–423. doi:10.1002/cncr.11464
- Nathanson, S.D., Rosso, K., Chitale, D., & Burke, M., 2017. Lymph node metastasis, in: introduction to cancer metastasis. Elsevier, pp. 235–261. doi:10.1016/B978-0-12-804003-4.00013-X
- Nguyen Van, D., Nguyen, T.B., Nguyen Thi, N.T., & Le Van, Q., 2021. Report on unusual sites of lymph node metastases in nasopharyngeal carcinoma. *Case Rep. Oncol.* 14: 1821–1826. doi:10.1159/000520977
- Notopuro H, Kentjono W, Handayani R, N.P., 2008. Karsinoma nasopharynx dan infeksi EBV di Indonesia; analisis aspek klinis, patologi dan biomolekular. *J. Yars.* 16: 1–19
- Nugraheni, R.A., & Waliyanti, E., 2019. Faktor risiko kanker nasofaring di kabupaten sleman yogyakarta. Available from: <http://repository.umy.ac.id/handle/123456789/28584>
- Okepa, S.I., S M N Mydin, R.B., Mangantig, E., Azmi, N.S.A., Zahari, S.N.S., Kaur, G., et al., 2019. Nasopharyngeal carcinoma (NPC) risk factors: a systematic review and meta-analysis of the Aassociation with lifestyle, Diets, Socioeconomic and Sociodemographic in Asian Region. *Asian Pacific J. Cancer Prev.* 20: 3505–3514. doi:10.31557/APJCP.2019.20.11.3505

- Pakoz, Z.B., Ustaoglu, M., Vatansever, S., Yuksel, E.S., & Topal, F., 2022. Serum immune-inflammation index assessment in the patients with ulcerative colitis. *Gastroenterol. Res. Pract.* 2022: 1–5. doi:10.1155/2022/9987214
- Pattanayak, S., Chatterjee, S., Ravikumar, R., Nijhawan, V.S., Vivek Sharma, & Debnath, J., 2018. Ultrasound evaluation of cervical lymphadenopathy: Can it reduce the need of histopathology/cytopathology? *Med. J. Armed Forces India* 74: 227–234. doi:10.1016/j.mjafi.2017.04.005
- Pisani, P., Airoidi, M., Allais, A., Aluffi Valletti, P., Battista, M., Benazzo, M., et al., 2020. Metastatic disease in head & neck oncology. *Acta Otorhinolaryngol. Ital.* 40: S1–S86. doi:10.14639/0392-100X-suppl.1-40-2020
- Quang Huy, H., 2020. Role of Computed tomography imaging in evaluation cervical lymph nodes in patients with nasopharyngeal carcinoma. *Biomed. J. Sci. Tech. Res.* 27: 20988–20992. doi:10.26717/BJSTR.2020.27.004546
- Rahman, S., Budiman, B.J., & Subroto, H., 2015. Faktor risiko non viral pada karsinoma nasofaring. *J. Kesehat. Andalas.* 4: 988–995.
- Ren, A., Li, Z., Zhang, X., & Deng, R., 2020. Inflammation-based prognostic scores in patients with Hepatitis B virus-related hepatocellular carcinoma after liver transplantation. *J. Hepatocell. Carcinoma* 7: 101–106
- RISKESDAS, 2013. Riset Kesehatan Dasar. Badan Penelitian dan Pengembangan Kesehatan Kementerian Kesehatan RI, Jakarta
- Rueda Domínguez, A., Cirauqui, B., García Castaño, A., Alvarez Cabellos, R., Carral Maseda, A., Castelo Fernández, B., et al., 2022. SEOM-TTCC clinical guideline in nasopharynx cancer (2021). *Clin. Transl. Oncol.* 24: 670–680. doi:10.1007/s12094-022-02814-x
- Salehiniya, H., Mohammadian, M., Hashejani, A., & MahdaviFar, N., 2018. Nasopharyngeal cancer in the world: epidemiology, incidence, mortality and risk factors. *World Cancer Res. J.* 5: 1–8.
- Sastroasmoro, S., & Ismael, S., 2011. Dasar-dasar metodologi penelitian klinis. CV. Sagung Seto, Jakarta
- Seo, J.W., Kim, Jun-hwee, Lee, S., Sh, B., Jw, S., Jh, K., et al., 2019. Prognostic value of cervical nodal necrosis observed in preoperative 437–443.
- Sharma, A., Jaiswal, A.A., Umredkar, G., Barle, R., Sharma, N., Banerjee, P.K., et al., 2017. Lymph node central necrosis on the computed tomography as the predictor of the extra capsular spread in metastatic head and neck squamous cell carcinoma. *Indian J. Otolaryngol. Head Neck Surg.* 69: 323–332. doi:10.1007/s12070-017-1131-4

- Sharma, T.D., Singh, T.T., Laishram, R.S., & Durlav, L., 2011. Nasopharyngeal carcinoma - a clinico-pathological study in a regional cancer centre of northeastern india. *APJCP*. 12 (6): 1583-7
- Sugiyono, D., 2013. Metode penelitian kuantitatif, kualitatif, dan R & D. Alfabeta, Bandung.
- Suta, D., Saputra, D., & Sutanegara, D., 2019. Profil penderita kanker nasofaring di Rumah Sakit Umum Pusat Sanglah Denpasar periode januari – desember tahun 2014. *E-Jurnal Med*. 8: 1–14.
- Takenaka, Y., Kitamura, T., Oya, R., Ashida, N., Shimizu, K., Takemura, K., et al., 2017. Prognostic role of neutrophil-lymphocyte ratio in nasopharyngeal carcinoma: a meta-analysis. *PLoS One* 12: e0181478. doi:10.1371/journal.pone.0181478
- Tomita, N., Fuwa, N., Arij, Y., Kodaira, T., & Mizoguchi, N., 2011. Factors associated with nodal metastasis in nasopharyngeal cancer: an approach to reduce the radiation field in selected patients. *Br. J. Radiol.* 84: 265–70. doi:10.1259/bjr/47164832
- Wang, H.-Y., Chang, Y.-L., To, K.-F., Hwang, J.S.G., Mai, H.-Q., Feng, Y.-F., et al., 2016. A new prognostic histopathologic classification of nasopharyngeal carcinoma. *Chin. J. Cancer* 35: 41. doi:10.1186/s40880-016-0103-5
- Wang, J., Zhou, D., Dai, Z., & Li, X., 2021. Association between systemic immune-inflammation index and diabetic depression. *Clin. Interv. Aging* 16: 97–105.
- Wang, P., Yue, W., Li, W., Luo, Y., Li, Z., Shao, Y., et al., 2019. Systemic immune-inflammation index and ultrasonographic classification of breast imaging-reporting and data system predict outcomes of triple-negative breast cancer. *Cancer Manag. Res.* Volume 11: 813–819. doi:10.2147/CMAR.S185890
- Wang, Q., & Zhu, D., 2019. The prognostic value of systemic immune-inflammation index (SII) in patients after radical operation for carcinoma of stomach in gastric cancer. *J. Gastrointest. Oncol.* 10: 965–978. doi:10.21037/jgo.2019.05.03
- Wang, Y.-T., Kuo, L.-T., Weng, H.-H., Hsu, C.-M., Tsai, M.-S., Chang, G.-H., et al., 2022. Systemic immun inflammation index as a predictor for head and neck cancer prognosis: a meta-analysis. *Front. Oncol.* 12: 1–9. doi:10.3389/fonc.2022.899518
- Wardhana, A.A., & Lesmana, T., 2021. Neutrophil lymphocyte ratio (NLR) as a predictive factor radiological response of neoadjuvant chemotherapy (NAC) in locally advanced rectal cancer (LARC). *Bali Med. J.* 10: 717. doi:10.15562/bmj.v10i2.2483
- Winter, M., Gibson, R., Ruszkiewicz, A., Thompson, S.K., & Thierry, B., 2015. Beyond conventional pathology: Towards preoperative and intraoperative lymph node staging. *Int. J. Cancer* 136: 743–751. doi:10.1002/ijc.28742

- Wu, J., Yan, L., & Chai, K., 2021. Systemic immune-inflammation index is associated with disease activity in patients with ankylosing spondylitis. *J. Clin. Lab. Anal.* 35: 1–8. doi:10.1002/jcla.23964
- Xie, C., Li, H., Yan, Y., Liang, S., Li, Y., Liu, L., et al., 2020. A nomogram for predicting distant metastasis using nodal-related features among patients with nasopharyngeal carcinoma. *Front. Oncol.* 10: 1–9. doi:10.3389/fonc.2020.00616
- Xu, Y., Huang, T., Fan, L., Jin, W., Chen, X., & Chen, J., 2019. Patterns and prognostic value of lymph node metastasis on distant metastasis and survival in nasopharyngeal carcinoma: a surveillance, epidemiology, and end results study, 2006–2015. *J. Oncol.* 2019: 1–8. doi:10.1155/2019/4094395
- Ying, M., & Ahuja, A.T., 2006. Ultrasound of neck lymph nodes: How to do it and how do they look? *Radiography* 12: 105–117. doi:10.1016/j.radi.2005.04.004
- Ying, M., Bhatia, K.S.S., Lee, Y.P., Yuen, H.Y., & Ahuja, A.T., 2014. Review of ultrasonography of malignant neck nodes: greyscale, doppler, contrast enhancement and elastography. *Cancer Imaging* 13: 658–669. doi:10.1102/1470-7330.2013.0056
- Zhang, Y., Chen, Z., Jin, F., Guo, D., Chen, Q., Liu, Z., et al., 2021. The value of the systemic immune-inflammation index in predicting survival outcomes in patients with brain metastases of non-small-cell lung cancer treated with stereotactic radiotherapy. *Mediators Inflamm.* 2021: 1–10. doi:10.1155/2021/2910892
- Zoumalan, R.A., Kleinberger, A.J., Morris, L.G.T., Ranade, A., Yee, H., DeLacure, M.D., et al., 2010. Lymph node central necrosis on computed tomography as predictor of extracapsular spread in metastatic head and neck squamous cell carcinoma: Pilot Study. *J. Laryngol. Otol.* 124: 1284–1288. doi:10.1017/S0022215110001453