

## DAFTAR PUSTAKA

- Ahmed, M. A. S. & Mohammed, S. A. 2018. Neutrophyl-to-lymphocyte ratio as a prognostic marker in critically-ill septic patients. *Research and Opinion in Anesthesia and Intensive Care*, Volume 5, pp. 279-286.
- Ayala, A., Herndon, C. & Lehman, D. 1996. Differential induction of apoptosis in lymphoid tissue during sepsis: variation in onset, frequency and nature of the mediators. *Blood*, 87(10), pp. 4261-4275.
- Chandra, A., Enkhbaatar, P., Nakano, Y., Traber, LD., Traber, D.L ., 2006. Sepsis: Emerging Role of Nitric oxide and Selectins. *CLINICS*, 61(1), pp. 71-6.
- Dahlan, M. S., 2018. *Langkah-langkah Membuat Proposal Penelitian Bidang Kedokteran dan Kesehatan*. 2 ed. Jakarta: CV Sagung Seto.
- De Jager, C. P. Wever, P, Gemen, E. Kusters, R. Gageldonk-Lafeber, V. 2012. The Neutrophyl-Lymphocyte Count Ratio in Patients with Community-Acquired Pneumonia. *PLOS ONE*, 7(10).
- Finfer, S. & Machado, F. R. 2016. The Global Epidemiology of Sepsis. Does It Matter That We Know So Little?. *ATS Journals*, Volume 193, pp. 228-230.
- Fleischmann, C., Scherag, A. & Adhikari, N. K. 2016. Assessment of Global Incidence and Mortality of Hospital-treated Sepsis. Current Estimates and Limitations. *American Journal of Respiratory and Critical Care Medicine*, Volume 193(3), pp. 259-72.
- Fujishima, S. 2016. Organ Dysfunction as A New Standard for Defining Sepsis. *Inflammation and Regeneration*, 36(24), pp. 1-6.
- Gul, F., Arslantas, M. K., Cinel, I. & Kumar, A. 2017. Changing Definitions of Sepsis. *Turk J Anesthesiol Reanim*, Volume 45, pp. 129-38.
- Gürağaç, A. & Demirel, Z. 2016. The Neutrophyl-to-Lymphocyte Ratio in Clinical Practice. *Can Urol Assoc J*, 10(3-4), pp. 141-142.
- Guyton, A. C. & Hall, E. J. 2006. *Buku Ajar Fisiologi Kedokteran*. 11 ed. Jakarta: EGC

Gyawali, Bishal. Ramakrishna, Karan. Dhamoon, Amit. 2019. Sepsis : The Evolution in Definition, Patophysiology, and Management. *SAGE Open Medicine* 7 : 1-13

Hotchkiss, R., Swanson, P. & Freeman, B. 1999. Apoptotic cell death in patients with sepsis, shock and multiple organ dysfunction. *Crit Care Med*, 27(7), pp. 1230-51.

Jilma, B., Blann, A., Pernerstorfer, T. & Stohlawetz, P. 1999. Regulation of Adhesion Molecules During Human Endotoxaemia. *Amer J Resp Crit*, Volume 159, pp. 857-863.

Kaushik, R. Gupta, M. Jash, D. Sharma, M. Jain, N. Sinha, N. et al. 2018. Diagnostic and Prognostic Role of Neutrophil-to-Lymphocyte Ratio in Early and Late Phase of Sepsis. *Indian J Crit Care Med* 22 (9) : 660-663

Kaml, G. J. & Davis, K. A. 2016. Surgical Critical Care for The Patient With Sepsis and Multiple Organ Dysfunction. *Anesthesiology Clin*, Volume 34, pp. 681-96.

Kim, H. R., G, B. R. & Lee, M. K. 2008. Effect of bacteria and yeast on WBC counting in three automated hematology counters. *Ann Hematol*, Volume 87, pp. 557-562.

Korkmaz, Pinar. Erarslan, Sertaz. Toka, Onur. 2016. Evaluation of the Association Between the Neutrophil to Lymphocyte Ratio and Mortality in The Patients Followed up With Diagnosis of Sepsis. *J Clin Anal Med* : 1-6

Koyama, K., Katayama, S., Muronoi, T. 2018. Time Course of Immature Platelet Count and Its Relation to Thrombocytopenia and Mortality in Patients with Sepsis. *Journal PLOS*. 1(30)., p : 1-16

Lie, K. C., Lau, C. Y., Chau, N., West, E. 2018. Utility of SOFA Score, Management and Outcomes of Sepsis in Southeast Asia: A Multinational Multicenter Prospective Observational Study. *Journal of Intensive Care*, 6(9), pp. 1-8.

Liu, X. Shen, Y. Wang, H. Qinmin, G. Fei, A. Pan, S. 2016. Prognostic Significance of Neutrophil-to-Lymphocyte Ratio in Patients with Sepsis : A Prospective Observational Study. Hindawi Publishing Corporation

Liu Y, Wang H, Li Y, Shu Y, Ni J. 2019. Neutrophil to Lymphocyte Ratio (NLR) as A Prognostic Marker for In-hospital Mortality of Patients with Sepsis. *Medicine* 98:46, pp. 1-8

Martins, Eduardo. Da De Silveira, Lilian. Viegas, Karin. 2019. Neutrophil-Lymphocyte Ratio in The Early Diagnosis of Sepsis in an Intensive Care Unit : A Case-Control Study. *Rev Bras Ter Intensiva* 31 (1) : 63-70

Nugroho, A., Suwarman & Nawawi, A. M. 2013. Hubungan antara rasio Neutrofil-Limfosit dan Skor Sequential Organ Failure Assessment pada Pasien yang Dirawat di Ruang Intensif Care Unit. *Jurnal Anestesi Perioperatif*, 1(3), pp. 189-196.

Okashah, A. Abdel, R. 2014. Ratio of Neutrophyl to Lymphocyte Counts as a Simple Marker for Sepsis and Severe Sepsis in Intensive Care Unit. *ROAIC*, Volume 2, pp. 39-46.

Reinhart, K., Bauer, M., Riedemann, N. C. & Hartog, C. S. 2012. New Approaches to Sepsis: Molecular Diagnostics and Biomarkers. *Clinical Microbiology Reviews*, 25(4), pp. 609-634.

Riandy, Muhammad David. Purba, Bernhard Arianto. Puspita, Yuni. Rizal, Zainal. 2019. Hubungan Rasio Neutrofil Limfosit dengan Mortalitas  $\leq 28$  Hari di General Intensive Care Unit RSUP Dr. Mohammad Husein Palembang. *Jurnal Anestesi Perioperatif* 7 (3) : 188-97

Rehman, F., Khan, A., Aziz, A. 2020. Neutrophil to Lymphocyte Ratio : Earliest and Efficacious Markers of Sepsis. *Cureus*. 12 (10)., pp 120-128

Rello Jordi, Valenzuela-Sanchez. 2017. Sepsis: A Review Of Advances in Management. *Adv Ther*, Volume 34, pp. 2393-2411.

Sakr, Y. 2018. Sepsis in Intensive care Unit Patients: Worldwide Data From the Intensive Care over Nations Audit. *Open Forum Infectious Disease*, pp. 1-9.

Saliccioli, J. Marshall, D. Pimental, M. Santos, M. Pollard, T. Celi, L. et al. 2015. The Association Between The Neutrophil-to-lymphocyte Ratio and Mortality in Critical Illness : An Observational Cohort Study. *Crit Care* 19 (1) : 13

Sarwono, J., 2006. *Metode Penelitian Kuantitatif dan Kualitatif*. Yogyakarta: Graha Ilmu.

Sastroasmoro, S. & Ismael, S. 2014. *Dasar-dasar Metodologi Penelitian Klinis*. 5 ed. Jakarta: Sagung Seto.

Shen, Xiao Fei. Cao, Ke. Jiang, Jin-Peng. 2017. Neutrophyl Disregulation During Sepsis : An Overview and Update. *J Cell Mol Med* 9 : 1686-1697

Savran, Y., Mengi, T. Yilmaz, B. Tokur, M. Ergan, B., Yaka, E. et al. 2018. Can Neutrophyl-Lymphocyte Ratio Predict Disease Severity and Mortality in Sepsis and Septic Shock Patients?. *Acta Medica Mediterranea*, Volume 34, pp. 877-81.

Singer, M. Deutschman, C. Seymour, C. Shankar-Hari, M. Annane, D. Bauer, M et al, 2016. The Third International Consensus Definitions for Sepsis and Septic Shock (Sepsis-3). *JAMA*, Volume 315, pp. 801-10.

Suwondo, Victoria. Jatmiko, Heru Dwi. Hendrianingtyas, Meita. 2015. Karakteristik Dasar Pasien Sepsis Yang Meninggal di ICU RSUP dr. Kariadi Semarang Periode 1 Januari-31 Desember 2014. *Media Medika Muda* 4 (4) : 1586-1596

Tambajong, Rheza. Lalenoh, Diana. Kumaat, L. 2016. Profil Penderita Sepsis di ICU RSUP Prof. Dr. RD Kandou Manado Periode Desember 2014-November 2015. *Jurnal e-Clinic* 4 (1) : 452-7

Torio, C. M. & Andrews, R. M. 2013. National Inpatient Hospital Cost: The Most Expensive Conditions by Payer, 2011.. *Statistical Brief #160. Healthcare Cost and Utilization Project (HCUP) Statistical Briefs*.

Vardon-Bounes, F., Ruiz, S., Gratacap, M. 2019. Platelets Are Critical Key Players in Sepsis. *Int J Mol Sci* 20 : 3494

Velissaris, D. Pantzaris, N-D. Bountouris, P. Gogos, C. 2018. Correlations Between Neutrophyl-to-Lymphocyte Ratio and Severity Scores Upon Hospital Admission. A series of 50 patients. *ROM. J. INTERN. MED*, 0(0), pp. 1-10.

Venkata, C., Kashyap, R., Farmer, J.C. 2013. Thrombocytopenia in Adult Patients With Sepsis : Incidence, Risk Factors, and Its Association With Clinical Outcome. *Journal of Intensive Care* 1 : 9 p : 1-10

Xu, J., Tong, L., Guo, Z. Lui, KY, Hu, X. Cao, L., et al. 2019. Association of Sex With Clinical Outcome in Critically Ill Sepsis Patients : A Retrospective Analysis of The Large Clinical Database MIMIC-III. *Shock*, 52 (2), pp.146-151

Zahorec, R., 2001. Ratio of Neutrophyl to Lymphocyte Counts - Rapid and Simple Parameter of Systemic Inflammation and Stress in Critically Ill. *BRATISL LEK LISTY*, 102(1), pp. 5-14

Zhou, Z., Feng, T., Xie, Y., et al. 2020. Prognosis and Rescue Therapy for Sepsis-Related Severe Thrombocytopenia in Critically Ill Patients. *Cytokine Vol. 136*. Pp 118-127