



DAFTAR PUSTAKA

- Adebawale, K.O. and Lawal, O.S. 2004. Comparative study of the functional properties of bambara groundnut (*Voandzeia subterranean*) jack bean (*Canavalia ensiformis*) and mucuna bean (*Mucuna pruriens*) flours. *Food Res. Int.* 37: 355–365.
- Ahmad Nadzri, F.N., Tawalbeh, D., and Sarbon, N.M. 2021. Physicochemical properties and antioxidant activity of enzymatic hydrolysed chickpea (*Cicer arietinum* L.) protein as influence by alcalase and papain enzyme. *Biocatal. Agric. Biotechnol.*, 36: 102-131.
- Ahmadifard, N., Murueta, J.H.C., Abedian-Kenari, A., Motamedzadegan, A., and Jamali, H. 2016. Comparison the effect of three commercial enzymes for enzymatic hydrolysis of two substrates (rice bran protein concentrate and soy-bean protein) with SDS-PAGE. *J. Food Sci. Technol* 53: 1279–1284.
- Akbarbaglu, Z., Ayaseh, A., Ghanbarzadeh, B., and Sarabandi, K.. 2022. Techno-functional , biological and structural properties of *Spirulina platensis* peptides from different proteases. *Algal Res* 102: 70-75.
- Akpossan, R., Digbeu, Y., Koffi, M., Kouadio, J., Dué, E., and Kouamé, P. 2015. Protein Fractions and Functional Properties of Dried *Imbrasia oyemensis* Larvae Full-Fat and Defatted Flours. *Int. J. Biochem. Res. Rev.* 5: 116–126.
- Al-Ruwaih, N., Ahmed, J., Mulla, M.F., and Arfat, Y.A. 2019. High-pressure assisted enzymatic proteolysis of kidney beans protein isolates and characterization of hydrolysates by functional, structural, rheological and antioxidant properties. *Lwt* 100: 231–236.
- Aluko, R.E. 2015. Amino acids, peptides, and proteins as antioxidants for food preservation. Elsevier Ltd.
- Amarowicz, R. 2008. Antioxidant activity of protein hydrolysates, *Eur. J. Lipid Sci. Technol.* 110: 489–490.
- Amri, E. and Mamboya, F. 2012. Papain, a plant enzyme of biological importance: A review. *Am. J. Biochem. Biotechnol.* 8: 99–104.
- Anwar, F., Latif, S., Przybylski, R., Sultana, B., and Ashraf, M. 2007. Chemical composition and antioxidant activity of seeds of different cultivars of mungbean. *J. Food Sci.* 72: 503–510.
- AOAC. 2005. Official Methods of Analysis of the Association of Official Agricultural Chemists. Cold Spring Harbor, New York.
- Atta, E.M., Mohamed, N.H., and Abdelgawad, A.A.M. 2017. Antioxidants: an



Overview on the Natural and Synthetic Types, Eur. Chem. Bull. 6: 365.

- Di Bernardini, R., Mullen, A.M., Bolton, D., Kerry, J., O'Neill, E., and Hayes, M. 2012. Assessment of the angiotensin-I-converting enzyme (ACE-I) inhibitory and antioxidant activities of hydrolysates of bovine brisket sarcoplasmic proteins produced by papain and characterisation of associated bioactive peptidic fractions. Meat Sci., 90: 226–235.
- Brishti, F.H., Zarei, M., Muhammad, S.K.S., Ismail-Fitry, M.R., Shukri, R., and Saari, N. 2017. Evaluation of the functional properties of mung bean protein isolate for development of textured vegetable protein, Int. Food Res. J., 24: 1595–1605.
- Carrasco-Castilla, J., Hernández-Álvarez, A.J., Jiménez-Martínez, C., Jacinto-Hernández, C., Alaiz, M., Girón-Calle, J., Vioque, J., and Dávila-Ortiz, G. 2012. Antioxidant and metal chelating activities of peptide fractions from phaseolin and bean protein hydrolysates. Food Chem. 135: 1789–1795.
- Chalamaiah, M., Dinesh Kumar, B., Hemalatha, R., and Jyothirmayi, T. 2012. Fish protein hydrolysates: Proximate composition, amino acid composition, antioxidant activities and applications: A review, Food Chem. 135: 3020–3038.
- Chang, Y.W. and Alli, I. 2012. In silico assessment: Suggested homology of chickpea (*Cicer arietinum L.*) legumin and prediction of ACE-inhibitory peptides from chickpea proteins using BLAST and BIOPEP analyses, Food Res. Int. 49: 477–486.
- CHEN, J. sheng, TIAN, J. chun, DENG, Z. ying, ZHANG, Y. xiang, FENG, S. li, YAN, Z. chen, ZHANG, X. ye, and YUAN, H. qing. 2012. Effects of Papain Hydrolysis on the Pasting Properties of Wheat Flour J. Integr. Agric., 11: 1948–1957.
- Chen, Y., Chen, Jianchu, Chang, C., Chen, Juan, Cao, F., Zhao, J., Zheng, Y., and Zhu, J. 2019. Physicochemical and functional properties of proteins extracted from three microalgal species, Food Hydrocoll. 96: 510–517.
- Coda, R., Rizzello, C.G., Pinto, D., and Gobbetti, M. 2012. Selected lactic acid bacteria synthesize antioxidant peptides during sourdough fermentation of cereal flours, Appl. Environ. Microbiol., 78: 1087–1096.
- Daliri, H., Ahmadi, R., Pezeshki, A., Hamishehkar, H., Mohammadi, M., Beyrami, H., Khakbaz Heshmati, M., and Ghorbani, M. 2021. Quinoa bioactive protein hydrolysate produced by pancreatin enzyme- functional and antioxidant properties. Lwt 150: 111-118.
- Du, M., Xie, J., Gong, B., Xu, X., Tang, W., Li, X., Li, C., and Xie, M. 2018. Extraction, physicochemical characteristics and functional properties of Mung bean protein. Food Hydrocoll., 76: 131–140.



- Durand, E., Beaubier, S., Ilic, I., Kapel, R., and Villeneuve, P. 2021. Current Research in Food Science Production and antioxidant capacity of bioactive peptides from plant biomass to counteract lipid oxidation. 4: 365–397.
- Esfandi, R., Walters, M.E., and Tsopmo, A. 2019. Antioxidant properties and potential mechanisms of hydrolyzed proteins and peptides from cereals. *Heliyon*, 5:15-20.
- Esfandi, R., Willmore, W.G., and Tsopmo, A. 2019. Antioxidant and Anti-Apoptotic Properties of Oat. 1–12.
- Fallah-Delavar, M. and Farmani, J. 2018. Recovery and Characterization of Enzymatic Protein Hydrolysates and Fat from Chicken Skin. *JAOCs, J. Am. Oil Chem. Soc.*, 95: 1151–1161.
- Fan, H., Liu, H., Zhang, Y., Zhang, S., Liu, T., and Wang, D. 2022. Review on plant-derived bioactive peptides: biological activities, mechanism of action and utilizations in food development. *J. Futur. Foods*, 2: 143–159.
- Fathollahy, I., Farmani, J., Kasaai, M.R., and Hamishehkar, H. 2021, Characteristics and functional properties of Persian lime (*Citrus latifolia*) seed protein isolate and enzymatic hydrolysates. *Lwt* 140: 110765.
- Felix, M., Cermeño, M., Romero, A., and FitzGerald, R.J. 2019. Characterisation of the bioactive properties and microstructure of chickpea protein-based oil in water emulsions. *Food Res. Int* 121: 577–585.
- Fennema. 2007. Food Chemistry. New York. Taylor and Francis.
- García-Mora, P., Martín-Martínez, M., Angeles Bonache, M., González-Múñiz, R., Peñas, E., Frias, J., and Martinez-Villaluenga, C. 2017. Identification, functional gastrointestinal stability and molecular docking studies of lentil peptides with dual antioxidant and angiotensin I converting enzyme inhibitory activities. *Food Chem.* 221: 464–472.
- Gbogouri, G.A., Linder, M., Fanni, J., and Parmentier, M. 2004. Food Chemistry and Toxicology of Medium-chain Triacylglycerols and, *J. Food Sci* 69: C615–C622.
- Grzonka Z, Kasprzykowski F, Wiczk. 2007. Cysteine Proteases. Di dalam: Polaina J, MacCabe AP, editor. Industrial Enzymes: Structure, Function and Application. Netherlands: Springer.
- Guimarães, R. de C.A., Favaro, S.P., Viana, A.C.A., Braga Neto, J.A., Neves, V.A., and Honer, M.R. 2012. Study of Protein in defatted flour and protein concentrate of baru nuts (*Dipteryx alata* Vog), *Cienc. e Tecnol. Aliment* 32: 464–470.
- Gupta, S., Kapoor, P., Chaudhary, K., Gautam, A., Kumar, R., and Raghava, G.P.S. 2013. In Silico Approach for Predicting Toxicity of Peptides and Proteins. *PLoS One*, 8: 1-10.



- Hernández-Ledesma, B., Miralles, B., Amigo, L., Ramos, M., and Recio, I. 2005. Identification of antioxidant and ACE-inhibitory peptides in fermented milk. *J. Sci. Food Agric* 85: 1041–1048.
- Hossain, F., Yea, S., and Muhammad, K. 2020. Effects of drying techniques on the physicochemical , functional , thermal , structural and rheological properties of mung bean (*Vigna radiata*) protein isolate powder. *138:* .
- HOYLE, N.T. and MERRITT, J.H. 1994. Quality of Fish Protein Hydrolysates from Herring (*Clupea harengus*) *J. Food Sci.* 59. 76–79.
- Huang, C., Tang, X., Liu, Z., Huang, W., and Ye, Y. 2022. Enzymes-dependent antioxidant activity of sweet apricot kernel protein hydrolysates. *LWT*. 154: 112-825.
- Humiski, L.M. and Aluko, R.E. 2007. Physicochemical and bitterness properties of enzymatic pea protein hydrolysates, *J. Food Sci.* 72: .
- Iwaniak, A., Minkiewicz, P., Darewicz, M., and Hrynkiewicz, M. 2016. Food protein-originating peptides as tastants - Physiological, technological, sensory, and bioinformatic approaches. *Food Res. Int.* 89: 27–38.
- Iwaniak, A., Minkiewicz, P., Darewicz, M., Sieniawski, K., and Starowicz, P. 2016. BIOPEP database of sensory peptides and amino acids, *Food Res. Int.*, 85: 155–161.
- Jideani, V.A. 2012. Soybean - Biochemistry, Chemistry and Physiology, *Soybean - Biochem. Chem. Physiol.*,.
- J. Kastin, A. and Pan, W., 2010, Concepts for Biologically Active Peptides, *Curr. Pharm. Des.*, 16, 3390–3400.
- Kandasamy, G., Karuppiah, S.K., and Rao, P.V.S. 2012. Salt- and pH-induced functional changes in protein concentrate of edible green seaweed Enteromorpha species, *Fish. Sci* 78: 169–176.
- Karami, Z., Peighambardoust, S.H., Hesari, J., Akbari-Adergani, B., and Andreu, D. 2019. Antioxidant, anticancer and ACE-inhibitory activities of bioactive peptides from wheat germ protein hydrolysates, *Food Biosci* 32: 100450.
- Kim, S.K. and Wijesekara, I. 2010. Development and biological activities of marine-derived bioactive peptides: A review, *J. Funct. Foods* 2: 1–9.
- Kou, X., Gao, J., Zhang, Z., Wang, H., and Wang, X. 2013. Purification and identification of antioxidant peptides from chickpea (*Cicer arietinum L.*) albumin hydrolysates. *Lwt*, 50: 591–598.
- Kudre, T.G., Benjakul, S., and Kishimura, H. 2013. Comparative study on chemical



- compositions and properties of protein isolates from mung bean, black bean and bambara groundnut/ *J. Sci. Food Agric.* 93: 2429–2436.
- Kusumah, J., Real Hernandez, L.M., and de Mejia, E.G. 2020. Antioxidant potential of mung bean (*Vigna radiata*) albumin peptides produced by enzymatic hydrolysis analyzed by biochemical and in silico methods. *Foods*, 9: 1–15.
- Lafarga, T. and Hayes, M., 2014, Bioactive peptides from meat muscle and by-products: Generation, functionality and application as functional ingredients, *Meat Sci.*, 98, 227–239.
- Laemmli, U.K., 1970, Cleavage of structural proteins during the assembly of the head of bacteriophage, *Nature* 227: 680–685.
- Li, G.H., Wan, J.Z., Le, G.W., and Shi, Y.H. 2006. Novel angiotensin I-converting enzyme inhibitory peptides isolated from Alcalase hydrolysate of mung bean protein, *J. Pept. Sci* 12: 509–514.
- Li, M., Zhang, Y., Xia, S., and Ding, X. 2019. Finding and isolation of novel peptides with anti-proliferation ability of hepatocellular carcinoma cells from mung bean protein hydrolysates, *J. Funct. Foods*, 62: 103557.
- Liu, C., Ren, D., Li, J., Fang, L., Wang, J., Liu, J., and Min, W. 2018. Cytoprotective effect and purification of novel antioxidant peptides from hazelnut (*C. heterophylla* Fisch) protein hydrolysates. *J. Funct. Foods*, 42: 203–215.
- López-Pedrouso, M., Borrajo, P., Pateiro, M., Lorenzo, J.M., and Franco, D. 2020. Antioxidant activity and peptidomic analysis of porcine liver hydrolysates using alcalase, bromelain, flavourzyme and papain enzymes. *Food Res. Int.*, 137: 109389.
- Lu, X., Zhang, L., Sun, Q., Song, G., and Huang, J. 2019. Extraction, identification and structure-activity relationship of antioxidant peptides from sesame (*Sesamum indicum* L.) protein hydrolysate. *Food Res. Int.* 116: 707–716.
- Mas-capdevila, A. 2022. Impact of centrifugation and vacuum filtration step on the yield and molecular weight distribution of protein hydrolysates from rapeseed and sunflower meals 165, 1-10 .
- Minkiewicz, P., Dziuba, J., and Michalska, J. 2011. Bovine meat proteins as potential precursors of biologically active peptides - A computational study based on the BIOPEP database, *Food Sci. Technol. Int.* 17: 39–45.
- Mokni Ghribi, A., Maklouf Gafsi, I., Sila, A., Blecker, C., Danthine, S., Attia, H., Bougatef, A., and Besbes, S. 2015. Effects of enzymatic hydrolysis on conformational and functional properties of chickpea protein isolate, *Food Chem.*, 187: 322–330.



- Mooney, C., Haslam, N.J., Pollastri, G., and Shields, D.C. 2012. Towards the Improved Discovery and Design of Functional Peptides: Common Features of Diverse Classes Permit Generalized Prediction of Bioactivity, PLoS One, 7: 1 -10 .
- Mora, L., Aristoy, M.C., and Toldrá, F. 2018. Bioactive peptides, Encycl. Food Chem 381–389.
- Mosafa, L., Moghadam, M., and Shahedi, M. 2013. Papain enzyme supported on magnetic nanoparticles: Preparation, characterization and application in the fruit juice clarification. Cuihua Xuebao/Chinese J. Catal 34: 1897–1904.
- Moure, A., Sineiro, J., Domínguez, H., and Parajó, J.C. 2006. Functionality of oilseed protein products: A review, Food Res. Int., 39: 945–963.
- Nazeer, R.A., Sampath Kumar, N.S., and Jai Ganesh, R. 2012. In vitro and in vivo studies on the antioxidant activity of fish peptide isolated from the croaker (*Otolithes ruber*) muscle protein hydrolysate. Peptides 35: 261–268.
- Ninan, G., Jose, J., and Abubacker, Z. 2011. Preparation and characterization of gelatin extracted from the skins of rohu (*Labeo rohita*) and common carp (*Cyprinus carpio*). J. Food Process. Preserv 35: 143–162.
- Nwachukwu, I.D. and Aluko, R.E. 2019. Structural and functional properties of food protein-derived antioxidant peptides. J. Food Biochem 43: 1–13.
- Ohata, M., Uchida, S., Zhou, L., and Arihara, K. 2016. Antioxidant activity of fermented meat sauce and isolation of an associated antioxidant peptide. Food Chem. 194: 1034–1039.
- Ortíz Cruz, R.A., Cárdenas López, J.L., González Aguilar, G.A., Astiazarán García, H., Gorinstein, S., Canett Romero, R., and Robles Sánchez, M. 2015. Influence of Sorghum Kafirin on Serum Lipid Profile and Antioxidant Activity in Hyperlipidemic Rats (In Vitro and in Vivo Studies). Biomed Res. Int. 20:15-20.
- Oyaizu, M. 1986. Studies on products of browning reaction. Antioxidative activities of products of browning reaction prepared from glucosamine. Japanese J. Nutr. Diet 44: 307–315.
- Picot, L., Ravallec, R., Martine, F.P., Vandanon, L., Jaouen, P., Chaplain-Derouiniot, M., Guérard, F., Chabeaud, A., Legal, Y., Alvarez, O.M., Bergé, J.P., Piot, J.M., Batista, I., Pires, C., Thorkelsson, G., Delannoy, C., Jakobsen, G., Johansson, I., and Bourseau, P. 2010. Impact of ultrafiltration and nanofiltration of an industrial fish protein hydrolysate on its bioactive properties. J. Sci. Food Agric 90: 1819–1826.
- Pizzino, G., Irrera, N., Cucinotta, M., Pallio, G., Mannino, F., Arcoraci, V., Squadrito, F., Altavilla, D., and Bitto, A. 2017. Oxidative Stress: Harms and Benefits for



Human Health. Oxid. Med. Cell. Longev 20: 17-27.

- Pooja, K., Rani, S., and Prakash, B. 2017. In silico approaches towards the exploration of rice bran proteins-derived angiotensin-I-converting enzyme inhibitory peptides, Int. J. Food Prop 20: 2178–2191.
- Ramos-Lopez, A., Mojica, L., Gomez-Ojeda, A., Macias-Cervantes, M., and Luevano-Contreras, C. 2020. Acute Effect of Black Bean (*Phaseolus vulgaris L.*) Hydrolyzed Protein on Glucose Levels in Adults with Prediabetes and Normal Glucose Tolerance. Curr. Dev. Nutr 4: 458–458.
- Roy, M., Sarker, A., Azad, M.A.K., Shaheb, M.R., and Hoque, M.M. 2020. Evaluation of antioxidant and antimicrobial properties of dark red kidney bean (*Phaseolus vulgaris*) protein hydrolysates, J. Food Meas Charact. 14: 303–313.
- Rui, X., Boye, J.I., Simpson, B.K., and Prasher, S.O. 2013. Purification and characterization of angiotensin I-converting enzyme inhibitory peptides of small red bean (*Phaseolus vulgaris*) hydrolysates, J. Funct. Foods 5: 1116–1124.
- Rutherford, S.M. 2010. Food Composition and Additives: Methodology for Determining Degree of Hydrolysis of Proteins in Hydrolysates: A Review, J. AOAC Int 93: 1–8.
- Ryder, K., Bekhit, A.E.D., McConnell, M., and Carne, A. 2016. Towards generation of bioactive peptides from meat industry waste proteins: Generation of peptides using commercial microbial proteases. Food Chem 208: 42–50.
- Saito, K., Jin, D.H., Ogawa, T., Muramoto, K., Hatakeyama, E., Yasuhara, T., and Nokihara, K. 2003. Antioxidative properties of tripeptide libraries prepared by the combinatorial chemistry. J. Agric. Food Chem 51: 3668–3674.
- Sarmadi, B.H. and Ismail, A. 2010. Antioxidative peptides from food proteins: A review. Peptides 31: 1949–1956.
- Seena, S. and Sridhar, K.R. 2005. Physicochemical, functional and cooking properties of under explored legumes, Canavalia of the southwest coast of India, Food Res. Int. 38: 803–814.
- Shahidi, F., Han, X.Q., and Synowiecki, J. 1995. Production and characteristics of protein hydrolysates from capelin (*Mallotus villosus*). Food Chem 53: 285–293.
- Shalaby, E.A. and Shanab, S.M.M. 2013. Comparison of DPPH and ABTS assays for determining antioxidant potential of water and methanol extracts of *Spirulina platensis*. Indian J. Mar. Sci. 42: 556–564.
- Shen, G., Chahal, B., Majumder, K., You, S.J., and Wu, J. 2010. Identification of novel antioxidative peptides derived from a thermolytic Hydrolysate of ovotransferrin by LC-MS/MS. J. Agric. Food Chem 58: 7664–7672.



- Shevkani, K., Singh, N., Kaur, A., and Rana, J.C. 2015. Structural and functional characterization of kidney bean and field pea protein isolates: A comparative study. *Food Hydrocoll* 43: 679–689.
- Singh, B.P., Bhushan, B., and Vij, S. 2020. Antioxidative, ACE inhibitory and antibacterial activities of soy milk fermented by indigenous strains of lactobacilli, *Legum. Sci.* 2: 1–7.
- Sonklin, C., Laohakunjit, N., and Kerdchoechuen, O. 2011. Physicochemical and flavor characteristics of flavoring agent from mungbean protein hydrolyzed by bromelain. *J. Agric. Food Chem.*, 59, 8475–8483.
- Suetsuna, K., Ukeda, H., and Ochi, H. 2000. Isolation and characterization of free radical scavenging activities peptides derived from casein, *J. Nutr. Biochem* 11: 128–131.
- Suresh Kumar, K., Ganesan, K., Selvaraj, K., and Subba Rao, P. V. 2014. Studies on the functional properties of protein concentrate of *Kappaphycus alvarezii* (Doty) Doty - An edible seaweed. *Food Chem* 153: 353–360.
- Tang, C.H. and Sun, X. 2010. Physicochemical and structural properties of 8S and/or 11S globulins from mungbean [*Vigna radiata* (L.) Wilczek] with various polypeptide constituents. *J. Agric. Food Chem* 58: 6395–6402.
- Tang, T., Du, H., Tang, S., Jiang, Y., Tu, Y., Hu, M., and Xu, M. 2021. Effects of incorporating different kinds of peptides on the foaming properties of egg white powder. *Innov. Food Sci. Emerg. Technol.* 72: 102742.
- Tavano, O.L. 2013. Protein hydrolysis using proteases: An important tool for food biotechnology, *J. Mol. Catal. B Enzym.* 90: 1–11.
- Tsumura, K., Saito, T., Tsuge, K., Ashida, H., Kugimiya, W., and Inouye, K. 2005. Functional properties of soy protein hydrolysates obtained by selective proteolysis, *Lwt*, 38: 255–261.
- Udenigwe, C.C., Udechukwu, M.C., Yiridoe, C., Gibson, A., and Gong, M. 2016. Antioxidant mechanism of potato protein hydrolysates against in vitro oxidation of reduced glutathione. *J. Funct. Foods* 20:, 195–203.
- Valdez-Ortiz, A., Fuentes-Gutiérrez, C.I., Germán-Báez, L.J., Gutiérrez-Dorado, R., and Medina-Godoy, S. 2012. Protein hydrolysates obtained from Azufrado (Sulphur yellow) beans (*Phaseolus vulgaris*): Nutritional, ACE-inhibitory and antioxidative characterization. *LWT - Food Sci. Technol.* 46: 91–96.
- Wahyuningtyas, W.S, Santoso U, Wagiman F.X, Ningrum, A., 2022. Functional properties of Oryctes rhinoceros larvae protein hydrolysates affected by different enzyme concentration and hydrolysis time, *Trends food Sci. Technol.*, 7: 120–



125.

- Waters Corporation. 2012. Acquity UPLC H-class and H-class bio amino acid analysis, Acquity UPLC H-class H-class bio Amin. acid Anal. 4–9.
- Wen, C., Zhang, J., Zhang, H., Duan, Y., and Ma, H. 2020. Plant protein-derived antioxidant peptides: Isolation, identification, mechanism of action and application in food systems: A review, Trends Food Sci. Technol. 105: 308–322.
- Wijatniko, B.D. and Murdiati, A. 2019. Antioxidant activity of bioactive peptides derived from the hydrolysates of jack bean (*Canavalia ensiformis* (L.) DC.) protein isolate, AIP Conf. Proc. 2099.
- Xie, J., Du, M., Shen, M., Wu, T., and Lin, L. 2019. Physico-chemical properties, antioxidant activities and angiotensin-I converting enzyme inhibitory of protein hydrolysates from Mung bean (*Vigna radiata*), Food Chem. 270: 243–250.
- Xie, J., Ye, H., Du, M., Yu, Q., Chen, Y., and Shen, M. 2020. Mung bean protein hydrolysates protect mouse liver cell line Nctc-1469 cell from hydrogen peroxide-induced cell injury. Foods, 9.
- Xu, X.P., Liu, H., Tian, L., Dong, X.B., Shen, S.H., and Qu, L.Q. 2015. Integrated and comparative proteomics of high-oil and high-protein soybean seeds. Food Chem. 172: 105–116.
- Yang, Juanjuan, Hu, L., Cai, T., Chen, Q., Ma, Q., Yang, Jie, Meng, C., and Hong, J. 2018. Purification and identification of two novel antioxidant peptides from perilla (*Perilla frutescens* L. Britton) seed protein hydrolysates, PLoS One. 13: 1–15.
- Yao, Y., Yang, X., Tian, J., Liu, C., Cheng, X., and Ren, G. 2013. Antioxidant and antidiabetic activities of black mung bean (*Vigna radiata* L.). J. Agric. Food Chem., 61: 8104–8109.
- Yi-Shen, Z., Shuai, S., and Fitzgerald, R. 2018. Mung bean proteins and peptides: Nutritional, functional and bioactive properties, Food Nutr. Res. 62: 1–11.
- You, L., Zhao, M., Liu, R.H., and Regenstein, J.M., 2011 Antioxidant and antiproliferative activities of loach (*Misgurnus anguillicaudatus*) peptides prepared by papain digestion, J. Agric. Food Chem. 59: 7948–7953.
- Yüctepe, A., Saroğlu, Ö., and Özçelik, B. 2019. Response surface optimization of ultrasound-assisted protein extraction from *Spirulina platensis*: investigation of the effect of extraction conditions on techno-functional properties of protein concentrates, J. Food Sci. Technol. 56: 3282–3292.
- Zang, X., Yue, C., Wang, Y., Shao, M., and Yu, G. 2019. Effect of limited enzymatic hydrolysis on the structure and emulsifying properties of rice bran protein. J.



Cereal Sci 85: 168–174.

Zayas, 1997. Functionality of Proteins in Food.

Zhang, L., Li, J., and Zhou, K. 2010. Chelating and radical scavenging activities of soy protein hydrolysates prepared from microbial proteases and their effect on meat lipid peroxidation, Bioresour. Technol. 101. 2084–2089.

Zhang, L., Song, C., Chang, J., Wang, Z., and Meng, X. 2022. Optimization of protein hydrolysates production from defatted peanut meal based on physicochemical characteristics and sensory analysis. Lwt. 163: 113572.

Zhang, L.Y., Liu, S., Zhao, X.J., Wang, N., Jiang, X., Xin, H.S., and Zhang, Y.G. 2019. Lactobacillus rhamnosus GG modulates gastrointestinal absorption, excretion patterns, and toxicity in Holstein calves fed a single dose of aflatoxin B 1, J. Dairy Sci. 102: 1330–1340.

Zhang, Q., Tong, X., Qi, B., Wang, Z., Li, Y., Sui, X., and Jiang, L., 2018. Changes in antioxidant activity of Alcalase-hydrolyzed soybean hydrolysate under simulated gastrointestinal digestion and transepithelial transport, J. Funct. Foods. 42: 298–305.

Zhang, Y., Ding, X., and Li, M.. 2021. Preparation, characterization and in vitro stability of iron-chelating peptides from mung beans. Food Chem 349: 129101.

Zheng, L., Yu, H., Wei, H., Xing, Q., Zou, Y., Zhou, Y., and Peng, J.. 2018. Antioxidative peptides of hydrolysate prepared from fish skin gelatin using ginger protease activate antioxidant response element-mediated gene transcription in IPEC-J2 cells. J. Funct. Foods 51: 104–112.

Zulvana, A.H., Andriati, N., Sri, A., and Widiastuti, S. 2019. In silico approach in evaluation of jack bean (*Canavalia ensiformis*) canavalin protein as precursors of bioactive peptides with dual antioxidant and angiotensin i-converting enzyme inhibitor, Mater. Sci. Forum. 948: 85–94.