

DAFTAR PUSTAKA

- Andrei, C., Margiocco, P., Poggi, A., Lotti, L.V., Torrisi, M.R., Rubartelli, A. (2004). Phospholipases C and A2 control lysosome-mediated IL-1 β secretion: Implications for inflammatory processes. *PNAS*, 101(26): 9745–9750.
- Anonim. (2019). *Mitochondrial DNA Isolation Kit, for the rapid isolation of Mitochondrial DNA in various cell and tissue samples*. Abcam®.
- Ali Abdelhamid, Y., Kar, P., Finnis, M.E., Phillips, L.K., Plummer, M.K., Shaw, J.E., *et al.* (2016). Stress hyperglycaemia in critically ill patients and the subsequent risk of diabetes: a systematic review and meta-analysis. *Critical Care*, 20:301.
- Al-Kafaji, G.A.L., Aljadaan, A., Kamal, A., Bakhiet, M. (2018). Peripheral blood mitochondrial DNA copy number as a novel potential biomarker for diabetic nephropathy in type 2 diabetes patients. *Exp Ther Med*, 16:1483–1492.
- American Diabetes Association (ADA). (2019). Classification and Diagnosis of Diabetes: Standards of Medical Care in Diabetes. *Diabetes Care*, 42(1):S13-S15.
- Amorim, R.G., Guedes, G.S., Vasconcelos, S.M.L., Santos, J.C.F. (2019). Kidney Disease in Diabetes Melitus: Cross-Linking between Hyperglycemia, Redox Imbalance and Inflammation. *Arq Bras Cardiol*, 112(5):577-597.
- Animaw, W., Seyoum, Y. (2017). Increasing prevalence of diabetes melitus in a developing country and its related factors. *PLOS ONE*, 7(11):1–11.
- Bakar, M.H.A., Hairunisa, N., & Huri, H.Z. (2018). Reduced mitochondrial DNA content in lymphocytes is associated with insulin resistance and infammation in patients with impaired fasting glucose. *Clin Exp Med*, 18:373-382.
- Bae, J.H., II, S.J., Kim, S.J., Lee, J.M., Jeong, J.H., Kang, J.S., *et al.* (2019). Circulating cell-free mtDNA contributes to AIM2 inflammasome-mediated chronic inflammation in patients with type 2 diabetes. *cells*, 8(328):1-12.
- Baynest, H.W. (2015). Classification, pathophysiology, diagnosis and management of diabetes melitus. *J Diabetes Metab*, 6(5):1-9.
- Bhat, A., Koul, A., Rai, E., Sharma, S., Dhar, M.K., Bamezai, R.N.K. (2007). PGC-1 α Thr394Thr and Gly482Ser variants are significantly associated with T2DM in two North Indian populations: a replicate case-control study. *Hum Genet*, 121: 609-614.

- Biosource. (2020). *Human IL-1 Beta ELISA Kit PicoKine™*. PicoKine™ ELISA catalog number: MBS175901.
- Biosource. (2020). Human Toll Like Receptor 9 (TLR9) ELISA Kit. Catalog No: MBS4502948.
- Bio-Rad. (2006). *Real-Time PCR Applications Guide*. Bio-Rad Laboratories.
- Blaslov K, Naranda F.S, Kruljac I, Renar I.P. (2018). Treatment approach to type 2 diabetes: Past, present and future. *World J Diabetes*, 9(12): 209-219.
- Boyapati, R.K., Tamborska, A., Dorward, D.A., & Ho, G. (2019). Advances in the understanding of mitochondrial DNA as a pathogenic factor in inflammatory diseases. *F1000Research*, 6(169):1–15.
- BPJS Kesehatan Cabang Surakarta. (2022). *Utilisation Review (UR) RS Universitas Sebelas Maret Periode Januari-November 2021*. Surakarta: Kantor BPJS Cabang Surakarta.
- Canizales, Y.G.C., Rivera, E.E.U., Jacobo, R.E.G., Perez, D.P.P., Yadira, B., Rivera, J.G.R. (2018). Increased levels of AIM2 and circulating mitochondrial DNA in type 2 diabetes. *Iran J Immunol*, 15(2):142-155.
- Cao, Z., & Cooper, M. E. (2011). Pathogenesis of diabetic nephropathy. Mini Review. *Journal of Diabetes Investigation*, 2(4):243–247.
- Chambers, J.M & Wingert, R.A. (2020). PGC-1 α in Disease: Recent Renal Insights into a Versatile Metabolic Regulator. Review. *Cells*, 9, 2234.
- Chou, Su-Tze & Tseng, Shih-Ting. (2017). Oxidative stress markers in type 2 diabetes patients with diabetic nephropathy. *Clin Exp Nephrol*, 21:283–292.
- Csep, K., Szigeti, E., Vitai, M., & Koranyi, L. (2017). The PPARGC1a - Gly482ser polymorphism (rs8192678) and the metabolic syndrome in a central romanian population. *Acta Endocrinol*, 13(2):161–167.
- Dai, Z., Jiao, Y., Fan, Q., Qi, A., Xiao, L., & Li, J. (2020). Homocysteine, interleukin-1 β , and fasting blood glucose levels as prognostic markers for diabetes melitus complicated with cerebral infarction and correlated with carotid intima-media thickness. *Exp Ther Med*, 19:1167–1174.
- Decroli, E. (2019). *Diabetes Melitus Tipe 2*. Edisi Pertama. Padang: Pusat Penerbitan Bagian Ilmu Penyakit Dalam Fakultas Kedokteran Universitas Andalas.

- Delanaye, P., Cavalier, E., Pottel, H. (2017). Serum Creatinine: Not So Simple!. *Nephron*, 136:302–308.
- Ding, S., Xu, S., Ma, Y., Liu, G., Jang, H., Fang, J. (2019). Modulatory Mechanisms of the NLRP3 Inflammasomes in Diabetes. *Biomolecules*, 9,850.
- Donate-correa, J., Luis-Rodriguez, D., Martin-Nunez, E., Tagua, V.G., Hernandez-Carballo, C., Ferri, C., *et al.* (2020). Inflammatory targets in diabetic nephropathy. *J. Clin. Med.* 458(9):1-23.
- Eynon, N., Meckel, Y., Sagiv, M., Yamin, C., Amir, R., Sagiv, M., *et al.* (2010). Do PPAR α and PPAR γ polymorphisms influence sprint or endurance phenotypes?. *Scand. J. Med. Sci. Sports*, 20:e145–e150.
- Eisele, P.S., Furrer, R., Beer, M., & Handschin, C. (2015). The PGC-1 coactivators promote an anti-inflammatory environment in skeletal muscle in vivo. *Biochem Biophys Res Commun*, 464(3): 692–697.
- Elmarakby, A.A. & Sullivan, J.C. (2012). Relationship between oxidative stress and inflammatory cytokines in diabetic nephropathy. *Cardiovasc Ther*, 30: 49–59.
- Filograna, R., Mennuni, M., Alsina, D., Larsson, N-G. (2020). Mitochondrial DNA copy number in human disease: the more the better?. Review Article. *FEBS Letters*, 595:976-1002.
- Fontecha-Barriuso, M., Martin-Sanchez, D., Martinez-Moreno, J.M., Monsalve, M., Ramoz, A.M., Sanchez-Nino, M.D., *et al.* (2020). The role of PGC-1 α and mitochondrial biogenesis in kidney diseases. *biomolecules*, 10(347): 1–24.
- Fukuda, D., Nishimoto, S., Aini, K., Tanaka, A., Nishiguchi, T., Kim-Kaneyama, J., *et al.* (2019). Toll-Like Receptor 9 Plays a Pivotal Role in Angiotensin II–Induced Atherosclerosis. *J Am Assoc*, 8: e010860.
- Gambardella, S., Limanaqi, F., Ferese, R., Biagioni, F., Campopiano, R., Centonze, D., *et al.* (2019). ccf-mtDNA as a potential link between the brain and immune system in neuro-immunological disorders. *frontiers in Immunology*, 10:1-9.
- García-garcía, P.M., Getino-melián, M.A., Domínguez-pimentel, V., & Navarro-gonzález, J.F. (2014). Inflammation in diabetic kidney disease. *World J Diabetes*, 5(4): 431–443.

- Gianotti, T.F., Sookoian, S., Dieuzeide, G., García, S.I., Gemma, C., González, C.D., *et al.* (2008). A decreased mitochondrial DNA content is related to insulin resistance in adolescents. *OBESITY*, 16:1591–1595.
- Gilkerson, R. (2017). Commentary: mitochondrial DNA damage and loss in diabetes. *Diabetes Metab Res Rev.* 32(7):672–674.
- Goh, K. & Sum, C. (2010). Connecting the dots: molecular and epigenetic mechanisms in type 2 diabetes. *Current Diabetes Review*, 6: 255–265.
- Gonzalez-hunt, C.P., Rooney, J.P., Ryde, I.T., Anbalagan, C., Joglekar, R., & Meyer, J.N. (2016.) PCR-based analysis of mitochondrial DNA copy number, mitochondrial DNA damage, and nuclear DNA damage. *Curr. Protoc. Toxicol.* 67:20.11.1-20.11.25.
- Goracy, I., Kaczmarczyk, M., Ciechanowicz, A., Lewandowska, K., Jakubiszyn, P., Bodnar, O. (2019). Polymorphism of interleukin 1 β may modulate the risk of ischemic stroke in Polish patients. *Medicina*, 55(558):1-11.
- Haneda, M., Utsunomiya, K., Koya, D., Babazono, T., Moriya, T., Makino, H *et al.* (2015). A new Classification of Diabetic Nephropathy 2014: a report from Joint Committee on Diabetic Nephropathy. *J Diabetes Invest*, 6:242-246.
- Inci, A., Olmaz, R., Sarı, F., Coban, M., Hy, E., Sarıkaya, M. (2016). Increased oxidative stress in diabetic nephropathy and its relationship with soluble Klotho levels. *HIPPOKRATIA*, 20(3):198–203.
- International Diabetes Federation. (2021). *IDF Diabetes Atlas*, 10th Edition. ISBN: 978-2-930229-98-0.
- Jin, J., Ding, G., Bao, H., Chen, Y., Han, Y., Zhao, F., *et al.* (2013). Correlation between PPAR gene polymorphisms and primary nephrotic syndrome in children. *PPAR Research*, 2013:1-8.
- Kaku, K. (2010). Pathophysiology of type 2 diabetes and its treatment policy. *JMAJ*, 53(1): 41–46.
- Kaneko, N., Kurata, M., Yamamoto, T., Morikawa, S., Masumoto, J. (2019). *The role of interleukin-1 in general pathology*. Inflammation and Regeneration, 39(12):1-16
- Kanungo, S., Morton, J., Neelakantan, M., Ching, K., Saeedian, J., Goldstein, A. (2018). Mitochondrial disorders. *Rev. Artic. Ann Transl Med*, 6(24):1-17.

- Kementerian Kesehatan RI. (2019). *Laporan provinsi Jawa Tengah Riskesdas 2018*. Lembaga Penerbit Badan Penelitian dan Pengembangan Kesehatan.
- Kementerian Kesehatan RI. (2018). *Hasil Utama Riskesdas 2018*. Jakarta : Badan Penelitian dan Pengembangan Kesehatan.
- Knez, D., Winckelmans, E., Plusquin, M., Thijs, L., Cauwenberghs, N., Gu, Y. (2015). Correlates of peripheral blood mitochondrial DNA content in a general population. *Am J Epidemiol*, 183(2):138-146.
- Kraja, A.T., Liu, C., Fetterman, J.L., Graff, M., Have, C.T., Gu, C., *et al.* (2019). Associations of mitochondrial and nuclear mitochondrial variants and genes with seven metabolic traits. *Am J Hum Genet*, 3(104): 112–138.
- Kruzliak, P., Haley, A.P., Starcevic, J.N., Gaspar, L., Petrovic, D. (2015). Polymorphisms of the Peroxisome Proliferator-Activated Receptor- γ (rs1801282) and its coactivator-1 (rs8192673) are associated with obesity indexes in subjects with type 2 diabetes melitus. *Cardiovascular Diabetology*, 14:42.
- Kusdiyah, E., Makmur, M.J., Aras, RBP. (2020). Karakteristik faktor-faktor risiko terjadinya komplikasi kronik nefropati diabetik dan atau penyakit pembuluh darah perifer pada penderita diabetes mellitus di rsud raden mattaher tahun 2018. *e-SEHAD*, 1(1):19-32.
- Lavoz, C., Rodrigues-diez, R.R., Plaza, A., Carpio, D., Egido, J., Ruiz-ortega, M., *et al.* (2020). VEGFR2 blockade improves renal damage in an experimental model of type 2 diabetic nephropathy. *J. Clin. Med.* 9(302):1-21.
- Lee, G., Uddin, J., Kim, Y., Ko, M., Yu, I., Ha, H. (2019). PGC -1 α , a potential therapeutic target against kidney aging. *Aging Cell*, 18:1–20.
- Lee, H.K., Song, J.H., Shin, C.S., Park, D.J., Park, K.S., Lee, K.U., *et al.* (1998). Decreased mitochondrial DNA content in peripheral blood precedes the development of non-insulin-dependent diabetes melitus. *Diabetes Res and Clin Pract*, 42: 161–167.
- Léveillé, M., Besse-Patin, A., Jouvet, N., Gunes, A., Sczelecki, S., Jeromson, S., *et al.* (2020). PGC-1 α isoforms coordinate to balance hepatic metabolism and apoptosis in inflammatory environments. *Mol Metab*, 34:72–84.
- Li, Z., Zhu, M., Du, J., Ma, H., Jin, G., Dai, J. (2018). Genetic variants in nuclear DNA along with environmental factors modify mitochondrial DNA copy number: A population-based exome-wide association study. *BMC Genomics*, 19 (752):1–9.

- Lin, Y-C., Chang, Y-H., Yang, S-Y., Wu, K-D., Chu, T-S. (2018). *Update of pathophysiology and management of diabetic kidney disease*. Journal of The Formosan Medical Association, 117:662-675.
- Lindinger, A., Peterli, R., Peters, T., Kern, B., Flüe, M., Calame, M. (2010). Mitochondrial DNA Content in Human Omental Adipose Tissue. *OBES SURG*, 20:84–92.
- Liu, C. & Lin, J.D. (2011). PGC-1 coactivators in the control of energy metabolism Structure and function of PGC-1. *Acta Biochim Biophys Sin*, 43:248–257.
- Liu, H., Li, Y., Lin, N., Dong, X., Li, W., Deng, Y., Ma, L. (2020). Interleukin-1 β Promotes Ox-LDL Uptake by Human Glomerular Mesangial Cells via LOX-1. *Int J. Med. Sci.*, 17(8): 1056-1061.
- Liu, M., Peng, J., Tai, N., Pearson, J.A., Hu, C., Guo, J., *et al.* (2018). Toll-like receptor 9 negatively regulates pancreatic islet beta cell growth and function in a mouse model of type 1 diabetes. *Diabetologia*, 61: 2333–2343.
- Lopez-Castejon, G and Brough, D. (2011). Understanding the mechanism of IL-1 β secretion. *Cytokine & Growth Factor Review*, 22:189-195.
- Lucia, A., Gómez-Gallego, F., Barroso, I., Rabadán, M., Bandrés, F., San Juan, A. F., *et al.* (2005). PPARGC1A genotype (Gly482Ser) predicts exceptional endurance capacity in European men. *J. Appl. Physiol.*, 99:344–348.
- Malik, A.N., Shahni, R., Iqbal, M.M. (2009). Increased peripheral blood mitochondrial DNA in type 2 diabetic patients with nephropathy. *Diabetes Research And Clinical Practice*, 86:e22– e24.
- Margaryan, S., Kriegova, E., Fillerova, R., Kraiczova, V.S., Manukyan, G. (2020). Hypomethylation of IL1RN and NFKB1 genes is linked to the dysbalance in IL1 β /IL-1Ra axis in female patients with type 2 diabetes mellitus. *PloS ONE*, 15(5): e0233737.
- Masser, D.R., Otalora, L., Clark, N.W., Kinter, M.T., Elliott, H., Freeman, W.M. (2018). Functional changes in the neural retina occur in the absence of mitochondrial dysfunction in a rodent model of diabetic retinopathy. *J Neurochem*, 143(5): 595–608.
- Meyer, A., Laverny, G., Bernardi, L., Charles, A.L., Alsaleh, G., Pottecher, J., *et al.* (2018). Mitochondria: An organelle of bacterial origin controlling inflammation. *Front Immunol*, 9(538): 1–8.

- Misra, A., Gopalan, H., Jayawardena, R., Hills, AP., Soares, M., Rezza-Albarran, AA., *et al.* (2019). Diabetes in developing countries. *J Diabetes*, 11(7):522-539.
- Molli, A.E.I., Bergonzi, M.F., Spalvieri, M.P., Linari, M.A., Frechtel, G.D., Cerrone, G.E. (2020). Relationship between the IL-1 β serum concentration, mRNA levels and rs16944 genotype in the hyperglycemic normalization of T2D patients. *Scientific Reports*, 10:9985.
- National Center for Biotechnology Information. *PPARGC1A* gene. Available at <http://www.ncbi.nlm.nih.gov> (diakses tanggal 16 Desember 2022).
- Nishimoto, S., Fukuda, D., & Sata, M. (2020). Emerging roles of Toll-like receptor 9 in cardiometabolic disorders. *Inflammation and Regeneration*, 40(18):1-13.
- Ohashi, N., Morino, K., Ida, S., Sekine, O., Lemecha, M., Kume, S., *et al.* (2017). Pivotal role of O -GlcNAc modification in cold-induced thermogenesis by brown adipose tissue through mitochondrial biogenesis. *Diabetes*, (66):2351–2362.
- Pan, H., Zhang, L., Guo, M., Sui, H., Li, H., Wu, W. (2010). The oxidative stress status in diabetes melitus and diabetic nephropathy. *Acta Diabetol*, 4(Suppl 1): 71-76.
- Papadopoulou-Marketou, N., Kanaka-Gantenbein, C., Marketos, N., Chrousos, G.P., Papassotiriou, I. (2017). Biomarkers of diabetic nephropathy : A 2017 update. *Crit Rev Lab Sci*, 54:326–342.
- Pasquinelly, A., Chico, L., Pasquali, L., Bisordi, C., Gerfo, A., Fabbrini, M. *et al.* (2016). Gly482Ser PGC-1 α Gene Polymorphism and Exercise-Related Oxidative Stress in Amyotrophic Lateral Sclerosis Patients. *Front. Cell. Neurosci*, 10:102
- Pavkov, ME., Collins, AJ., Coresh, J., Nelson, RG. (2015). *Diabetes in America*, 3rd Edition. Chapter 22 Kidney Disease in Diabetes.
- Peiró, C., Lorenzo, óscar, Carraro, R., & Sánchez-Ferrer, C.F. (2017). IL-1 β inhibition in cardiovascular complications associated to diabetes melitus. *Front Pharm*, 8(363):1–13.
- Perkumpulan Endokrinologi Indonesia (PERKENI). (2021). *Pedoman Pengelolaan dan Pencegahan Diabetes Melitus Tipe 2 di Indonesia 2021*. PB PERKENI, ISBN: 978-602-53035-5-5.

- Perkumpulan Endokrinologi Indonesia (PERKENI). (2019). *Pedoman Pengelolaan dan Pencegahan Diabetes Melitus Tipe 2 Dewasa di Indonesia 2019*. PB PERKENI, ISBN: 978-602-53035-2-4.
- Perkumpulan Endokrinologi Indonesia (PERKENI). (2015). *Konsensus Pengelolaan Dan Pencegahan Diabetes Melitus Tipe 2 Di Indonesia 2015*. PB PERKENI, ISBN: 978-979-19388-6-0.
- Peterson, S.B., & Hart, G.W., 2016. New insights : A role for O -GlcNAcylation in diabetic complications. *Crit. Rev. Biochem. Mol. Biol*, 51(3):150–161.
- Prior, S.L., Clark, A.R., Jones, D.A., Bain, S.C., Hurel, S.J., Humphries, S.E., *et al.* (2012). Association of the PGC-1 α rs8192678 variant with microalbuminuria in subjects with type 2 diabetes melitus. *Disease Markers*, 32:363–369.
- Pusat Data dan Informasi Kementerian Kesehatan RI (Infodatin). (2020). *Tetap produktif, cegah, dan atasi diabetes melitus*, ISSN 2442-7659. Jakarta Selatan: Kementerian Kesehatan RI Pusat Data dan Informasi.
- Qiagen. (2016). QIAamp® DNA Mini and Blood Mini Handbook. QIAGEN Sample and Assay Technologies.
- Ruscitti, P., Ursini, F., Cipriani, P., Greco, M., Alvaro, S., Vasiliki, L., *et al.* (2019). IL-1 inhibition improves insulin resistance and adipokines in rheumatoid arthritis patients with comorbid type 2 diabetes. *Medicine*, 98(7):1-9.
- Saini, D.C., Kochar, A., Poonia, R. (2021). Clinical correlation of diabetic retinopathy with nephropathy and neuropathy. *Indian J Ophthalmol*, 69:3364-8.
- Samsu, N. (2021). Diabetic Nephropathy: Challenges in Pathogenesis, Diagnosis, and Treatment. *Biomed Research International*, Vol. 2021:1-17.
- Saremi, L., Lotfipannah, S., Mohammadi, M., Hosseinzadeh, H., Hosseini-Khah, Z., Johari, B., Saltanatpour, Z. (2019). Association between PPARGC1A single nucleotide polymorphisms and increased risk of nonalcoholic fatty liver disease among Iranian patients with type 2 diabetes melitus. *Turk J Med Sci*, 49:1089-1094.
- Satitapoj, B., and Adler, SG. (2015). *Prevalence and Management of Diabetic Nephropathy in Western Countries*, *Kidney Dis*, 1:61-70.
- Scarpulla, R.C. (2011). Metabolic control of mitochondrial biogenesis through the PGC-1 family regulatory network. *Biochim Biophys Acta* 1813(7):1269–1278.

- Schmidth, AM. (2018). Highlighting Diabetes Mellitus The Epidemic Continues. *Arterioscler Thromb Vasc*, 38:e1-e8.
- Schwartz, S.S., Eipstein, S., Corkey, B.E., Grant, S.F.A., Gavin III, J.R., Aguilar, R.B. (2016). The time is right for a new classification system for diabetes: rationale and implications of the β -cell-centric classification schema. *Diabetes Care*, 39:176-186.
- Scurt, F.G., Menne, J., Brandt, S., Bernhardt, A., Mertens, P.R., Haller, H., *et al.* (2019). Systemic inflammation precedes microalbuminuria in diabetes. *Kidney Int Rep*, 4:1373–1386.
- Shoar, Z., Goldenthal, M.J., Luca, F. De, & Suarez, E. (2016). Mitochondrial DNA content and function, childhood obesity, and insulin resistance. *Endocr Res*, 41(1):49–56.
- Shokouhi, S., Haghani, K., Borji, P., Bakhtiyari, S. (2015). Association Between PGC-1 α Gene Polymorphisms and Type 2 Diabetes Risk: A Case-Control Study of an Iranian Population. *Can J Diabetes*, 39:65-72.
- Siew, E,D and Matheny, M.E. (2015). Choice of Reference Serum Creatinine in Defining AKI. *Nephron*, 131(2):107–112.
- Sivitz, W.I. & Yorek, M.A. (2010). Mitochondrial dysfunction in diabetes: from molecular mechanism to functional significance and therapeutic opportunities. *Antioxid Redox Signal*, 12(4):537-577.
- Skyler, J.S., Bakris, G.L., Bonifacio, E., Darsow, T., Eckel, R.H., Groop, L., *et al.* (2017). Differentiation of diabetes by pathophysiology, natural history, and prognosis. *Diabetes*, 66:241–255.
- Song, J; Oh, JY; Sung, Y; Pak, YK; Park, KS; Lee, H. (2001). Peripheral blood mitochondrial DNA content is related to insulin sensitivity in offspring of type 2 diabetic patients. *Diabetes Care*, 24(5): 865–870.
- Soyal, S., Krempler, F., Oberkofler, H., Patsch, W. (2006). PGC-1 α : a potent transcriptional cofactor involved in the pathogenesis of type 2 diabetes. *Diabetologia*, 49: 1477–1488.
- Tang, Z., Zeng, F., & Zhang, X. (2015). Human genetics of diabetic nephropathy. *Ren. Fail*, 37(3): 363–371.

- Testa, R., Bonfigli, A.R., Prattichizzo, F., Sala, L.L., Nigris, V.D., Ceriello, A. (2017). The “Metabolic Memory” Theory and the Early Treatment of Hyperglycemia in Prevention of Diabetic Complications. *Nutrients*, 9(437):1-9.
- Thipsawat, S. (2021). Early detection of diabetic nephropathy in patient with type 2 diabetes mellitus: A review of the literature. *Diabetes & Vascular Disease Research*, 18(6):1-9.
- Tobina, T., Mori, Y., Doi, Y., Nakayama, F., Kiyonaga, A., Tanaka, H. (2017). Peroxisome proliferator-activated receptor gamma co-activator 1 gene Gly482Ser polymorphism is associated with the response of low-density lipoprotein cholesterol concentrations to exercise training in elderly Japanese. *J. Physiol. Sci.* 67: 595–602.
- Tomino, Y. and Gohda, T. (2015). *The Prevalence and Management of Diabetic Nephropathy in Asia*. *Kidney Dis*, 1:52-60.
- Umanath, K., & Lewis, J. B. (2018). Update on diabetic nephropathy: core curriculum 2018. *AJKD*, 71(6):884–895.
- Vandenbeek, R., Khan, N.P., & Estall, J.L. (2018). Linking metabolic disease with the PGC-1 α Gly482Ser polymorphism. *Endocrinology* 159(2):853–865.
- Varshney, P., Parveen, R., Khan, M.A., Kohli, S., Agarwal, N.B. (2019). Increased serum interleukin-9 and interleukin-1 β are associated with depression in type 2 diabetes patients. *Arq Neuropsiquiatr*, 78(5):255-261.
- Wang, W., Sun, W., Cheng, Y., Xu, Z., Cai, L. (2019). Role of sirtuin-1 in diabetic nephropathy. *J Mol Med*, 97:291–309.
- Wang, Y., Xu, W., Zhu, D., Zhang, N., Wang, Y., Ding, M., *et al.* (2018). Specific expression network analysis of diabetic nephropathy kidney tissue revealed key methylated sites. *J Cell Physiol*, 233(10): 7139–7147.
- Weng, S-W., Lin, T-K., Liou, C-W., Chen, S-D., Wei, Y-H., Lee, H-C. (2008). Peripheral blood mitochondrial DNA content and dysregulation of glucose metabolism. *Diabetes Research and Clinical Practice*, 83:94–99.
- Wifi, M.N.A., Assem, M., Elsherif, R.H., El-Azab, H.A.F., Saif, A. (2017). Toll-like receptors-2 and -9 (TLR2 and TLR9) gene polymorphism in patients with type 2 diabetes and diabetic foot. *Medicine*, 96:17.

World Health Organization. (2019). *Classification of diabetes melitus*. Geneva: Licence: CC BY-NC-SA 3.0 IGO. ISBN 978-92-4-151570-2.

World Health Organization. (2016). *Global report of diabetes*. Geneva: WHO Library Cataloguing-in-Publication Data, available on <http://www.who.int>.

World Health Organization & International Diabetes Federation. (2006). *Definition and Diagnosis of Diabetes Melitus and Intermediate Hyperglycemia*. Geneva: WHO Document Production Services, ISBN: 92 4 159493 4.

Wu, H., Deng, X., Shi, Y., Su, Y., Wei, J., Duan, H. (2016). PGC-1 α , glucose metabolism and type 2 diabetes melitus. *J Endocrinol*, 229(3): 99-115.

Xia, W., Chen, N., Peng, W., Jia, X., Yu, Y., Wu, X., *et al.* (2019). Systematic meta-analysis revealed an association of PGC-1 α rs8192678 polymorphism in type 2 diabetes melitus. *Dis Markers*, 2019(2970401):1-10.

Xu, F.X., Zhou, X., Shen, F., Pang, R., Liu, S.M. (2012). Decreased peripheral blood mitochondrial DNA content is related to HbA 1c, fasting plasma glucose level and age of onset in type 2 diabetes melitus. *Diabet Met*, 29(7):47–54.

Yuan, Y., Fernandes, V., Newsholme, P., Cheng, J., Chen, Y., Lu, Y. (2016). Regulation of SIRT1 in aging: roles in mitochondrial function and biogenesis. *Mech Ageing Dev*, 155:10–21.

Zhang, S., Lu, W., Yan, Li., Wu, M., Xu, M., *et al.* (2007). Association between peroxisome proliferator-activated receptor- γ coactivator-1 α gene polymorphisms and type 2 diabetes in southern Chinese population: role of altered interaction with myocyte enhancer factor 2C. *Chin Med J*, 120(21): 1878-1885.



UNIVERSITAS
GADJAH MADA

Potensi PGC-1 α ; Gly482Ser, DNA Mitokondria Copy Number (mtDNA-CN), Toll-like Receptor 9

(TLR9), dan Interleukin-1 α ; (IL-1 α) sebagai Biomarker Prediktif Diabetes Melitus Tipe 2 dengan Nefropati

RIANDINI AISYAH, dr. Ahmad Hamim Sadewa, PhD; dr. Suryono Yudha Patria, Sp.A(K), PhD

Universitas Gadjah Mada, 2023 | Diunduh dari <http://etd.repository.ugm.ac.id/>