



UNIVERSITAS  
GADJAH MADA

INDUKSI DAN PERKEMBANGAN MIKROSPORA EMBRIOGENIK PADA PADI HITAM TRANSGENIK  
(*Oryza sativa L.*)  
& Ireng & Iren  
IRENE KSPRESI OsRKD3  
MERI HANDAYANI, Dr. Yekti Asih Purwestri, S.Si., M.Si; Dr. Tri Rini Nuringtyas, S.Si., M.Sc  
Universitas Gadjah Mada, 2023 | Diunduh dari <http://etd.repository.ugm.ac.id/>

## DAFTAR PUSTAKA

- Bajpai, R., & Chaturvedi, R. (2018). *Haploid Embryogenesis in Tea Haploid Embryogenesis in Tea* (Issue May). Springer International Publishing. <https://doi.org/10.1007/978-3-319-79087-9>
- Bhojwani, S. S., & Dantu, P. K. (2013). Plant tissue culture: An introductory text. In *Plant Tissue Culture: An Introductory Text*. <https://doi.org/10.1007/978-81-322-1026-9>
- Boutilier, K., Offringa, R., Sharma, V. K., Kieft, H., Ouellet, T., Zhang, L., Hattori, J., Liu, C., Lammeren, A. A. M. Van, Miki, B. L. A., Custers, J. B. M., & Lookeren, M. M. Van. (2002). *Ectopic Expression of BABY BOOM Triggers a Conversion from Vegetative to Embryonic Growth*. 14(August), 1737–1749. <https://doi.org/10.1105/tpc.001941.tissue>
- Budiman, B., Arisoesiloningsih, E., & Bagus Edi Wibowo, R. (2012). Growth Adaptation of Two Indonesian Black Rice Origin NTT Cultivating In Organic Paddy Field, Malang-East Java. *Journal of Tropical Life Science*, 2(3), 77–80. <https://doi.org/10.11594/jtls.02.03.04>
- Chang, T., & Bardenas, E. A. (1965). The Morphology and Varietal Characteristics of The Rice Plant. *Irri*, 40. [https://www.google.co.uk/books/edition/Morphology\\_and\\_Varietal\\_Characteristics/xoR0r5Nam9QC?hl=en&gbpv=0](https://www.google.co.uk/books/edition/Morphology_and_Varietal_Characteristics/xoR0r5Nam9QC?hl=en&gbpv=0)
- Chardin, C., Girin, T., Roudier, F., Meyer, C., & Krapp, A. (2014). The plant RWP-RK transcription factors: Key regulators of nitrogen responses and of gametophyte development. *Journal of Experimental Botany*, 65(19), 5577–5587. <https://doi.org/10.1093/jxb/eru261>
- Chaudhry, B., & Rustagi, A. (2015). *Anther*. University of Delhi.
- Cho, M. S., & Zapata, F. J. (1990). *Plant Regeneration from Isolated Microspore of Indica Rice*. 31(6), 881–885.
- Cordewener, J. H. G., Hause, G., Grge, E., Busink, R., Hause, B., Dons, H. J. M., Lammeren, A. A. M. Van, Campagne, M. M. V. L., & Pechan, P. (1995). *Changes in synthesis and localization of members of the 70-kDa class of heat-shock proteins accompany the induction of embryogenesis in Brassica napus L. microspores*. 31, 747–755.
- Custers, J. B. M., Cordewener, J. H. G., Fiers, M. A., Maassen, B. T. H., Van Lookeren



UNIVERSITAS  
GADJAH MADA

INDUKSI DAN PERKEMBANGAN MIKROSPORA EMBRIOGENIK PADA PADI HITAM TRANSGENIK  
(*Oryza sativa L.*)  
& Ireng & Cempo  
MERI HANDAYANI, Dr. Yekti Asih Purwestri, S.Si., M.Si; Dr. Tri Rini Nuringtyas, S.Si., M.Sc  
Universitas Gadjah Mada, 2023 | Diunduh dari <http://etd.repository.ugm.ac.id/>

- Campagne, M. M., & Liu, C. M. (2001). Androgenesis in Brassica. *Current Trends in the Embryology of Angiosperms*, 451–470. [https://doi.org/10.1007/978-94-017-1203-3\\_18](https://doi.org/10.1007/978-94-017-1203-3_18)
- Custers, Jan B.M., Cordewener, J. H. G., Nöllen, Y., Dons, H. J. M., & Van Lockeren Campagne, M. M. (1994). Temperature controls both gametophytic and sporophytic development in microspore cultures of *Brassica napus*. *Plant Cell Reports*, 13(5), 267–271. <https://doi.org/10.1007/BF00233317>
- Datta, S. K., Datta, K., & Potrykus, I. (1990). Embryogenesis and plant regeneration from microspores of both “Indica” and “Japonica” rice (*Oryza sativa*). *Plant Science*, 67, 83–88. [https://doi.org/10.1016/0168-9452\(90\)90053-q](https://doi.org/10.1016/0168-9452(90)90053-q)
- Dewi, I. S., & Purwoko, B. S. (2016). Kultur Antera untuk Percepatan Perakitan Varietas Padi di Indonesia. *Jurnal AgroBiogen*, 8(2), 78. <https://doi.org/10.21082/jbio.v8n2.2012.p78-88>
- Dewi, I. S., Purwoko, B. S., & Aswidinnoor, H. (2004). Kultur antera padi pada beberapa formulasi media yang mengandung poliamin. *Jurnal Bioteknologi Pertanian*, 9(1), 14–19.
- Dubas, E., Castillo, A. M., Żur, I., Krzewska, M., & Vallés, M. P. (2021). Microtubule organization changes severely after mannitol and n-butanol treatments inducing microspore embryogenesis in bread wheat. *BMC Plant Biology*, 21(1), 1–19. <https://doi.org/10.1186/s12870-021-03345-3>
- Fauzia, A. N. (2017). *Overekspresi Gen OsRKD4 pada Padi Hitam (*Oryza sativa L.* ‘Cempo Ireng’) untuk Induksi Embriogenesis Somatik*. Universitas Gadjah Mada.
- Feng, K., Hou, X., Xing, G., Liu, J., Duan, A., Xu, Z., Li, M., Zhuang, J., & Xiong, A. (2020). Critical Reviews in Biotechnology Advances in AP2 / ERF superfamily transcription factors in plant. *Critical Reviews in Biotechnology*, 0(0), 1–27. <https://doi.org/10.1080/07388551.2020.1768509>
- Germanà, M. A. (2011). Anther culture for haploid and doubled haploid production. *Plant Cell, Tissue and Organ Culture*, 104(3), 283–300. <https://doi.org/10.1007/s11240-010-9852-z>
- Goldberg, R. B., Beals, T. P., & Sanders, P. M. (1993). *Anther Development : Basic Principles and Practical Applications*. 5(October), 1217–1229.
- Guo-Wei, F., & Hai-Men, L. (1985). Studies on function of cold pretreatment affecting the efficiency of rice anther culture. *Acta Phytophysiologica Sin*, 11, 366–380.



Haddadi, P., Medicines, H., Karimzadeh, G., & Abdollahi, M. R. (2008). *Effects of gibberellin , abscisic acid and embryo desiccation on normal plantlet regeneration , secondary embryogenesis and callogenesis in microspore Effects of gibberellin , abscisic acid and embryo desiccation on normal plantlet regeneration , secondar*. April.

Halbritter, H., Ulrich, S., Grímsson, F., Weber, M., Zetter, R., Hesse, M., Buchner, R., Svojtka, M., & Frosch-Radivo, A. (2018). Illustrated Pollen Terminology. In *Illustrated Pollen Terminology*. <https://doi.org/10.1007/978-3-319-71365-6>

Hoekstra, S., van Zijderveld, M. H., Louwerse, J. D., Heidekamp, F., & van der Mark, F. (1992). Anther and microspore culture of *Hordeum vulgare* L. cv. Igri. *Plant Science*, 86(1), 89–96. [https://doi.org/10.1016/0168-9452\(92\)90182-L](https://doi.org/10.1016/0168-9452(92)90182-L)

Honya, D., & Twell, D. (2004). Transcriptome analysis of haploid male gametophyte development in *Arabidopsis*. *Genome Biology*, 5(1). <https://doi.org/https://doi.org/10.1186/gb-2004-5-11-r85>

Hu, T.-X., Yu, M., & Zhao, J. (2010). Comparative transcriptional profiling analysis of the two daughter cells from tobacco zygote reveals the transcriptome differences in the apical and basal cells. *BMC Plant Biology*, 10, 167. <https://doi.org/10.1186/1471-2229-10-167>

Huang, B., & Keller, W. A. (1989). Microspore culture technology. *Journal of Tissue Culture Methods*, 12(4), 171–178. <https://doi.org/10.1007/BF01404445>

Huang, M. Der, Wei, F. J., Wu, C. C., Hsing, Y. I. C., & Huang, A. H. C. (2009). Analyses of advanced rice anther transcriptomes reveal global tapetum secretory functions and potential proteins for lipid exine formation. *Plant Physiology*, 149(2), 694–707. <https://doi.org/10.1104/pp.108.131128>

Indrianto, A., Barinova, I., Touraev, A., & Heberle-bors, E. (2001). Tracking individual wheat microspores *in vitro*: identification of embryogenic microspores and body axis formation in the embryo. *Planta*, 212(2), 163–174. <https://doi.org/10.1007/s004250000375>

Indrianto, A., Heberle-bors, E., & Touraev, A. (1999). Assessment of various stresses and carbohydrates for their effect on the induction of embryogenesis in isolated wheat microspores. 143, 71–79.

Islam, M., Haque, E., & Alam, S. M. (2013). Morphological and Histological Observation of Embryogenic Calli Derived from Immature Embryo of BRRI Dhan28 (*Oryza sativa L.*) Variety. 3(5), 21–27.



UNIVERSITAS  
GADJAH MADA

INDUKSI DAN PERKEMBANGAN MIKROSPORA EMBRIOGENIK PADA PADI HITAM TRANSGENIK  
(*Oryza sativa L.*)  
Ireng & Cempo  
MERI HANDAYANI, Dr. Yekti Asih Purwestri, S.Si., M.Si; Dr. Tri Rini Nuringtyas, S.Si., M.Sc  
Universitas Gadjah Mada, 2023 | Diunduh dari <http://etd.repository.ugm.ac.id/>

- Jacquard, C., Mazeyrat-Gourbeyre, F., Devaux, P., Boutilier, K., Baillieul, F., & Clement, C. (2009). Microspore embryogenesis in barley: anther pre-treatment stimulates plant defence gene expression. *Planta*, 229, 393–402. <https://doi.org/10.1007/s00425-008-0838-6>
- Jang, S., & Li, H. Y. (2018). Overexpression of *OsAP2* and *OSWRKY24* in arabidopsis results in reduction of plant size. *Plant Biotechnology*, 35(3), 273–279. <https://doi.org/10.5511/plantbiotechnology.18.0508a>
- Jung, K. H., Han, M. J., Lee, Y. S., Kim, Y. W., Hwang, I., Kim, M. J., Kim, Y. K., Nahm, B. H., & An, G. (2005). Rice Undeveloped Tapetum1 is a major regulator of early tapetum development. *Plant Cell*, 17(10), 2705–2722. <https://doi.org/10.1105/tpc.105.034090>
- Kasha, K. J., Simion, E., Oro, R., Yao, Q. A., Hu, T. C., & Carlson, A. R. (2001). An improved *in vitro* technique for isolated microspore culture of barley. 1994, 379–385.
- Kim, Y. J., Jang, M. G., Zhu, L., Silva, J., Zhu, X., Sukweenadhi, J., Kwon, W. S., Yang, D. C., & Zhang, D. (2016). Cytological characterization of anther development in Panax ginseng Meyer. *Protoplasma*, 253(4), 1111–1124. <https://doi.org/10.1007/s00709-015-0869-3>
- Kimber, G., & Riley, R. (1963). Haploid Angiosperms. *Botanical Review*, 29(4), 480–531.
- Köszegi, D., Johnston, A. J., Rutten, T., Czihal, A., Altschmied, L., Kumlehn, J., Wüst, S. E. J., Kirioukhova, O., Gheyselinck, J., Grossniklaus, U., & Bäumlein, H. (2011). Members of the RKD transcription factor family induce an egg cell-like gene expression program. *Plant Journal*, 67(2), 280–291. <https://doi.org/10.1111/j.1365-313X.2011.04592.x>
- Kristamtini, K., Sutarno, S., Wiranti, E. W., & Widayanti, S. (2016). Kemajuan Genetik dan Heritabilitas Karakter Agronomi Padi Beras Hitam pada Populasi F2. *Jurnal Penelitian Pertanian Tanaman Pangan*, 35(2), 119. <https://doi.org/10.21082/jpptp.v35n2.2016.p119-124>
- Krogaard, H., & Andersen, A. S. (1983). Free amino acids of Nicotiana alata anthers during development *in vivo*. *Physiologia Plantarum*, 57(4), 527–531. <https://doi.org/10.1111/j.1399-3054.1983.tb02780.x>
- Kumar, A., Batra, R., Gahlaut, V., Gautam, T., Kumar, S., Sharma, M., Tyagi, S., Singh, K. P., Balyan, H. S., Pandey, R., & Gupta, P. K. (2018). Genome-wide identification and characterization of gene family for RWP-RK transcription



UNIVERSITAS  
GADJAH MADA

INDUKSI DAN PERKEMBANGAN MIKROSPORA EMBRIOGENIK PADA PADI HITAM TRANSGENIK  
(*Oryza sativa L.*)  
Ireng & Cempo  
MERI HANDAYANI, Dr. Yekti Asih Purwestri, S.Si., M.Si; Dr. Tri Rini Nuringtyas, S.Si., M.Sc  
Universitas Gadjah Mada, 2023 | Diunduh dari <http://etd.repository.ugm.ac.id/>

factors in wheat (*Triticum aestivum L.*). *PLoS ONE*, 13(12), 1–28. <https://doi.org/10.1371/journal.pone.0208409>

Kumar, V., & Giridhar, P. (2009). *AgNO3 - A potential regulator of ethylene activity and plant growth modulator*. 12(May 2014). <https://doi.org/10.2225/vol12-issue2-fulltext-1>

Kurniawan, A. A. (2016). *Pengaruh auksin endogen terhadap embriogenesis mikrospora padi (*Oryza sativa L. 'Fatmawati*')*. Universitas Gadjah Mada.

Kyo, M., & Harada, H. (1986). Control of the developmental pathway of tobacco pollen *in vitro*. *Planta*, 168(4), 427–432. <https://doi.org/10.1007/BF00392260>

Lestari, A. P. (2008). *Evaluasi Mutu Beras 18 Galur Padi Hasil Kultur Antera*. 1449–1455.

Li, N., Liu, H., Yin, C., Li, X., Liang, W., Yuan, Z., Xu, B., Chu, H., Wang, J., Wen, T., & Huang, H. (2006). *The Rice Tapetum Degeneration Retardation Gene Is Required for Tapetum Degradation and Anther Development*. 18(November), 2999–3014. <https://doi.org/10.1105/tpc.106.044107>

Licausi, F., Ohme-takagi, M., & Perata, P. (2013). Factors : Mediators of Stress Responses and Developmental Programs. *New Phytologist*, 199(3), 639–649.

Liu, J. X., Wang, M., Chen, B. X., Jin, P., Li, J. Y., & Zeng, K. (2012). Microsporogenesis, microgametogenesis, and pollen morphology of Ambrosia artemisiifolia L. in China. *Plant Systematics and Evolution*, 298(1), 43–50. <https://doi.org/10.1007/s00606-011-0521-4>

Liu, L., White, M. J., & Macrae, T. H. (1999). Transcription factors and their genes in higher plants: Functional domains, evolution and regulation. *Eur. j. Biochem*, 262, 247–257.

Ma, Z., Wu, T., Huang, K., Jin, Y. M., Li, Z., Chen, M., Yun, S., Zhang, H., Yang, X., Chen, H., Bai, H., Du, L., Ju, S., Guo, L., Bian, M., Hu, L., Du, X., & Jiang, W. (2020). A Novel AP2/ERF Transcription Factor, OsRPH1, Negatively Regulates Plant Height in Rice. *Frontiers in Plant Science*, 11(May), 1–15. <https://doi.org/10.3389/fpls.2020.00709>

Maraschin, A. S. F., Priester, W. De, Spaink, H. P., Wang, M., Maraschin, S. F., Priester, W. De, Spaink, H. P., & Wang, M. (2005). Androgenic switch : an example of plant embryogene from the male gametophyte perspective. *Journal of Experimental Botany*, 56(417), 1711–1726. <https://doi.org/10.1093/jxb/eri190>



UNIVERSITAS  
GADJAH MADA

INDUKSI DAN PERKEMBANGAN MIKROSPORA EMBRIOGENIK PADA PADI HITAM TRANSGENIK  
(*Oryza sativa L.*)  
& Ireng & Cempo  
IRENG & CEMPO  
MERI HANDAYANI, Dr. Yekti Asih Purwestri, S.Si., M.Si; Dr. Tri Rini Nuringtyas, S.Si., M.Sc  
Universitas Gadjah Mada, 2023 | Diunduh dari <http://etd.repository.ugm.ac.id/>

Maraschin, S. D. F., Gaussand, G., Pulido, A., Olmedilla, A., Lamers, G. E. M., Korthout, H., Spaink, H. P., & Wang, M. (2005). Programmed cell death during the transition from multicellular structures to globular embryos in barley androgenesis. *Planta*, 221(4), 459–470. <https://doi.org/10.1007/s00425-004-1460-x>

Mishra, R. (2016). *In-vitro Androgenesis in Rice : Advantages , Constraints and Future Prospects. March*. <https://doi.org/10.1016/j.rsci.2016.02.001>

Mizoi, J., Shinozaki, K., & Yamaguchi-Shinozaki, K. (2012). AP2/ERF family transcription factors in plant abiotic stress responses. *Biochimica et Biophysica Acta - Gene Regulatory Mechanisms*, 1819(2), 86–96. <https://doi.org/10.1016/j.bbagr.2011.08.004>

Müller, M., & Munné-bosch, S. (2015). *Ethylene Response Factors : A Key Regulatory Hub in Hormone and Stress Signaling 1*. 169(September), 32–41. <https://doi.org/10.1104/pp.15.00677>

Mustari, E., Diningrat, D. S., Ratnashih, R., & Widiyanto, S. M. (2016). APETALA2 and APETALA3 genes expression profiling on floral development of teak (*tectona grandis* Linn f.). *Journal of Plant Sciences*, 11(4–5), 61–68. <https://doi.org/10.3923/jps.2016.61.68>

Nafisa, A. R. (2021). *Analisis Faktor Transkripsi OsERF3, OsAP2, OsEREBP2 Tanaman Padi Hitam Transgenik (*Oryza sativa L. 'Cempo Ireng'*) Overekspresi OsRKD3*. Universitas Gadjah Mada.

Neves, M., Correia, S., Cavaleiro, C., & Canhoto, J. (2021). Modulation of Organogenesis and Somatic Embryogenesis by Ethylene: An Overview. *Plants*, 10, 1–30.

Nitsch, J. P., & Nitsch, C. (1969). Haploid Plants from Pollen Grains. *Science*, 163(3862), 85–87. <https://doi.org/https://doi.org/10.1126/science.163.3862.85>

Nurbaiti, S., Indrianto, A., Purwestri, Y. .., & Daryono, B. S. (2021). Characterization of Putative RKD Homologous Gene With Microspore Embryogenesis in The Local Pigmented Rice Cultivar ‘Segreng.’ *Sabrao Journal of Breeding and Genetics*, 53(2), 527–534.

Nurbaiti, Siti. (2022). Induksi dan Perkembangan Mikrospora Embriogenik Padi (*Oryza sativa L. 'Segreng'*). Universitas Gadjah Mada.

Nurbaiti, Siti, Purwestri, Y. A., Daryono, B. S., Semiarti, E., & Indrianto, A. (2019). Developmental the pattern of embryogenic microspore of rice (*Oryza sativa L.*)



UNIVERSITAS  
GADJAH MADA

INDUKSI DAN PERKEMBANGAN MIKROSPORA EMBRIOGENIK PADA PADI HITAM TRANSGENIK  
(*Oryza sativa L.*)  
& Ireng & Irleng  
MERI HANDAYANI, Dr. Yekti Asih Purwestri, S.Si., M.Si; Dr. Tri Rini Nuringtyas, S.Si., M.Sc  
Universitas Gadjah Mada, 2023 | Diunduh dari <http://etd.repository.ugm.ac.id/>

based on morphological characteristic. *Journal of Biological Researches*, 25(1), 7–11. <https://doi.org/10.23869/bphjbr.25.1.20192>

Okamuro, J. K., Caster, B., Villarroel, R., Montagu, M. Van, & Jofuku, K. D. (1997). The AP2 domain of APETALA 2 defines a large new family of DNA binding proteins in *Arabidopsis*. *Plant Biology*, 94(June), 7076–7081. <https://doi.org/10.1073/pnas.94.13.7076>

Ouakfaoui, S. El, Schnell, J., Abdeen, A., Colville, A., Labbe, H., Han, S., Baum, B., Laberge, S., & Miki, B. (2010). Control of somatic embryogenesis and embryo development by AP2 transcription factors. *Plant Molecular Biology*, 74, 313–326. <https://doi.org/10.1007/s11103-010-9674-8>

Pauk, J., Puolimatka, M., Lökös Tóth, K., & Monostori, T. (2000). *In vitro* androgenesis of triticale in isolated microspore culture. *Plant Cell, Tissue and Organ Culture*, 61(3), 221–229. <https://doi.org/10.1023/A:1006416116366>

Pechan, P. M., & Keller, W. A. (1988). Identification of potentially embryogenic microspores in *Brassica napus*. *Physiologia Plantarum*, 74(2), 377–384. <https://doi.org/10.1111/j.1399-3054.1988.tb00646.x>

Pérez-Pérez, Y., El-Tantawy, A. A., Solís, M. T., Risueño, M. C., & Testillano, P. S. (2019). Stress-Induced Microspore Embryogenesis Requires Endogenous Auxin Synthesis and Polar Transport in Barley. *Frontiers in Plant Science*, 10(September), 1–16. <https://doi.org/10.3389/fpls.2019.01200>

Piyatrakul, P., Putranto, R., Martin, F., Rio, M., Dessailly, F., Leclercq, J., Dufayard, J., Lardet, L., & Montoro, P. (2012). Some ethylene biosynthesis and AP2 / ERF genes reveal a specific pattern of expression during somatic embryogenesis in *Hevea brasiliensis*.

Pratiwi, R., & Purwestri, Y. A. (2017). Black rice as a functional food in Indonesia. *Functional Foods in Health and Disease*, 7(3), 182–194. <https://doi.org/10.31989/ffhd.v7i3.310>

Rajcan, I., Boersma, J. G., & Shaw, E. J. (2011). Plant Genetic Techniques: Plant Breeder's Toolbox. In *Comprehensive Biotechnology, Second Edition* (Second Edi, Vol. 4). Elsevier B.V. <https://doi.org/10.1016/B978-0-08-088504-9.00252-X>

Sakuma, Y., Liu, Q., Dubouzet, J. G., Abe, H., Yamaguchi-Shinozaki, K., & Shinozaki, K. (2002). DNA-binding specificity of the ERF/AP2 domain of *Arabidopsis* DREBs, transcription factors involved in dehydration- and cold-



inducible gene expression. *Biochemical and Biophysical Research Communications*, 290(3), 998–1009. <https://doi.org/10.1006/bbrc.2001.6299>

Saleh, A., & Pages, M. (2003). Plant AP2/ERF transcription factors. *Genetika*, 35(1), 37–50. <https://doi.org/10.2298/GENS0301037S>

Scagliusi, S. M. (2014). Establishing isolated microspore culture to produce doubled haploid plants in Brazilian wheat (*Triticum aestivum L.*) Sandra Mansur Scagliusi EMBRAPA - Brazilian Agricultural Research Corporation , National Wheat Research Center , Passo Fundo – RS , 9900. *Australian Journal of Crop Science*, 8(6), 887–894. [http://www.cropj.com/scaglius-8\\_6\\_2014\\_887\\_894.pdf](http://www.cropj.com/scaglius-8_6_2014_887_894.pdf)

Segui, M., & Nuez, F. (2008). How microspores transform into haploid embryos : changes associated with embryogenesis induction and microspore- derived embryogenesis. *Physiologia Planta*, 134, 1–12. <https://doi.org/10.1111/j.1399-3054.2008.01113.x>

Serra, T. S., Figueiredo, D. D., Cordeiro, A. M., Almeida, D. M., Lourenço, T., Abreu, I. A., Sebastián, A., Fernandes, L., Contreras-Moreira, B., Oliveira, M. M., & Saibo, N. J. M. (2013). OsRMC, a negative regulator of salt stress response in rice, is regulated by two AP2/ERF transcription factors. *Plant Molecular Biology*, 82(4–5), 439–455. <https://doi.org/10.1007/s11103-013-0073-9>

Sholikhah, U., Parjanto, Handoyo, T., & Yunus, A. (2018). Anthocyanin Content in Some Black Rice Cultivars. In *IOP Conference Series: Materials Science and Engineering* (Vol. 161, Issue 1, pp. 0–5). <https://doi.org/10.1088/1755-1315/709/1/012976>

Soriano, M., Li, H., & Boutilier, K. (2013). Microspore embryogenesis: Establishment of embryo identity and pattern in culture. *Plant Reproduction*, 26(3), 181–196. <https://doi.org/10.1007/s00497-013-0226-7>

Suaib, S., & Arma, M. J. (2012). Pengembangan Kultur Mikrospora pada Varietas Padi Ladang Lokal Asal Kendari Development of Microspore Culture of Local Upland Rice Varieties from Kendari. *J. Agro Indonesia*, 40(2), 99–104.

Sunderland, A. N., & Wicks, F. M. (2022). *Embryoid Formation in Pollen Grains of Nicotiana tabacum* Linked references are available on JSTOR for this article : *Embryoid Formation in Pollen Grains of Nicotiana tabacum*. 22(70), 213–226.

Sunderland, N., Huang, B., & Hills, G. J. (1984). Disposition of pollen in situ and its relevance to anther/pollen culture. *Journal of Experimental Botany*, 35(4), 521–530. <https://doi.org/10.1093/jxb/35.4.521>



UNIVERSITAS  
GADJAH MADA

INDUKSI DAN PERKEMBANGAN MIKROSPORA EMBRIOGENIK PADA PADI HITAM TRANSGENIK  
(*Oryza sativa L.*)  
Ireng & Cempo  
MERI HANDAYANI, Dr. Yekti Asih Purwestri, S.Si., M.Si; Dr. Tri Rini Nuringtyas, S.Si., M.Sc  
Universitas Gadjah Mada, 2023 | Diunduh dari <http://etd.repository.ugm.ac.id/>

- Tang, X., Liu, Y., He, Y., Ma, L., & Sun, M. X. (2013). Exine dehiscing induces rape microspore polarity, which results in different daughter cell fate and fixes the apicalbasal axis of the embryo. *Journal of Experimental Botany*, 64(1), 215–228. <https://doi.org/10.1093/jxb/ers327>
- Tedeschi, F., Rizzo, P., Rutten, T., Altschmied, L., & Bäumlein, H. (2017). RWP-RK domain-containing transcription factors control cell differentiation during female gametophyte development in *Arabidopsis*. *New Phytologist*, 213(4), 1909–1924. <https://doi.org/10.1111/nph.14293>
- Testillano, P. S. (2019). Microspore embryogenesis: targeting the determinant factors of stress-induced cell reprogramming for crop improvement. *Journal of Experimental Botany*, 70(11), 2965–2978. <https://doi.org/10.1093/JXB/ERY464>
- Thanuja, B., & Parimalavalli, R. (2018). Role of Black Rice in Health and Diseases. *International Journal of Health Sciences and Research*, 8(2), 241–248. [https://www.ijhsr.org/IJHSR\\_Vol.8\\_Issue.2\\_Feb2018/IJHSR\\_Abstract.031.htm](https://www.ijhsr.org/IJHSR_Vol.8_Issue.2_Feb2018/IJHSR_Abstract.031.htm)
- Touraev, A., Indrianto, A., Wratschko, I., Vicente, O., & Heberle-Bors, E. (1996). Efficient microspore embryogenesis in wheat (*Triticum aestivum L.*) induced by starvation at high temperature. *Sexual Plant Reproduction*, 9(4), 209–215. <https://doi.org/10.1007/s004970050033>
- Touraev, Alisher, Ilham, A., Vicente, O., & Heberle-Bors, E. (1996). Stress-induced microspore embryogenesis in tobacco: An optimized system for molecular studies. *Plant Cell Reports*, 15(8), 561–565. <https://doi.org/10.1007/BF00232453>
- Touraev, Alisher, Pfosser, M., & Heberle-Bors, E. (2001). The microspore: A haploid multipurpose cell. In *Advances in Botanical Research* (Vol. 35). [https://doi.org/10.1016/s0065-2296\(01\)35004-8](https://doi.org/10.1016/s0065-2296(01)35004-8)
- Touraev, Alisher, Vicente, O., & Heberle-Bors, E. (1997). Initiation of microspore embryogenesis by stress. *Trends in Plant Science*, 2(8), 297–302. [https://doi.org/10.1016/S1360-1385\(97\)01083-2](https://doi.org/10.1016/S1360-1385(97)01083-2)
- Tripathi, D. K. K., Warrier, D. R., Govila, D. O. P., & Ahuja, D. V. (2011). *Biology\_of\_Rice.pdf*. In *Series of Crop Specific Biology Documents* (p. 54).
- Vijay, D., & Roy, B. (2013). Rice (*Oryza sativa L.*). In *Breeding, Biotechnology and Seed Production of Field Crops* (Issue December, pp. 71–122). [https://www.researchgate.net/publication/281152980\\_Rice\\_Oryza\\_sativa\\_L](https://www.researchgate.net/publication/281152980_Rice_Oryza_sativa_L)



UNIVERSITAS  
GADJAH MADA

INDUKSI DAN PERKEMBANGAN MIKROSPORA EMBRIOGENIK PADA PADI HITAM TRANSGENIK  
(*Oryza sativa L.*)  
& Ireng & Irena  
MERI HANDAYANI, Dr. Yekti Asih Purwestri, S.Si., M.Si; Dr. Tri Rini Nuringtyas, S.Si., M.Sc  
Universitas Gadjah Mada, 2023 | Diunduh dari <http://etd.repository.ugm.ac.id/>

Vrinten, P. L., Nakamura, T., & Kasha, K. J. (1999). Characterization of cDNAs expressed in the early stages of microspore embryogenesis in barley (*Hordeum vulgare L.*) *Plant Molecular Biology*, 41, 455–463.

Waki, T., Hiki, T., Watanabe, R., Hashimoto, T., & Nakajima, K. (2011). The arabidopsis RWP-RK protein RKD4 triggers gene expression and pattern formation in early embryogenesis. *Current Biology*, 21(15), 1277–1281. <https://doi.org/10.1016/j.cub.2011.07.001>

Wang, L. S., & Stoner, G. D. (2008). Anthocyanins and their role in cancer prevention. *Cancer Letters*, 269(2), 281–290. <https://doi.org/10.1016/j.canlet.2008.05.020>

Xie, J. H., Gao, M. W., Liang, Z. Q., Shu, Q. Y., Cheng, X. Y., & Xue, Q. Z. (1997). The effect of cool-pretreatment on the isolated microspore culture and the free amino acid change of anthers in japonica rice (*Oryza sativa L.*). *Journal of Plant Physiology*, 151(1), 79–82. [https://doi.org/10.1016/S0176-1617\(97\)80040-5](https://doi.org/10.1016/S0176-1617(97)80040-5)

Xie, Z., Nolan, T. M., Jiang, H., & Yin, Y. (2019). AP2 / ERF Transcription Factor Regulatory Networks in Hormone and Abiotic Stress Responses in Arabidopsis. 10(February), 1–17. <https://doi.org/10.3389/fpls.2019.00228>

Yaish, M. W., El-Kereamy, A., Zhu, T., Beatty, P. H., Good, A. G., Bi, Y. M., & Rothstein, S. J. (2010). The APETALA-2-like transcription factor *OsAP2-39* controls key interactions between abscisic acid and gibberellin in rice. *PLoS Genetics*, 6(9). <https://doi.org/10.1371/journal.pgen.1001098>

Yamuangmorn, S., & Prom-U-thai, C. (2021). The potential of high-anthocyanin purple rice as a functional ingredient in human health. *Antioxidants*, 10(6), 1–21. <https://doi.org/10.3390/antiox10060833>

Zapata, F. ., Aldemita, R. ., Ella, E. ., & Cho, M. S. (1990). Rice Genetics II. In *Proceedings of the Second International Genetics Symposium*.

Zhai, L., Xu, L., Wang, Y., Zhu, X., Feng, H., Li, C., & Luo, X. (2016). Transcriptional identification and characterization of differentially expressed genes associated with embryogenesis in radish (*Raphanus sativus L.*). *Nature Publishing Group*, September 2015, 1–13. <https://doi.org/10.1038/srep21652>

Zhang, H., Zhang, J., Quan, R., Pan, X., Wan, L., & Huang, R. (2013). EAR motif mutation of rice OsERF3 alters the regulation of ethylene biosynthesis and drought tolerance. *Planta*, 237. <https://doi.org/10.1007/s00425-013-1852-x>

Zhang, Z., Lu, A. E. Y., & Liu, A. E. X. (2005). Nuclear and cell migration during pollen development in rice (*Oryza sativa L.*). December 2015.



UNIVERSITAS  
GADJAH MADA

INDUKSI DAN PERKEMBANGAN MIKROSPORA EMBRIOGENIK PADA PADI HITAM TRANSGENIK  
(*Oryza sativa L.*)  
Ireng & Cempo  
MERI HANDAYANI, Dr. Yekti Asih Purwestri, S.Si., M.Si; Dr. Tri Rini Nuringtyas, S.Si., M.Sc  
Universitas Gadjah Mada, 2023 | Diunduh dari <http://etd.repository.ugm.ac.id/>

<https://doi.org/10.1007/s00497-005-0241-4>

Zhao, L., Xu, A. S., & Chai, A. T. (2006). *OsAP2-1 , an AP2 -like gene from Oryza sativa , is required for flower development and male fertility.* 197–206. <https://doi.org/10.1007/s00497-006-0036-2>

Zorinians, S., Tashpulatov, A. S., Heberle-Bors, E., & Touraev, A. (2005). The role of stress in the induction of haploid microspore embryogenesis. *Biotechnology in Agriculture and Forestry*, 56, 35–52. [https://doi.org/10.1007/3-540-26889-8\\_3](https://doi.org/10.1007/3-540-26889-8_3)

Žur, I., Dubas, E., Krzewska, M., Sánchez-Díaz, R. A., Castillo, A. M., & Vallés, M. P. (2014). Changes in gene expression patterns associated with microspore embryogenesis in hexaploid triticale ( $\times$ Triticosecale Wittm.). *Plant Cell, Tissue and Organ Culture*, 116(2), 261–267. <https://doi.org/10.1007/s11240-013-0399-7>