

REFERENCES

- Abdelgader, H., & Heimbach, U. (1992). The effect of some insect growth regulators (IGRs) on first instar larvae of the carabid beetle *Poecilus cupreus* (Coleoptera: Carabidae) using different application methods. *Aspects of Applied Biology*, 31, 171–177.
- Abduchalek, B., Rauf, A., & . P. (2017). Kutu Putih Singkong, *Phenacoccus Manihoti* Matile-Ferrero (Hemiptera: Pseudococcidae): Persebaran Geografi Di Pulau Jawa Dan Rintisan Pengendalian Hayati. *Jurnal Hama Dan Penyakit Tumbuhan Tropika*, 17(1), 1. <https://doi.org/10.23960/j.hptt.1171-8>
- Adriani, E., & Lestari, T. W. W. (2021). Utilization of Natural Enemy (Parasitoid *Anagyrus lopezi*) Introduction from Bogor to Control Pink Cassava Mealybug (*Phenacoccus manihoti*) in Gorontalo. *Jurnal Pertanian Berkelanjutan*, 9(1), 18–26.
- Adriani, E., Rauf, A., & Pudjianto, P. (2016). Laju enkapsulasi parasitoid *Anagyrus lopezi* (De Santis) (Hymenoptera: Encyrtidae) oleh kutu putih singkong *Phenacoccus manihoti* Matile-Ferrero (Hemiptera: Pseudococcidae). *Jurnal Entomologi Indonesia*, 13(3), 147–155. <https://doi.org/10.5994/jei.13.3.147>
- Alves, A. A. C. (2002). Cassava botany and physiology. *Cassava: Biology, Production and Utilization*, 67–89. <https://doi.org/10.1079/9780851995243.0067>
- Anonymus. (1985). *Technical information. Applaud 25 WP on citrus scales and mealybugs.*
- Anonymus. (1987). *Applaud, new pesticide (insect growth regulator), technical information.*
- Bellotti, A. ., Smith, L., & Lapointe, A. (1999). Recent Advances in Cassava Pest Management. *Annual Review of Entomology*, 343–370.
- Bellotti, A. C. (2002). Arthropod pests. In R. J. Hillocks, J. M. Thresh, & A. C. Bellotti (Eds.), *Cassava: Biology, Production and Utilization* (pp. 209–235). Centro Internacionalde Agricultura Tropical (CIAT).
- Bellotti, A. C., Herrera, C. J., Hernández, M. del P., Arias, B., Guerrero, J. M., & Melo, E. L. (2011). Cassava Pests in Latin America, Africa and Asia. In R. H. Howeler (Ed.), *The Cassava Handbook. A Reference Manual based on the Asian Regional Cassava Training Course, held in Thailand.* (pp. 199–257). Centro Internacional de Agricultura Tropical (CIAT).
- Bellotti, A. C., Melo, E. L., Arias, B., Herrera, C. J., Hernández, M. del P., Holguín, C. M., Guerrero, J. M., & Trujillo, H. (2005). Biological control in the neotropics: A selective review with emphasis on cassava. *Second International Symposium on Biological Control of Arthropods; September 12-16, 2005, Davos, Switzerland*, 1(Table 2), 206–227.
- Bellotti, A., Campo, H. V. B., & Hyman, G. (2012). Cassava Production and Pest Management: Present and Potential Threats in a Changing Environment. *Tropical Plant Biology*, 5(1), 39–72. <https://doi.org/10.1007/s12042-011-9091-4>
- Ben-Dov, Y. (1994). A systematic catalogue of the mealybugs of the world (Insecta: Homoptera: Coccoidea: Pseudococcidae and Putoidae) with data on geographical distribution, host plants, biology and economic importance. *Intercept Limited Andover UK*, 686.
- Bertschy, C., Turlings, T. C. J., Bellotti, A. C., & Dorn, S. (2001). The Role of Mealybug-Induced Cassava Plant Volatiles in the Attraction of the Encyrtid Parasitoids *Aenasius vexans* and *Apoanagyrus diversicornis*. *Journal Of Insect Behavior*, 14(3). <https://doi.org/10.3929/ethz-a-010025751>
- Bigler, F., & Waldburger, M. (1994). Effects of pesticides on *Chrysoperla carnea* Steph. (Neuroptera, Chrysopidae) in the laboratory and semi-field. *Bulletin OILB SROP (France)*, 17, 55–69.
- Blank, R. H., Olson, M. H., Clark, J. B., & Gill, G. S. C. (1993). Investigating two bee-safe materials for controlling latania scale on avocados during pollination. *In Proceedings of*

the New Zealand Plant Protection Conference, 46, 80–85.

- Braun, A. R., Bellotti, A. C., & Lozano, C. . (1993). Implementation of IPM for small-scale cassava farmers. In Miguel A. Altieri (Ed.), *Crop Protection Strategies For Subsistence Farmers* (1st ed., pp. 103–115).
- Buss, E. A., & Turner, J. C. (2006). Scale Insects and Mealybugs on Ornamental Plants. *ENY-323 (MG005)*, University of Florida, 1–6.
- Calatayud, P., & Rü, B. Le. (2006). *Cassava-Mealybug Interactions*.
- Carvalho, G. A., Salgado, L. O., Rigitano, R. L. O., & Velloso, A. H. P. P. (1994). Effects of insect growth regulators on adults of *Ceraeochrysa cubana* (Hagen)(Neuroptera: Chrysopidae). *ANAIS-SOCIEDADE ENTOMOLOGICA DO BRASIL*, 23(2), 335–335.
- Carvalho, G. A., Tironi, P., Rigitano, R. L., & Salgado, L. O. (1994). Selectivity of insect growth regulators to *Trichogramma pretiosum* Riley (Hymenoptera: Trichogrammatidae). *An Soc Entomol Bras*, 23.
- Chakupurakal, J., Markham, R. H., Neuenschwander, P., Sakala, M., Malambo, C., Mulwanda, D., Banda, E., Chalabesa, A., Bird, T., & Haug, T. (1994). Biological control of the cassava mealybug, *Phenacoccus manihoti* (Homoptera: Pseudococcidae), in Zambia. *Biological Control*, 4(3), 254–262. <https://doi.org/10.1006/bcon.1994.1032>
- Cloyd, R. A., & Dickinson, A. (2006). Effect of insecticides on mealybug destroyer (Coleoptera: Coccinellidae) and parasitoid *Leptomastix dactylopii* (Hymenoptera: Encyrtidae), natural enemies of citrus mealybug (Homoptera: Pseudococcidae). *Journal of Economic Entomology*, 99(5), 1596–1604. <https://doi.org/10.1093/jee/99.5.1596>
- Das, G., & Islam, T. (2014). Mortality and growth inhibition of brinjal shoot and fruit borer , *Leucinodes orbonalis* (Guen .) by buprofezin , a potent chitin synthesis inhibitor. *Journal of Entomology and Zoology Studies*, 2(6), 282–287.
- De Santis, L. (1963). Encírtidos de la República Argentina (Hymenoptera: Chalcidoidea). *Anales de La Comisión de Investigación Científica de La Gobernación de La Provincia de Buenos Aires*, 4, 9–422.
- Desneux, N., Decourtye, A., & Delpuech, J. M. (2007). The sublethal effects of pesticides on beneficial arthropods. *Annual Review of Entomology*, 52, 81–106. <https://doi.org/10.1146/annurev.ento.52.110405.091440>
- Desneux, N., Denoyelle, R., & Kaiser, L. (2006). A multi-step bioassay to assess the effect of the deltamethrin on the parasitic wasp *Aphidius ervi*. *Chemosphere*, 65(10), 1697–1706. <https://doi.org/10.1016/j.chemosphere.2006.04.082>
- Downie, D. A., & Gullan, P. J. (2004). Phylogenetic analysis of mealybugs (Hemiptera : Coccoidea : Pseudococcidae) based on DNA sequences from three nuclear genes , and a review of the higher classification. *Systematic Entomology*, 238–259.
- Drobnjakovic, T., & Marcic, D. (2020). Acute toxicity and sublethal effects of pymetrozine on the whitefly parasitoid *Encarsia formosa* Gahan. *Pesticidi i Fitomedicina*, 35(2), 81–95. <https://doi.org/10.2298/pif2002081d>
- Du Toit, W. J., & De Villiers, E. A. (1990). Effects of insect growth regulators on the development of heart-shaped scale (Hemiptera: Coccidae) on avocados in South Africa. *Annals of Applied Biology*, 11(6), 4–5.
- Egho, E. ., Ojianwuna, C. C., & Enujeke, E. C. (2013). Cassava Mealybug's Incidence, Species, Status and Alternative Host Plants in Ethiopie East and Oshimili South Agro-Ecological Zones, Delta State. *International Journal of Science and Nature*, 4(3), 425–430.
- Endo, S., & Tsurumachi, M. (2001). Insecticide Susceptibility of the Brown Planthopper and the White-backed Plant-hopper Collected from Southeast Asia. *Journal of Pesticide Science*, 26(1), 82–86. <https://doi.org/10.1584/jpestics.26.82>
- Fabres, G., & Boussiengue, J. (1981). Bioecology of the cassava mealybug (*Phenacoccus manihoti* Hom. Pseudococcidae) in the People's Republic of Congo. 1-Development

- cycle and biological parameters. *Agronomie Tropicale*, 36(1), 82–89.
- Fanani, M. Z., Rauf, A., Maryana, N., Nurmansyah, A., & Hindayana, D. (2019). Geographic distribution of the invasive mealybug *Phenacoccus manihoti* and its introduced parasitoid *Anagyrus lopezi* in parts of Indonesia. *Biodiversitas*, 20(12), 3751–3757. <https://doi.org/10.13057/biodiv/d201238>
- FAOSTAT. (2019). FAOSTAT. FAO. <http://www.fao.org/faostat/en/#data/QC>
- Garrido, A., Beitia, F., & Gruenholz, P. (1984). Effects of PP618 on immature stages of *Encarsia formosa* and *Cales noacki* (Hymenoptera: Aphelinidae). In *British Crop Protection Conference-Pests and Diseases*, 305–311. Brighton, UK
- Gerling, D., & Sinai, P. (1994). Buprofezin effects on two parasitoid species of whitefly (Homoptera: Aleyrodidae). *Journal of Economic Entomology*, 87(4), 842–846. <https://doi.org/10.1093/jee/87.4.842>
- Gholamzadeh, M., Ghadamyari, M., Salehi, L., & Hoseininaveh, V. (2012). Effects of amitraz, buprofezin and propargite on some fitness parameters of the parasitoid *Encarsia formosa* (Hym.: Aphelinidae), using life table and IOBC methods. *Journal of Entomological Society of Iran*, 31(2), 1–14.
- Giordanengo, P., & Nénon, J. P. (1990). A study of the life cycle of *Epidinocarsis* (= *Apoanagyrus*) *lopezi* (Hymenoptera: Encyrtidae) introduced parasitoid on the cassava mealybug *Phenacoccus manihoti* (Homoptera: Pseudococcidae) in Africa. *Journal of African Zoology*, 104(6), 549–555.
- Gogi, M. D., Syed, A. H., Atta, B., Sufyan, M., Arif, M. J., Arshad, M., Nawaz, A., Khan, M. A., Mukhtar, A., & Liburd, O. E. (2021). Efficacy of biorational insecticides against *Bemisia tabaci* (Genn.) and their selectivity for its parasitoid *Encarsia formosa* Gahan on Bt cotton. *Scientific Reports*, 11(1), 1–12. <https://doi.org/10.1038/s41598-021-81585-x>
- Grafton-cardwell, E. E., Lee, J. E., Stewart, J. R., & Olsen, K. D. (2009). Role of Two Insect Growth Regulators in Integrated Pest Management of Citrus Scales. *Journal of Economic Entomology*, 99(3), 733–744. <https://doi.org/10.1603/0022-0493-99.3.733>
- Gravena, S., Fernandes, O. D., Santos, A. C., Pinto, A. S., & Paiva, P. S. B. (1992). Effect of buprofezin and abamectin on *Pentilia egena* (Muls.) (Coleoptera, Coccinellidae) and *chrysopdis* in citrus. *Anais Da Sociedade Entomologica Do Brasil*, 21, 215–222.
- Grout, T. G., & Richards, G. (1991). Effect of Buprofezin Applications at Different Phenological Times on California Red Scale (Homoptera: Diaspididae). *Journal of Economic Entomology*, 84(6), 1802–1805. <https://doi.org/10.1093/jee/84.6.1802>
- Hassan, S. A. (1994). Comparison of three different laboratory methods and one semi-field test method to assess the side effects of pesticides on *Trichogramma cacoeciae*. *Bulletin OILB/SROP*, 17(10), 133–141.
- Heidari, A., Alford, L., & Kishani Farahani, H. (2015). Effects of three insecticides on adult bionomics of the parasitoid *Encarsia formosa*. *Archives of Phytopathology and Plant Protection*, 48(17–20), 898–909. <https://doi.org/10.1080/03235408.2016.1142926>
- Hennessey, R. D., & Muaka, T. (1987). Field biology of the cassava mealybug, *Phenacoccus manihoti*, and its natural enemies in Zaire. *International Journal of Tropical Insect Science*, 8(4–5–6), 899–903. <https://doi.org/10.1017/s1742758400023158>
- Herren, H. R. (1981). Biological Control of the Cassava Mealybug. In E. R. TERRY, K. A. ODURO, & F. CAVENESS (Eds.), *Tropical Root Crop: RESEARCH STRATEGIES FOR THE 1980s* (pp. 79–80). International Development Research Centre.
- Herren, H. R., & Neuenschwander, P. (1991). Biological control of cassava pests in Africa. *Annual Review of Entomology*. Vol. 36, 257–283. <https://doi.org/10.1146/annurev.ento.36.1.257>
- Hodde, M. S., Van Driesche, R. G., Lyon, S. M., & Sanderson, J. P. (2001). Compatibility of insect growth regulators with *Eretmocerus eremicus* (Hymenoptera: Aphelinidae) for

- whitefly (Homoptera: Aleyrodidae) control on poinsettias: I. Laboratory assays. *Biological Control*, 20(2), 122–131. <https://doi.org/10.1006/bcon.2000.0885>
- Holguín, C., & Bellotti, A. (2004). Efecto de la aplicación de insecticidas químicos en el control de la mosca blanca *Aleurotrachelus socialis* Bondar en el cultivo de yuca *Manihot esculenta* Crantz. *Rev Colomb Entomol*, 30(1), 37–42.
- Iheagwam, E. U. (1981). The influence of temperature on increase rates of the cassava mealybug *Phenacoccus manihoti* Mat.-Ferr. (Homoptera, Pseudococcidae). *Revue de Zoologie Africaine*, 95(4), 959–967.
- Irulandi, S., Kumar, P. K. V., & Sreedharan, K. (2006). Effect of Buprofezin (Applaud) 25 EC on coffee mealy bug, *Planococcus lilacinus* (Cockerell) and its parasitoid, *Leptomastix dactylopii* mulus. In *Journal of Coffee Research* (Vol. 34, pp. 64–71).
- Ishaaya, I., Blumberg, D., & Yarom, I. (1989). Buprofezin: a novel IGR for controlling whiteflies and scale insects. *Mededelingen van de Faculteit Landbouwwetenschappen. Rijksuniversiteit Gent*, 54(3b), 1003–1008.
- Ishaaya, I., Mendel, Z., & D. Blumberg. (1992). Effect of Buprofezin on California red scale, *Aonidiella aurantii* (Maskell), in a citrus orchard. *Israel Journal of Entomology*, 25(3146), 67–71.
- Ishaaya, I., Mendelson, Z., & Melamed-Madjar, V. (1988). Effect of Buprofezin on Embryo genesis and Progeny Formation of Sweet potato Whitefly (Homoptera: Aleyrodidae). *Journal of Economic Entomology*, 81(3), 781–784. <https://doi.org/10.1093/jee/81.3.781>
- Izawa, Y., Uchida, M., Sugimoto, T., & Asai, T. (1985). Inhibition of chitin biosynthesis by buprofezin analogs in relation to their activity controlling *Nilaparvata lugens* Stål. *Pesticide Biochemistry and Physiology*, 24(3), 343–347. [https://doi.org/10.1016/0048-3575\(85\)90145-2](https://doi.org/10.1016/0048-3575(85)90145-2)
- Iziquel, Y., & Le Rü, B. (1992). FECUNDITY, LONGEVITY, AND INTRINSIC NATURAL RATE OF INCREASE OF EPIDINOCARSIS LOPEZI (DE SANTIS)(HYMENOPTERA: ENCYRTIDAE). *The Canadian Entomologist*, 124(6), 1115–1121.
- James, D. G. (2004). Effect of buprofezin on survival of immature stages of *Harmonia axyridis*, *Stethorus punctum picipes* (Coleoptera: Coceinellidae), *Orius tristicolor* (Hemiptera: Anthocoridae), and *Geocoris* spp. (Hemiptera: Geocoridae). *Journal of Economic Entomology*, 97(3), 900–904. <https://doi.org/10.1093/jee/97.3.900>
- Karmakar, P., & Shera, P. S. (2017). Toxicity of insecticides to *Aenasius arizonensis* (Girault) (= *Aenasius bambawalei* Hayat), a solitary endoparasitoid of *Phenacoccus solenopsis* Tinsley on Bt cotton under semi-field conditions. *Journal of Biological Control*, 31(1), 5–9. <https://doi.org/10.18311/jbc/2017/15594>
- Karmakar, P., & Shera, P. S. (2020). Lethal and sublethal effects of insecticides used in cotton crop on the mealybug endoparasitoid *Aenasius arizonensis*. *International Journal of Pest Management*, 66(1), 13–22. <https://doi.org/10.1080/09670874.2018.1538544>
- Karyani, R. D., Maryana, N., & Rauf, A. (2016). Pengujian kekhususan inang parasitoid *Anagyrus lopezi* (De Santis) (Hymenoptera: Encyrtidae) pada empat spesies kutu putih yang berasosiasi dengan tanaman singkong. *Jurnal Entomologi Indonesia*, 13(1), 30–39. <https://doi.org/10.5994/jei.13.1.30>
- Ko, K., Liu, Y., Hou, M., Babendreier, D., Zhang, F., & Song, K. (2015). Toxicity of insecticides targeting rice planthoppers to adult and immature stages of *trichogramma chilonis* (Hymenoptera: Trichogrammatidae). *Journal of Economic Entomology*, 108(1), 69–76. <https://doi.org/10.1093/jee/tou053>
- KOGAN, M., GERLING, D., & MADDOX, J. V. (1999). Enhancement of Biological Control in Annual Agricultural Environments. In T. S. Bellows & F. W. Fisher (Eds.), *Handbook of biological control. Principles and applications of biological control*. (pp. 789–818). San Diego: Academic Press. <http://marefateadyan.nashriyat.ir/node/150>
- Kosztarab, M., & Kozar, F. (1988). *Scale Insects of Central Europe* (S. K.A (ed.); Vol. 41,

Issue 2). <https://doi.org/10.2307/4899>

- Latif, S., & Müller, J. (2014). Cassava – how to explore the “all-sufficient.” *Rural* 21, 30–31.
- Le Rü, B., & Fabres, G. (1987). Influence of temperature and relative humidity on the capacity for increase and population dynamics of the cassava mealybug, *Phenacoccus manihoti* (Hom., Pseudococcidae), in the Congo. *Acta Oecologica, Oecologia Applicata*, 8(2), 165–174.
- Le Rü, B., Iziquel, Y., Biassangama, A., & Kiyindou, A. (1991). Variations d’abondance et facteurs de régulation de la Cochenille du manioc *Phenacoccus manihoti* [Hom.: Pseudococcidae] cinq ans après l’introduction d’*Epidinocarsis lopezi* [Hym.: Encyrtidae] parasitoïde néotropical au Congo en 1982. *Entomophaga*, 6(4), 499-511.
- Lema, K. M., & Herren, H. R. (1985). The influence of constant temperature on population growth rates of the cassava mealybug, *Phenacoccus manihoti*. *Entomologia Experimentalis et Applicata*, 38(2), 165. <https://doi.org/10.1111/j.1570-7458.1985.tb03514.x>
- Lewis, W. J., Stapel, J. O., Cortesero, A. M., & Takasu, K. (1998). Understanding how parasitoids balance food and host needs: Importance to biological control. *Biological Control*, 11(2), 175–183. <https://doi.org/10.1006/bcon.1997.0588>
- Liu, T. X., & Stansly, P. A. (2004). Lethal and sublethal effects of two insect growth regulators on adult *Delphastus catalinae* (Coleoptera: Coccinellidae), a predator of whiteflies (Homoptera: Aleyrodidae). *Biological Control*, 30(2), 298–305. <https://doi.org/10.1016/j.biocontrol.2004.01.007>
- Löhr, B., Neuenschwander, P., Varela, A. M., & Santos, B. (1988). Interactions between the female parasitoid *Epidinocarsis lopezi* De Santis (Hym., Encyrtidae) and its host, the cassava mealybug, *Phenacoccus manihoti* Matile-Ferrero (Hom., Pseudococcidae). *Journal of Applied Entomology*, 105(1–5), 403–413. <https://doi.org/10.1111/j.1439-0418.1988.tb00203.x>
- Löhr, B., Varela, A. M., & Santos, B. (1990). Exploration for natural enemies of the cassava mealybug, *Phenacoccus manihoti* (Homoptera: Pseudococcidae), in South America for the biological control of this introduced pest in Africa. *Bulletin of Entomological Research*, 80(4), 417–425. <https://doi.org/10.1017/S0007485300050677>
- Maharani, J. S., Rauf, A., & Maryana, N. (2020). Masa hidup imago, progeni, dan kemampuan parasitisasi *Anagyrus lopezi* (De Santis) (Hymenoptera: Encyrtidae), parasitoid kutu putih singkong. *Jurnal Entomologi Indonesia*, 16(3), 138. <https://doi.org/10.5994/jei.16.3.138>
- Mani, M., & Shivaraju, C. (2016). Biology. In M. Mani & C. Shivaraju (Eds.), *Mealybugs and their management in agricultural and horticultural crops* (pp. 87–106). Springer India.
- Matazaemon, U., Toshiro, A., & Tatsuyoshi, S. (1985). Inhibition of cuticle deposition and chitin biosynthesis by a new insect growth regulator, buprofezin, in *Nilaparvata lugens* stal. *Agricultural and Biological Chemistry*, 49(4), 1233–1234. <https://doi.org/10.1080/00021369.1985.10866885>
- Matile-Ferrero D. (1977). Une cochenille nouvelle nuisible au manioc en Afrique équatoriale, *Phenacoccus manihoti* n. sp. (Homoptera, Coccoidea, Pseudococcidae). *Ann Soc Entomol Fr New Ser* 13:, 145–152.
- Mendel, Z., Blumberg, D., & Ishaaya, I. (1991). Effect of buprofezin on *Icerya purchasi* and *Planococcus citri*. *Phytoparasitica*, 19(3198), 103–112.
- Mendel, Z., Blumberg, D., & Ishaaya, I. (1994). Effects of some insect growth regulators on natural enemies of scale insects (Hom.: Coccoidea). *Entomophaga*, 39(2), 199–209.
- Meyer, J. B., Kasdorf, G. G. F., Nel, L. H., & Pietersen, G. (2008). Transmission of activated-episomal Banana streak OL (badna)virus (BSOLV) to cv. Williams banana (*Musa* sp.) by three mealybug species. *Plant Disease*, 92(8), 1158–1163. <https://doi.org/10.1094/PDIS-92-8-1158>



- Mgocheki, N., & Addison, P. (2009). Effect of contact pesticides on vine mealybug parasitoids, *Anagyrus* sp. near *pseudococci* (Girault) and *Coccidoxenoides perminutus* (Timberlake) (Hymenoptera: Encyrtidae). *South African Journal of Entology and Viticulture*, 30(2), 110–116. <https://doi.org/10.21548/30-2-1430>
- Miller, D. R., Miller, G. L., & Watson, G. W. (2002). Invasive species of mealybugs (Hemiptera: Pseudococcidae) and their threat to U.S. agriculture. *Proceedings of the Entomological Society of Washington*, 104(4), 825–836.
- Muniappan, R., Shepard, B. M., Watson, G. W., Carner, G. R., Rauf, A., Sartiami, D., Hidayat, P., Afun, J. V. K., Goergen, G., & Rahman, A. K. M. Z. (2009). New records of invasive insects (Hemiptera: Sternorrhyncha) in Southeast Asia and West Africa. *Journal of Agricultural and Urban Entomology*, 26(4), 167–174. <https://doi.org/10.3954/1523-5475-26.4.167>
- Muniappan, R., Shepard, B. M., Watson, G. W., Carner, G. R., Rauf, A., Sartiami, D., Hidayat, P., Afun, J. V. K., Goergen, G., & Rahman, A. K. M. Z. (2011). New records of invasive insects (Hemiptera: Sternorrhyncha) in Southeast Asia and West Africa. *Journal of Agricultural and Urban Entomology*, 26(4), 167–174. <https://doi.org/10.3954/1523-5475-26.4.167>
- Muthukrishnan, N., Manoharan, T., Thevan, P., & Anbu, S. (2005). Evaluation of buprofezin for the management of grape mealy bug, *Maconellicoccus hirsutus* (Green). *J Entomol Res*, 29(4), 339–344.
- Nagai, K. (1990). Effect of insecticides on *Orius* sp., the natural enemy of *Thrips palmi* Karny. *Japanese Journal of Applied Entomology and Zoology, (Japan)*, 34(4), 321–324. <https://doi.org/10.1303/jjaez.34.321>
- Nagata, T. (1986). Timing of buprofezin application for control of the brown planthopper, *Nilaparvata lugens* (Stål) (Homoptera: Delphacidae). *Applied Entomology and Zoology*, 14, 357–368.
- Nagrare, V. S., Kranthi, S., Kranthi, K. R., Naik, V. C. B., Deshmukh, V., Naikwadi, B., & Dahekar, A. (2016). Relative toxicity of insecticides against cotton mealybug *Phenacoccus solenopsis* Tinsley (Hemiptera: Pseudococcidae) and its fortuous parasitoid *Aenasius bambawalei* Hayat (Hymenoptera: Encyrtidae). *Journal of Applied and Natural Science*, 8(2), 987–994. <https://doi.org/10.31018/jans.v8i2.909>
- Narayana, M. L., & Babu, T. R. (1992). Evaluation of five insect growth regulators on the egg parasitoid *Trichogramma chilonis* (Ishii) (Hym., Trichogrammatidae) and the hatchability of *Corcyra cephalonica* Staint (Lep., Galleriidae). *Journal of Applied Entomology*, 113(1–5), 56–60. <https://doi.org/10.1111/j.1439-0418.1992.tb00635.x>
- Neuenschwander, P. (1994). Control of the cassava mealybug in Africa: Lessons from a biological control project. *African Crop Science Journal*, 2(4), 369–383.
- Neuenschwander, P. (2001). Biological control of the cassava mealybug in Africa: A review. *Biological Control*, 21(3), 214–229.
- Neuenschwander, P., & Hammond, W. N. O. (1988). Natural Enemy Activity Following the Introduction of *Epidinocarsis lopezi* (Hymenoptera: Encyrtidae) Against the Cassava Mealybug, *Phenacoccus manihoti* (Homoptera: Pseudococcidae), in Southwestern Nigeria. *Environmental Entomology*, 17(5), 894–902.
- Nguyen Tuan, D., Sam, L., Zhang, C., Nguyen Ngoc Bao, C., Takano, S. ichiro, & Takasu, K. (2020). Taro *Colocasia esculenta* as an alternative host plant for rearing cassava mealybug (Hemiptera: Pseudococcidae) and its parasitoid *Anagyrus lopezi* (Hymenoptera: Encyrtidae). *Applied Entomology and Zoology*, 55(3), 355–359. <https://doi.org/10.1007/s13355-020-00690-x>
- Noyes, J. S., & Hayat, M. (1994). *Oriental Mealybug Parasitoids of the Anagyrini (Hymenoptera: Encyrtidae)*. Cab International.
- Nwanze, K. F. (1978). Biology of the cassava mealybug, *Phenacoccus manihoti* Mat.-Ferr. in

- the Republic of Zaire. *International Institute of Tropical Agriculture*, 20–28.
- Nwanze, K. F. (1982). Relationships between cassava root yields and crop infestations by the mealybug, *phenacoccus manihoti*. *Tropical Pest Management*, 28(1), 27–32. <https://doi.org/10.1080/09670878209370669>
- Odebiyi, J. A., & Bokonon-Ganta, A. H. (1986). Biology of *Epidinocarsis* [=Apoanagyrus] *lopezi* [Hymenoptera: Encyrtidae] an exotic parasite of cassava mealybug, *Phenacoccus manihoti* [Homoptera: Pseudococcidae] in Nigeria. *Entomophaga*, 31(3), 251–260. <https://doi.org/10.1007/BF02373334>
- Olsen, K. M., & Schaal, B. A. (1999). Evidence on the origin of cassava: Phylogeography of *Manihot esculenta*. *Proceedings of the National Academy of Sciences of the United States of America*, 96(10), 5586–5591. <https://doi.org/10.1073/pnas.96.10.5586>
- Parsa, S., Kondo, T., & Winotai, A. (2012). The Cassava Mealybug (*Phenacoccus manihoti*) in Asia: First Records, Potential Distribution, and an Identification Key. *PLoS ONE*, 7(10). <https://doi.org/10.1371/journal.pone.0047675>
- Poletti, M., Maia, A. H. N., & Omoto, C. (2007). Toxicity of neonicotinoid insecticides to *Neoseiulus californicus* and *Phytoseiulus macropilis* (Acari: Phytoseiidae) and their impact on functional response to *Tetranychus urticae* (Acari: Tetranychidae). *Biological Control*, 40(1), 30–36. <https://doi.org/10.1016/j.biocontrol.2006.09.001>
- Prabhaker, N., Morse, J. G., Castle, S. J., Naranjo, S. E., Henneberry, T. J., & Toscano, N. C. (2007). Toxicity of seven foliar insecticides to four insect parasitoids attacking citrus and cotton pests. *Journal of Economic Entomology*, 100(4), 1053–1061. [https://doi.org/10.1603/0022-0493\(2007\)100\[1053:TOSFIT\]2.0.CO;2](https://doi.org/10.1603/0022-0493(2007)100[1053:TOSFIT]2.0.CO;2)
- Rashidi, F., & Ganbalani, G. N. (2018). Toxicity and sublethal effects of selected insecticides on life parameters of *encarsia formosa* (Hymenoptera: Aphelinidae), a parasitoid of *trialeurodes vaporariorum* (Hemiptera: Aleyrodidae). *Journal of Entomological Science*, 53(4), 543–553. <https://doi.org/10.18474/JES18-03.1>
- Rauf, A. (2016). *Infestations and management of the cassava mealybug, Phenacoccus manihoti (Hemiptera: Pseudococcidae), a newly invasive pest in Indonesia*. 26(4), 2014. <https://doi.org/10.1603/ice.2016.117544>
- Rill, S. M., Grafton-Cardwell, E. E., & Morse, J. G. (2008). Effects of two insect growth regulators and a neonicotinoid on various life stages of *Aphytis melinus* (Hymenoptera: Aphelinidae). *BioControl*, 53(4), 579–587. <https://doi.org/10.1007/s10526-007-9097-x>
- Rivnay, E., & Perzelan, J. (1943). Insects associated with *Pseudococcus* spp. (Homoptera) in Palestine, with notes on their biology and economic status. *Journal of the Entomological Society of Southern Africa*, 6, 9–28. <http://content.ajarchive.org/cgi-bin/showfile.exe?CISOROOT=/0012-8789&CISOPTR=1648>
- Sartiami, D., Watson, G. W., Mohamad Roff, M. N., Mohd Hanifah, Y., & Idris, A. B. (2015). First record of cassava mealybug, *Phenacoccus manihoti* (Hemiptera: Pseudococcidae), in Malaysia. *Zootaxa*, 3957(2), 235–238. <https://doi.org/10.11646/zootaxa.3957.2.8>
- Sataloff, R. T., Johns, M. M., & Kost, K. M. (2014). *Sustainable Management of Cassava in Asia, From Research to Practice*. Centro Internacional de Agricultura Tropical (CIAT).
- Schulthess, F., Baumgärtner, J. U., Delucchi, V., & Gutierrez, A. P. (1991). The influence of the cassava mealybug, *Phenacoccus manihoti* Mat.-Ferr. (Hom., Pseudococcidae) on yield formation of cassava, *Manihot esculenta* Crantz. *Journal of Applied Entomology*, 111(1–5), 155–165. <https://doi.org/10.1111/j.1439-0418.1991.tb00306.x>
- Schulthess, F., Baumgärtner, J. U., & Herren, H. R. (1987). Factors influencing the life table statistics of the cassava mealybug *Phenacoccus manihoti*. *International Journal of Tropical Insect Science*, 8(4–5), 851–856. <https://doi.org/10.3929/ethz-a-010025751>
- Shylesha, A. N., & Mani, M. (2016). Natural Enemies of Mealybugs. In M. Mani & C. Shivaraju (Eds.), *Mealybugs and their management in agricultural and horticultural*

- crops* (pp. 149–171). Springer India.
- Simsek, V. M., Uygun, N., & Satar, S. (2008). *Investigation of Side Effects of Some Pesticides under Laboratory Conditions on Important Parasitoids and Predators in Citrus Ecosystem*. 38(November 2006), 9067. <https://doi.org/10.1002/ps.2780430308>.Tuzcu
- Smith, D. (1995). *Effect of the insect growth regulator buprofezin against citrus pests *Coccus viridis* Green , *Polyphagotarsonemus latus* (Banks) and *Aonidiella aurantii* (Maskell) and the predatory Coccinellid *Chilocorus circumdatus* Gyllenhal*. 10(3), 112–115.
- Sohrabi, F., Shishehbor, P., Saber, M., & Mosaddegh, M. S. (2012). Lethal and sublethal effects of buprofezin and imidacloprid on the whitefly parasitoid *Encarsia inaron* (Hymenoptera: Aphelinidae). *Crop Protection*, 32, 83–89. <https://doi.org/10.1016/j.cropro.2011.10.005>
- Sohrabi, F., Shishehbor, P., Saber, M., & Mosaddegh, M. S. (2013). Lethal and sublethal effects of imidacloprid and buprofezin on the sweetpotato whitefly parasitoid *Eretmocerus mundus* (Hymenoptera: Aphelinidae). *Crop Protection*, 45, 98–103. <https://doi.org/10.1016/j.cropro.2012.11.024>
- Sontakke, B. K., Mohapatra, L. N., & Swain, L. K. (2013). *Comparative bioefficacy of buprofezin 25 EC against sucking pests of cotton and its safety to natural enemies*. 75Sontakke(4), 325–329.
- Suma, P., Zappalà, L., Mazzeo, G., & Siscaro, G. (2009). Lethal and sub-lethal effects of insecticides on natural enemies of citrus scale pests. *BioControl*, 54(5), 651–661. <https://doi.org/10.1007/s10526-009-9215-z>
- Takasu, K. E. I. J. I., & Lewis., W. J. (1993). Host-and food-foraging of the parasitoid *Microplitis croceipes*: learning and physiological state effects. *Biological Control*, 3,(1), 70–74.
- Uchida, M., Izawa, Y., & Sugimoto, T. (1987). Inhibition of prostaglandin biosynthesis and oviposition by an insect growth regulator, buprofezin, in *Nilaparvata lugens* Stål. *Pesticide Biochemistry and Physiology*, 27(1), 71–75. [https://doi.org/10.1016/0048-3575\(87\)90097-6](https://doi.org/10.1016/0048-3575(87)90097-6)
- Van De Veire, M. (1995). Integrated pest management in glasshouse tomatoes, sweet peppers and cucumbers in Belgium. (*Doctoral Dissertation, Ghent University*).
- Van Driesche, R. G., & Bellows, T. S. (1996). *Biology of Arthropod Parasitoids and Predators*. In: *Biological Control*.
- Walker, A., Hoy, M., & Meyerdirk, D. (2003). Papaya mealybug , *Paracoccus marginatus* Williams and Granara de Willink. *EENY-302, One of a Series of Featured Creatures from the Entomology and Nematology Department, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida*, 1–7.
- Walker, J. T. S., White, V., & Charles, J. G. (1993). Field control of chlorpyrifos-resistant mealybugs (*Pseudococcus affinis*) in a Hawkes Bay orchard. *In Proceedings of the New Zealand Plant Protection Conference*, 46, 126–128.
- Wang. Q. L., & Liu, T.-X. (2016). Effects of three insect growth regulators on *Encarsia formosa* (Hymenoptera: Aphelinidae), an endoparasitoid of *Bemisia tabaci* (Hemiptera: Aleyrodidae). *Journal of Economic Entomology*, 109(6), 2290–2297. <https://doi.org/10.1093/jee/tow216>
- Wang, Z., Dai, P., Yang, X., Ruan, C. C., Biondi, A., Desneux, N., & Zang, L. S. (2019). Selectivity of novel and traditional insecticides used for management of whiteflies on the parasitoid *Encarsia formosa*. *Pest Management Science*, 75(10), 2716–2724. <https://doi.org/10.1002/ps.5380>
- Wardani, N. (2015). *Kutu Putih Ubi Kayu, Phenacoccus manihoti Matile-Ferrero (Hemiptera: Pseudococcidae), Hama Invasif Baru di Indonesia*. Institut Pertanian Bogor, Bogor.
- Williams, D. J., & Willink, M. C. G. de. (1992). *Mealybugs of Central and South America*.



Impact of Buprofezin on Performance *Anagyrus lopezi* (Hymenoptera: Encyrtidae) the Parasitoid of Cassava Mealybug *Phenacoccus manihoti* (Hemiptera: Pseudococcidae)

Mesa Mon, Prof. Dr. Ir. Edhi Martono, M.Sc.; Prof. Dr. Ir. FX Wagiman, S.U.

Universitas Gadjah Mada, 2023 | Diunduh dari <http://etd.repository.ugm.ac.id/>

UNIVERSITAS
GADJAH MADA

- Willink, M. C. G. De. (2003). New records and host plants of *Phenacoccus* from Argentina (Hemiptera: Pseudococcidae). *Journal of the Argentine Entomological Society* ., 62(3/4), 80–82.
- Winotai, W., Goergen, G., Tamò, M., & Neuenschwander, P. (2010). Cassava mealybug has reached Asia. *Biocontrol News and Information*.
- Wyckhuys, K., & Rauf, A. (2014). Parasitoids introduced into Indonesia: part of a region-wide campaign to tackle emerging cassava pests and diseases. *Biocontrol News and Information*. <https://cgspace.cgiar.org/handle/10568/57006>
- Yarom, I., Blumberg, D., & Ishaaya, I. (1988). Effects of Buprofezin on California Red Scale (Homoptera: Diaspididae) and Mediterranean Black Scale (Homoptera: Coccidae). *Journal of Economic Entomology*, 81(6), 1581–1585. <https://doi.org/10.1093/jee/81.6.1581>
- Yonow, T., Kriticos, D. J., & Ota, N. (2017). The potential distribution of cassava mealybug (*Phenacoccus manihoti*), a threat to food security for the poor. *PLoS ONE*, 12(3), 1–17. <https://doi.org/10.1371/journal.pone.0173265>
- Zeddies, J., Schaab, R. P., Neuenschwander, P., & Herren, H. R. (2001). Economics of biological control of cassava mealybug in Africa. *Agricultural Economics*, 24(2), 209–219. [https://doi.org/10.1016/S0169-5150\(00\)00064-5](https://doi.org/10.1016/S0169-5150(00)00064-5)