



DAFTAR PUSTAKA

- Aboonabi, A., Rahmat, A., & Othman, F. (2014). Antioxidant effect of pomegranate against streptozotocin-nicotinamide generated oxidative stress induced diabetic rats. *Toxicology Reports*, 1, 915-922. <https://dx.doi.org/10.1016%2Fj.toxrep.2014.10.022>.
- Adebawale, K. O., Afolabi, T. A., & Lawal, O. S. (2002). Isolation, chemical modification and physicochemical characterisation of Bambara groundnut (*Voandzeia subterranean*) starch and flour. *Food chemistry*, 78(3), 305-311. [https://doi.org/10.1016/S0308-8146\(02\)00100-0](https://doi.org/10.1016/S0308-8146(02)00100-0)
- Adebawale, K. O., Olu-Owolabi, B. I., kehinde Olawumi, E., & Lawal, O. S. (2005). Functional properties of native, physically and chemically modified breadfruit (*Artocarpus atilis*) starch. *Industrial crops and products*, 21(3), 343-351. <https://doi.org/10.1016/j.indcrop>.
- Adewumi, D. F., Lajide, L., Afolabi, O. B., & Ojo, A. O. (2015). Comparative Study on the Pasting Properties of Native and Acetylated Cocoyam (*Xanthosoma sagittifolium*) Starch Contents. *International Journal of Applied Research and Technology*, 4(7), 147-151. <https://www.researchgate.net/publication/282327915>
- Adriawan, I. R., Andrie, M., Susilowati, R., Pramono, S., dan Nugroho, A. E. (2014). Evaluasi efek anti-diabetes mellitus ekstrak terpurifikasi *Andrographis paniculata* (burn,f.) nees dan andrografolid dengan parameter indeks HOMA-IR, *Traditional Medicine Journal*, 19(11): 19-23. ISSN : 1410-5918
- Alvani, K., Qi, X., Tester, R. F., & Snape, C. E. (2011). Physico-chemical properties of potato starches. *Food Chemistry*, 125(3), 958-965. <https://doi.org/10.1016/j.foodchem.2010.09.088>
- Anderson, J., Briefel, G., Jones, J. M., Ryu, J. H., McGuire, M., & Yun, Y. P. (1991). Effects of acetate dialysate on transforming growth factor β 1, interleukin, and β 2-microglobulin plasma levels. *Kidney international*, 40(6), 1110-1117.
- Annison, G., Illman, R. J., & Topping, D. L. (2003). Acetylated, propionylated or butyrylated starches raise large bowel short-chain fatty acids preferentially when fed to rats. *The Journal of nutrition*, 133(11), 3523-3528. <https://doi.org/10.1093/jn/133.11.3523>
- AOAC: Official Methods of Analysis. (1990). 15th Edition (Vol 1), Association of office analytical chemist. Inc, Suite 400. 2200 Wilson Boulevard, Arlington, Virginia 22201 USA



- Arifin, H., Delvita, V., & Almahdy, A. (2007). Pengaruh pemberian vitamin C terhadap fetus pada mencit diabetes. *Jurnal Sains dan Teknologi Farmasi*, 12(1), 32-40. Akreditasi DIKTI Depdiknas RI No. 49/DIKTI/Kep/2003. ISSN : 1410 – 0177
- Aparicio-Saguilán, A., Gutiérrez-Meraz, F., García-Suárez, F. J., Tovar, J., & Bello-Pérez, L. A. (2008). Physicochemical and functional properties of cross-linked banana resistant starch. Effect of pressure cooking. *Starch-Stärke*, 60(6), 286-291.
<https://doi.org/10.1002/star.200600584>
- Aparicio-Saguilán, A., Flores-Huicochea, E., Tovar, J., García-Suárez, F., Gutiérrez-Meraz, F., & Bello-Pérez, L. A. (2005). Resistant starch-rich powders prepared by autoclaving of native and ligninized banana starch: Partial characterization. *Starch-Stärke*, 57(9), 405-412. DOI 10.1002/star.200400386
- Apriyantono, A., Fardiaz, D., Puspitasari, N. L., Sedarnawati, S. B., & Budiyantono, S. (1989). Analisis Pangan: Petunjuk Laboratorium. *IPB, Bogor*.
- Ashogbon, A. O. (2014). Chemical and functional properties of cocoyam starch and wheat starch blends. *International Journal of Biotechnology and Food Science*, 2(5), 94-101.
- Ashogbon, A. O., & Akintayo, E. T. (2014). Recent trend in the physical and chemical modification of starches from different botanical sources: A review. *Starch-Stärke*, 66(1-2), 41-57.
<https://doi.org/10.1002/star.201300106>
- Ashwar, B. A., Gani, A., Shah, A., & Masoodi, F. A. (2017). Physicochemical properties, in-vitro digestibility and structural elucidation of RS4 from rice starch. *International journal of biological macromolecules*, 105, 471-477.
<https://doi.org/10.1016/j.ijbiomac.2017.07.057>
- ASP, N. G. (1992). Resistant starch. Proceeding from the second plenary meeting of EURESTA: European FLAIR concerted action no. 11 on physiological implications of the consumption of resistant starch in man. *Eur, J. Clin, Nutr.*, 46, S1. ISBN 90-5485-651-3
- Arisandy, O. M. P., & Estiasih, T. (2016). Beras tiruan berbasis tepung kimpul (*Xanthosoma sagittifolium*): Kajian pustaka . *Jurnal Pangan dan Agroindustri*, 4(1); 253-261. 326-Article Text-760-1-10-20150519.pdf



Aryanti, N., Kusumastuti, Y. A., & Rahmawati, W. (2017). Pati talas (*Colocasia esculenta* (L.) Schott) sebagai alternatif sumber pati industri. *Majalah Ilmiah MOMENTUM*, 13(1); 46-52. ISSN 0216-7395

Awokoya, K. N., Moronkola, B. A., & Tovide, O. O. (2012). Characterization of starches from red cocoyam (*Colocasia esculenta*) and white cocoyam (*Colocasia antiquorum*) cormels. *Food Sc. Qual. Mgt*, 5, 27-35. ISSN 2224-6088 (Paper), ISSN 2225-0557 (Online).

Ayucitra, A. (2012). Preparation and characterisation of acetylated corn starches. *International Journal of Chemical Engineering and Applications*, 3(3), 156. DOI:[10.7763/IJCEA.2012.V3.178](https://doi.org/10.7763/IJCEA.2012.V3.178)

Ayu, D. C., & Yuwono, S. S. (2014). Oengaruh suhu blansing dan lama perendaman terhadap sifat fisik kimia tepung kimpul (*Xanthosoma Sagittifolium*)[IN PRESS APRIL 2014]. *Jurnal Pangan dan Agroindustri*, 2(2), 110-120.

Banerjee, M., & Vats, P. (2014). Reactive metabolites and antioxidant gene polymorphisms in type 2 diabetes mellitus. *Redox biology*, 2, 170-177. DOI: [10.1016/j.redox.2013.12.001](https://doi.org/10.1016/j.redox.2013.12.001)

Babic, J., Subasic, D., Ackar, D., Kovacevic, D., Pilizota, V., & Kopjar, M. (2007). Preparation and characterization of acetylated tapioca starches. *deutsche lebensmittelrundschau*, 103(12), 580. https://www.researchgate.net/publication/279710937_Preparation_and_characterization_of_acetylated_tapioca_starches

Bartz, J., Goebel, J. T., Giovanaz, M. A., da Rosa Zavareze, E., Schirmer, M. A., & Dias, A. R. G. (2015). Acetylation of barnyardgrass starch with acetic anhydride under iodine catalysis. *Food chemistry*, 178, 236-242. <https://doi.org/10.1016/j.foodchem.2015.01.095>

Behall, K. M., Scholfield, D. J., Hallfrisch, J. G., & Liljeberg-Elmståhl, H. G. (2006). Consumption of both resistant starch and β -glucan improves postprandial plasma glucose and insulin in women. *Diabetes care*, 29(5), 976-981. <https://doi.org/10.2337/diacare.295976>

Belitz, H. D., and W. Grosch. (1999). Food chemistry. Springer Verlag, Berlin.

Bello-Pérez, L. A., Agama-Acevedo, E., Zamudio-Flores, P. B., Mendez-Montealvo, G., & Rodriguez-Ambriz, S. L. (2010). Effect of low and high acetylation degree in the morphological, physicochemical and structural characteristics of barley starch. *LWT-Food Science and Technology*, 43(9), 1434-1440. <https://doi.org/10.1016/j.lwt.2010.04.003>



- Betancur, A. D., Chel, G. L., & Cañizares, H. E. (1997). Acetylation and characterization of *Canavalia ensiformis* starch. *Journal of Agricultural and Food Chemistry*, 45(2), 378-382. <https://doi.org/10.1021/jf960272e>
- Bird, A. R., Brown, I. L., & Topping, D. L. (2006). Low and high amylose maize starches acetylated by a commercial or a laboratory process both deliver acetate to the large bowel of rats. *Food hydrocolloids*, 20(8), 1135-1140. <https://doi.org/10.1016/j.foodhyd.2005.12.009>
- Birt, D. F., Boylston, T., Hendrich, S., Jane, J. L., Hollis, J., Li, L., ... & Whitley, E. M. (2013). Resistant starch: promise for improving human health. *Advances in nutrition*, 4(6), 587-601. <https://doi.org/10.3945/an.113.004325>
- Blazek, J., & Copeland, L. (2008). Pasting and swelling properties of wheat flour and starch in relation to amylose content. *Carbohydrate polymers*, 71(3), 380-387. <https://doi.org/10.1016/j.carbpol.2007.06>
- Bourquin, L. D., Titgemeyer, E. C., & Fahey Jr, G. C. (1993). Vegetable fiber fermentation by human fecal bacteria: cell wall polysaccharide disappearance and short-chain fatty acid production during in vitro fermentation and water-holding capacity of unfermented residues. *The Journal of nutrition*, 123(5), 860-869. <http://dx.doi.org/10.1093/jn/123.5.860>
- Bradbury, H. J., and W. D. Holloway. (2000). Chemistry of tropical root crops. Australian centre for international agriculture research, Canberra.
- Brites, C. M., Trigo, M. J., Carrapico, B., Alviña, M., & Bessa, R. J. (2011). Maize and resistant starch enriched breads reduce postprandial glycemic responses in rats. *Nutrition Research*, 31(4), 302-308. <https://doi.org/10.1016/j.nutres.2011.02.001>
- Bukabi - Deptan. (2009). Umbi-umbian. Direktorat budidaya kacang - kacangan dan umbi - umbian. Departemen Pertanian.
- BPS Maluku. (2017). Badan Pusat Statistik (BPS) Propinsi Maluku. Maluku Dalam Angka. BPS Maluku. Maluku.
- Canfora, E. E., Jocken, J. W., & Blaak, E. E. (2015). Short-chain fatty acids in control of body weight and insulin sensitivity. *Nature Reviews Endocrinology*, 11(10), 577-591. DOI: [10.1038/nrendo.2015.128](https://doi.org/10.1038/nrendo.2015.128)
- Chandrashekhar, K. N. (2009). Evidence of oxidative stress and mitochondrial dysfunctions in the testis of prepubertal diabetic rats. *International journal of impotence research*, 21(3), 198-206.
- Chaplin, M. 2002. Starch. <http://www.sbu.ac.uk/>



Chávez-Murillo, C. E., Wang, Y. J., & Bello-Pérez, L. A. (2008). Morphological, physicochemical and structural characteristics of oxidized barley and corn starches. *Starch-Stärke*, 60(11), 634-645.
<http://dx.doi.org/10.1002/star.200800016>

Chen, Z., Schols, H. A., & Voragen, A. G. (2004). Differently sized granules from acetylated potato and sweet potato starches differ in the acetyl substitution pattern of their amylose populations. *Carbohydrate Polymers*, 56(2), 219-226. <https://doi.org/10.1016/j.carbpol.2004.02.004>

Chen, J., Wu, F., Long, Y., & Yu, W. (2015). Glutathione supplementation attenuates oxidative stress and improves vascular hyporesponsiveness in experimental obstructive jaundice. *Oxidative Medicine and Cellular Longevity*, 2015. <https://doi.org/10.1155/2015/486148>

Cheng, H. H., & Lai, M. H. (2000). Fermentation of resistant rice starch produces propionate reducing serum and hepatic cholesterol in rats. *The Journal of nutrition*, 130(8), 1991-1995. <https://doi.org/10.1093/jn/130.8.1991>

Cooke, D., & Gidley, M. J. (1992). Loss of crystalline and molecular order during starch gelatinisation: origin of the enthalpic transition. *Carbohydrate research*, 227, 103-112. [https://doi.org/10.1016/0008-6215\(92\)85063-6](https://doi.org/10.1016/0008-6215(92)85063-6)

Chung, H. J., Shin, D. H., & Lim, S. T. (2008). In vitro starch digestibility and estimated glycemic index of chemically modified corn starches. *Food research international*, 41(6), 579-585. <https://doi.org/10.1016/j.foodres.2008.04.006>

Cook, S. I., & Sellin, J. H. (1998). Short chain fatty acids in health and disease. *Alimentary pharmacology & therapeutics*, 12(6), 499-507.

Colussi, R., Pinto, V. Z., El Halal, S. L. M., Vanier, N. L., Villanova, F. A., e Silva, R. M., ... & Dias, A. R. G. (2014). Structural, morphological, and physicochemical properties of acetylated high-, medium-, and low-amylose rice starches. *Carbohydrate Polymers*, 103, 405-413. <https://doi.org/10.1016/j.carbpol.2013.12.070>

Colussi, R., El Halal, S. L. M., Pinto, V. Z., Bartz, J., Gutkoski, L. C., da Rosa Zavareze, E., & Dias, A. R. G. (2015). Acetylation of rice starch in an aqueous medium for use in food. *LWT-Food Science and Technology*, 62(2), 1076-1082. <https://doi.org/10.1016/j.lwt.2015.01.053>

Cornejo-Ramírez, Y. I., Martínez-Cruz, O., Del Toro-Sánchez, C. L., Wong-Corral, F. J., Borboa-Flores, J., & Cinco-Moroyoqui, F. J. (2018). The structural characteristics of starches and their functional properties. *CyTA-*



Journal of Food, 16(1), 1003-1017.
<https://www.tandfonline.com/action/showCitFormats?doi=10.1080/19476337.2018.1518343>

Coronell-Tovar, D. C., Chavez-Jauregui, R. N., Bosques-Vega, A., & Lopez-Moreno, M. L. (2018). Characterization of cocoyam (*Xanthosoma* spp.) corm flour from the Nazareno cultivar. *Food Science and Technology*, 39, 349-357. <https://doi.org/10.1590/fst.30017>

Cummings, J. H., & Bingham, S. A. (1987). Dietary fibre, fermentation and large bowel cancer. *Cancer surveys*, 6(4), 601-621. doi: 10.1136/gut.28.10.1221.

Czuchajowska, Z., Sievert, D., & Pomeranz, Y. (1991). Enzyme-resistant starch. IV. Effects of complexing lipids. *Cereal Chem*, 68(5), 537-542.

Damat, D., Marsono, M., Haryadi, H., & Cahyanto, C. (2008). Efek hipokolesterolemik dan hipoglikemik patigarut butirat pada tikus Sprague Dawley. *Majalah Farmasi Indonesia*, 19(3), 109-116.

Damat, D., & Kurniawati, K. (2016). Karakteristik fisiko-kimia pati garut (marantha arundinaceae) termodifikasi secara fisik melalui proses gelatinisasi retrogradasi berulang. 160-166. *Seminar Nasional Hasil Penelitian*, 2016-UNMUH.

Djunaidi, C. S., Affandi, D. R., & Praseptiangga, D. (2014). Efek hipoglikemik tepung komposit (ubijalar ungu, jagung kuning, dan kacang tunggak) pada tikus diabetes induksi streptozotocin. *Jurnal Gizi Klinik Indonesia*, 10(3), 119-126. <https://doi.org/10.22146/ijcn.18859>

Deetae, P., P. Chinachoti., S. Shobsngob., O. Naivikul., and W. Varanyanond., Varavinit, S. (2008). Preparation, pasting properties and freeze-thaw stability of dual modified crosslink-phosphorilated rice starch *Carbohydrate Polymers*, 73: 351-358. <http://dx.doi.org/10.1016/j.carbpol.2007.12.004>

Den Besten, G., Van Eunen, K., Groen, A. K., Venema, K., Reijngoud, D. J., & Bakker, B. M. (2013). The role of short-chain fatty acids in the interplay between diet, gut microbiota, and host energy metabolism. *Journal of lipid research*, 54(9), 2325-2340. <https://doi.org/10.1194/jlr.R036012>

Dian Kartika Sari. Pati tahan cerna untuk pencegahan diabetes Tipe-2. Program Studi IKM, Program Pascasarjana, Fakultas Kedokteran, Universitas Padjadjaran.pustaka_unpad_Pati_-Tahan_-Cerna.pdf.pdf

Dolas, K. A., Ranveer, R. C., Tapre, A. R., Nandane, A. S. and Sahoo, A. K. (2020). Effect of starch modification on physico-chemical, functional and



structural characterization of cassava starch (*Manihot esculenta* Crantz).
Food Research, 4(4): 1265 – 1271 DOI:[10.26656/fr.2017.4\(4\).075](https://doi.org/10.26656/fr.2017.4(4).075)

Eerlingen, R. C., & Delcour, J. A. (1995). Formation, analysis, structure and properties of type III enzyme resistant starch. *Journal of Cereal Science*, 22(2), 129-138. [https://doi.org/10.1016/0733-5210\(95\)90042-X](https://doi.org/10.1016/0733-5210(95)90042-X)

Elayat, A. A., el-Naggar, M. M., & Tahir, M. (1995). An immunocytochemical and morphometric study of the rat pancreatic islets. *Journal of anatomy*, 186(Pt 3), 629. <http://www.ncbi.nlm.nih.gov/pmc/articles/pmc1167020/>

Falade, K. O., & Okafor, C. A. (2013). Physicochemical properties of five cocoyam (*Colocasia esculenta* and *Xanthosoma sagittifolium*) starches. *Food Hydrocolloids*, 30(1), 173-181. <https://doi.org/10.1016/j.foodhyd.2012.05.006>

Faridah, D. N., Fardiaz, D., Andarwulan, N., & Sunarti, T. C. (2014). Karakteristik sifat fisikokimia pati garut (Maranta arundinaceae). *Agritech*, 34(1), 14-21. <https://doi.org/10.22146/agritech.9517>

Fennema OR.1985. Food Chemistry. Marcel Dekker Inc. New York. Basel

Fiqtinovri, S. M., Benita, A. M., Marseno, D. W., & Pranoto, Y. (2020). Sifat Fisik, Amilograf, dan Morfologi Pati Biji Lai (Durio kutejensis) Asetilasi Menggunakan Asetat Anhidrat. *agriTECH*, 40(1), 74-83. <http://dx.doi.org/10.22146/agritech.33809>

Fuentes-Zaragoza, E., Riquelme-Navarrete, M. J., Sánchez-Zapata, E., & Pérez-Álvarez, J. A. (2010). Resistant starch as functional ingredient: A review. *Food Research International*, 43(4), 931-942. <https://doi.org/10.1016/j.foodres.2010.02.004>

Fleche, G. (1985). Chemical modification and degradation of starch. *Di dalam* G.M.A. Van Beynum dan J.A. Roels, ed. Starch conversion technology. London: Applied Science Publ

Fortuna, T., Juszczak, L., & Palasinski, M. (2001). Properties of corn and wheat starch phosphates obtained from granules segregated according to their size. *Electronic journal of polish agricultural universities*, 4(2). <http://www.ejpau.media.pl/volume4/issue2/food/art-05.html>

Gao, Z., Yin, J., Zhang, J., Ward, R. E., Martin, R. J., Lefevre, M., ... & Ye, J. (2009). Butyrate improves insulin sensitivity and increases energy expenditure in mice. *Diabetes*, 58(7), 1509-1517. <https://doi.org/10.2337/db08-1637>



- Ganjyal, G. M., Hanna, M. A., & Jones, D. D. (2003). Modeling selected properties of extruded waxy maize cross-linked starches with neural networks. *Journal of food science*, 68(4), 1384-1388. <https://doi.org/10.1111/j.1365-2621.2003.tb09654.x>
- Ganong, W. F. (1993). Review of medica; physiologi. 16^{ed}. Lange Medical Publication. San Fransisco, California.
- Goñi, I., Garcia-Diz, L., Mañas, E., & Saura-Calixto, F. (1996). Analysis of resistant starch: a method for foods and food products. *Food chemistry*, 56(4), 445-449. [https://doi.org/10.1016/0308-8146\(95\)00222-7](https://doi.org/10.1016/0308-8146(95)00222-7)
- Green, C. J., Henriksen, T. I., Pedersen, B. K., & Solomon, T. P. (2012). Glucagon like peptide-1-induced glucose metabolism in differentiated human muscle satellite cells is attenuated by hyperglycemia. *PLoS One*. 2012;7(8):e44284. <https://doi.org/10.1371/journal.pone.0044284>
- Greenwood, C. T dan D. N. Munro. (1979). Carbohydrates. *Di dalam* R.J. Priestley,ed. Effects of heat on foodstufs. *Applied Science* Publ. Ltd., London
- Golachowski, A., Zięba, T., Kapelko-Żeberska, M., Drożdż, W., Gryszkin, A., & Grzechac, M. (2015). Current research addressing starch acetylation. *Food chemistry*, 176, 350-356. <https://doi.org/10.1016/j.foodchem.2014.12.060>
- Gujska, E., D.-Reinhard, W., & Khan, K. (1994). Physicochemical properties of field pea, pinto and navy bean starches. *Journal of Food Science*, 59(3), 634-636. <https://doi.org/10.1111/j.1365-2621.1994.tb05580.x>
- Gunaratne, A., R. Hoover. Effect of heta-moisture treatment on the stucture and physicochemical properties of tuber and root starches. *Carbohydrate Polymers*, 49: 425-437. [http://dx.doi.org/10.1016/S0144-8617\(01\)00354-X](http://dx.doi.org/10.1016/S0144-8617(01)00354-X)
- Guyton, A. C. Hall, J. E. (2008). Metabolisme karbohidrat dan pembentukan adenosin tripospat *dalam* Buku ajar fisiologi kedokteran. Jakarta: EGC
- Harianingsih, H., & Wibowo, W. A. (2016). Produksi pati sorgum termodifikasi dengan metode asetilasi. *Majalah Ilmiah MOMENTUM*, 12(1); 26-29 ISSN 0216-7395, e-ISSN 2406-9329.
- Harijono, Estiasih, T., Sunarharum, W. B., & Suwita, I. K. (2012). Efek hipoglikemik polisakarida larut air gembili (*Dioscorea esculenta*) yang diekstrak dengan berbagai metode [Hypoglycaemic Effect of water soluble polysaccharides extracted from gembili (*Dioscorea esculenta*) by various methods]. *Jurnal Teknologi dan Industri Pangan*, 23(1), 1-8.



- Hariyanto B., Purwa. T. Cahyana., Agus. T. Putranto., Sri B. Wahyuningsih dan Y. Marsono. (2017). Penggunaan beras sagu untuk penderita pradiabetes (Use of rice sagu for patients of prediabetes). *Jurnal Pangan*, p-ISSN 0852-0607/ e-ISSN 2527-6239. <http://jurnalpangan.com/index.php/pangan/article/view/361/311>. Artikel.
- Herdianita, N. R., Ong H. L., E. A. Subroto., B. Priadi. (1999). Pengukuran kristalinitas silika berdasarkan difraktometer sinar-X. Proc. ITB, 31(1).
- Hermansson, A. M., & Svegmark, K. (1996). Developments in the understanding of starch functionality. *Trends in Food Science & Technology*, 7(11), 345-353. DOI:10.1016/S0924-2244(96)10036-4
- Hirata, A., Maeda, N., Hiuge, A., Hibuse, T., Fujita, K., Okada, T., & Shimomura, I. (2009). Blockade of mineralocorticoid receptor reverses adipocyte dysfunction and insulin resistance in obese mice. *Cardiovascular research*, 84(1), 164-172. DOI: [10.1093/cvr/cvp191](https://doi.org/10.1093/cvr/cvp191)
- Hizukuri, S. (1986). Polymodal distribution of the chain lengths of amylopectins, and its significance. *Carbohydrate Research*, 147: 342-347. [https://doi.org/10.1016/S0008-6215\(00\)90643-8](https://doi.org/10.1016/S0008-6215(00)90643-8)
- Hoover, Ratnajothi., and T. Vasantha. (1994). Effect of heat-moisture treatment on the structure and physicochemical properties of cereal, legume, and tuber starches. *Carbohydrate Research*, 252: 33-53. [https://doi.org/10.1016/0008-6215\(94\)90004-3](https://doi.org/10.1016/0008-6215(94)90004-3)
- Huang, J., Schols, H. A., Jin, Z., Sulmann, E., & Voragen, A. G. (2007). Pasting properties and (chemical) fine structure of acetylated yellow pea starch is affected by acetylation reagent type and granule size. *Carbohydrate Polymers*, 68(3), 397-406. <https://doi.org/10.1016/j.carbpol.2006.12.019>
- Hustiany, R. (2006). Modifikasi asilasi dan suksinilasi pati tapioka sebagai bahan enkapsulasi komponen flavor. Disertasi, Institut Pertanian Bogor.
- If'all., Hasanuddin, A., Rahim, A., Kadir, S. (2019). Modifikasi pati secara asetilasi terhadap gugus fungsi asetyl dan kristalinitas pati ubi banggai asetat. REKAYASA. Journal of Science and Technology, 12(2): 135-140. <https://doi.org/10.21107/rekayasa.v12i2.5857>
- Imanningsih, N. (2012). Profil gelatinisasi beberapa formulasi tepung-tepungan untuk pendugaan sifat pemasakan (Gelatinisation profile of several flour formulations for estimating cooking behaviour). *Nutrition and Food Research*, 35(1), 13-22.



Institute of Medicine (U.S.) and Subcommittee on Technical Specifications for a High-Energy Emergency Relief Ration. (2002). *High-energy, nutrient-dense emergency relief food product*, ISBN-10: 0-309-08315-X

Jacobs, H., & Delcour, J. A. (1998). Hydrothermal modifications of granular starch, with retention of the granular structure: A review. *Journal of agricultural and food chemistry*, 46(8), 2895-2905.
<https://doi.org/10.1021/jf980169k>

Jatmiko, G. P., Teti, T. (2014). Mie dari umbi kimpul (*Xanthosoma sagittifolium*): Kajian Pustaka (Noodles from cocoyam (*Xanthosoma sagittifolium*)): A Review. *Jurnal Pangan dan Agroindustri*, 2(2): 127-134.
<https://jpa.ub.ac.id/index.php/jpa/article/view/45>

Jyoshna, E., & Hymavathi, T. V. (2017). Review of studies on effect of resistant starch supplementation on glucose and insulin. *J Pharmacog Phytochem*, 6, 55-58.
<https://www.phytojournal.com/archives/2017/vol6issue3/PartA/6-5-189-982.pdf>

Karam, L. B., Grossmann, M. V. E., Silva, R. S. S., Ferrero, C., & Zaritzky, N. E. (2005). Gel textural characteristics of corn, cassava and yam starch blends: a mixture surface response methodology approach. *Starch-Stärke*, 57(2), 62-70. DOI:[10.1002/star.200400328](https://doi.org/10.1002/star.200400328)

Kaur, L., Singh, N., & Sodhi, N. S. (2002). Some properties of potatoes and their starches II. Morphological, thermal and rheological properties of starches. *Food chemistry*, 79(2), 183-192. [http://dx.doi.org/10.1016/S0308-8146\(02\)00130-9](http://dx.doi.org/10.1016/S0308-8146(02)00130-9)

King, G. L., & Loeken, M. R. (2004). Hyperglycemia-induced oxidative stress in diabetic complications. *Histochemistry and cell biology*, 122(4), 333-338. DOI 10.1007/s00418-004-0678-9

Kleessen, B., Stoof, G., Proll, J., Schmiedl, D., Noack, J., & Blaut, M. (1997). Feeding resistant starch affects fecal and cecal microflora and short-chain fatty acids in rats. *Journal of animal science*, 75(9), 2453-2462.
<https://doi.org/10.2527/1997.7592453x>

Kolawole, S. A., Igwemmar, N. C., & Bello, H. A. (2013). Comparison of the physicochemical properties of starch from ginger (*Zingiber officinale*) and maize (*Zea mays*). *International Journal of Science and Research*, 2(11), 71-76.



Kulp K. 1975. Carbohydrate. Didalam G Reed (Ed). Enzymes in Food Processing. Academic Press. New York.

Kumoro, A. C., Amalia, R., Retnowati, D. S., Budiyati, C. S., & Ratnawati, R. (2014). Preparation and physicochemical characterization of modified (acetylated) Gadung (*Dioscorea Hispida Dennst*) flours. *Scientific Study & Research. Chemistry & Chemical Engineering, Biotechnology, Food Industry*, 15(2), 135. <https://dx.doi.org/10.1007%2Fs13197-015-1723-5>

Kurniawati, M., Budijanto, S., & Yuliana, N. D. (2016). Karakterisasi dan indeks glikemik beras analog berbahan dasar tepung jagung. *Jurnal Gizi dan Pangan*, 11(3), 169-174. ISSN 1978-1059 EISSN 2407-0920. Terakreditasi SK Menristek Dikti 12/M/Kp/II/2015. 16450-49483-1-SMLOWGI.pdf

Kusnandar, F. 2010. Kimia Pangan Komponen Mikro. Jakarta. Dian Rakyat.

Kwak, J. H., Paik, J. K., Kim, H. I., Kim, O. Y., Shin, D. Y., Kim, H. J., ... & Lee, J. H. (2012). Dietary treatment with rice containing resistant starch improves markers of endothelial function with reduction of postprandial blood glucose and oxidative stress in patients with prediabetes or newly diagnosed type 2 diabetes. *Atherosclerosis*, 224(2), 457-464. <https://doi.org/10.1016/j.atherosclerosis.2012.08.003>

Lawal, O. S. (2004). Composition, physicochemical properties and retrogradation characteristics of native, oxidised, acetylated and acid-thinned new cocoyam (*Xanthosoma sagittifolium*) starch. *Food chemistry*, 87(2), 205-218. <https://doi.org/10.1016/j.foodchem.2003.11.013>

Lauzon, R. D., Shiraishi, K., Yamazaki, M., Sawayama, S., Sugiyama, N., & Kawabata, A. (1995). Physicochemical properties of cocoyam starch. *Food hydrocolloids*, 9(2), 77-81. 10.1016/s0268-005x(09)80268-3

Lee, H. (2007). *The isolation and characterisation of starches from legume grains and their application in food formulations* (Doctoral dissertation, RMIT University).p. 217

Leclerc, C. J., Champ, M., Boillot, J., Guille, G., Lecannu, G., Molis, C., ... & Galmiche, J. P. (1994). Role of viscous guar gums in lowering the glycemic response after a solid meal. *The American journal of clinical nutrition*, 59(4), 914-921. <https://doi.org/10.1093/ajcn/59.4.914>

Le Thanh-Blicharz, J., Anioła, J., Kowalczewski, P., Przygoński, K., Zaborowska, Z., & Lewandowicz, G. (2014). Type IV resistant starch increases cecum short chain fatty acids level in rats. *Acta Biochimica Polonica*, 61(1): 109-114. https://doi.org/10.18388/abp.2014_1931



Lingga, P. (1995). Bertanam Ubi-ubian. Penebar Swadaya. Jakarta

Liu, Z., L. Peng., and J.F. Kennedy. (2005). The technology of molecular manipulation and modification. Asisted by Microwaves as Applied to Starch Granules. *Carbohydrate Polymers*, 61: 374 - 378. <http://dx.doi.org/10.1016/j.carbpol.2005.06.004>

Liu, H., L. Ramsden, & H. Corke. (1997). Physical Properties and enzymatic digestibility of acetylated ae, wx, and normal Maize Starch. *Carbohydrate Polymer*, 34: 283-289. <https://doi.org/10.1021/jf00113a029>

Masharani,U., German, M. S. (2007). dalam a lange greenspan's basic and clinical endocrinology (8th ed), McGraw Hill Companies, USA. 18:661-747

Marsono, Y. (1998). Perubahan kadar resisten starch (RS) dan komposisi kimia beberapa bahan pangan kaya karbohidrat dalam pengolahan. (Changes of resistant starch (RS) and chemical composition of selected starchy food during processing). *Agritech*, 19(3). <https://doi.org/10.22146/agritech.13729>

Marsono, Y. (2002). Indeks glikemik umbi-umbian (Glycemic index of tubers foods). *Agritech*. 22 (1): 13-16. <https://doi.org/10.22146/agritech.13574>

Marsono, Y. Zuheid Noor, Fitri Rahrnawati. 2003. Pengaruh diet kacang merah terhadap kadar gula darah (Effect of red bean diet on blood glucose concentration of alloxan-induced diabetic rats). *Jurnal.Teknol dan Industri Pangan*. 14 (1).

Marsono, Y. (2012). Zat gizi makro : Karbohidrat. Buku ajar kuliah gizi makro. Program Studi Ilmu dan Teknologi Pangan. SPs UGM. Yogyakarta.

Marsono, Y. (2016). The role and mechanism of resistant starch in reduction plasma glucose concentratiton. Proceeding. International food conference 2016. Innovation of food tecnology to improved food security and health. Okt. 20-2, 2016. Universitas Katolik Widya Mandala. Surabaya.

Marjani, A. (2010). Lipid peroxidation altearions in type 2 diabetic patients. *Pakistan Journal of Biological Science*, 13(15): 723-730. <https://dx.doi.org/10.3923/pjbs.2010.723.730>

Martínez, P., Peña, F., Bello-Pérez, L. A., Núñez-Santiago, C., Yee-Madeira, H., & Velezmoro, C. (2019). Physicochemical, functional and morphological characterization of starches isolated from three native potatoes of the Andean region. *Food chemistry*: X, 2, 100030. <https://doi.org/10.1016/j.fochx.2019.100030>



Maulani, R.R., R. Budiasih., N. Immaningsih. (2012). Karakterisasi fisik dan kimia rimpang dan pati garut (*Marantha arundinaceae l.*) pada berbagai umur panen. Prosiding Seminar Nasional: Kedaulatan Pangan dan Energi Fakultas Pertanian Universitas Trunojoyo Madura.

Maya K., Slamet Budijanto., Nancy Dewi Yuliana. (2015). Karakterisasi dan indeks glikemik beras analog berbahan dasar tepung jagung (Characteritation and glycemic index of rice analog form corn flour). *J. Gizi Pangan*, 3:169-174. ISSN 1978-1059 EISSN 2407-0920. Terakreditasi SK Menristek Dikti 12/M/Kp/II/2015. 16450-49483-1-SMLOWGI.pdf

Mboungeng P. D., D. Tenin., J. Scher., and C. Tchiegang. (2008). Physicochemical and functional properties oof some cultivars of irish potato and cassava starch. *Journal of Food Technology* 6(3): 139-146. <https://medwelljournals.com/abstract/?doi=jftech.2008.139.146>

Mbougueng, P. D., D. Tenin., J. Scher., C. Tchiegang. (2012). Influence of acetylation on phycochemical, functional and thermal properties of potato and cassava starches. *Journal of Food Engineering*. 108: 320-326. <http://dx.doi.org/10.1016/j.jfoodeng.2011.08.006>

Mepba, H. D., Eboh, L., Eko, C. B., & Ukpabi, U. J. (2009). Composition and pasting properties of starch from two cocoyam cultivars. *Journal of food quality*, 32(4), 522-537. <https://doi.org/10.1111/J.1745-4557.2009.00257.X>

Mooradian, A. D. (2009). Dyslipidemia in type 2 diabetes mellitus. *Nature Clinical Practice Endocrinology & Metabolism*. 5: 150-159. <http://dx.doi.org/10.1038/ncpendmet1066>

Muchtadi T. R, Sugiyono. (1992). Petunjuk Laboratorium Ilmu Pengetahuan Bahan Pangan. PAU Pangan dan Gizi. IPB. Bogor

Muhandri, T. (2007). Pengaruh ukuran partikel, kadar padatan, NaCl dan Na²CO³ terhadap sifat amilografi tepung dan pati jagung. *Jurnal Teknologi dan Industri Pangan*, 18(2) : 109-117

Murilo-Chávez C. E., Ya-Jane Wangb., Luis A. Bello-Pérez. (2008). Morphological, physicochemical and structural characteristics of oxidized barley and corn starches. *Starch/Stärk*, 60; 634–645. <http://dx.doi.org/10.1002/star.200800016>

Murtiningrum, M., Lisangan, M. M., & Edoway, Y. (2012). Pengaruh preparasi ubi jalar (ipomoea batatas) sebagai bahan pengental terhadap komposisi kimia dan sifat organoleptik saus buah merah (Pandanus conoideusL). *Agrointek: Jurnal Teknologi Industri Pertanian*, 6(1), 1-7. <https://doi.org/10.21107/agrointek.v6i1.1947>



- Muzaifa M., Muhammad I, S., & Liyuza. (2014). Evaluasi sifat fisik pati ganyong (*Canna edulis* kerr.) sebagai bahan baku pembuatan kwietaiw pada tingkat substitusi yang berbeda. *SAGU*, 13(2) : 35-40. ISSN 1412-44242014.
- Mweta, D. E., Labuschagne, M. T., Bonnet, S., Swarts, J. and Saka, J. D. K. (2010). Isolation and physicochemical characterisation of starch from cocoyam (*Colocasia esculenta*) grown in Malawi. *Journal of the Science of Food and Agriculture*, 90(11): 1886-1896. DOI: [10.1002/jsfa.4029](https://doi.org/10.1002/jsfa.4029)
- Nadia, L., Wirakartakusumah, M. A., Andarwulan, N. dan Purnomo, E. H. (2013). Karakterisasi sifat fisikokimia dan fungsional fraksi pati uwi ungu (*Dioscorea alata*). *Nutrition and Food Research*, 36(2): 91-102.
- Nastiti, A., Murdiati, A., & Marsono, Y. (2017). The effect of autoclaved-cooled jack bean (*canavalia ensiformis* (L.) dc.) high rs-4 starch on lowering glucose level and characteristics of digesta of stz-na induced type-2 diabetes mellitus rats. *Indonesian Food and Nutrition Progress*, 14(2), 77-84. ISSN: 0854-6177
- Ningtyas, N. P. A. (2010). Karakteristik sifat fisikokimia pati jagung termodifikasi dengan proses asetilasi. Skripsi Fakultas Teknologi Industri Universitas Pembangunan Nasional Veteran, Surabaya.
- Noviasari, S., Kusnandar, F., Setiyono, A., & Budijanto, S. (2015). Beras analog sebagai pangan fungsional dengan indeks glikemik rendah. *Jurnal Gizi dan Pangan*, 10(3). <https://doi.org/10.25182/jgp.2015.10.3.%25p>
- Nurani, S., & Yuwono, S. S. (2014). Pemanfaatan tepung kimpul (*xanthosoma sagittifolium*) sebagai bahan baku cookies (kajian proporsi tepung dan penambahan margarin). *Jurnal Pangan dan Agroindustri*, 2(2), 50-58. 37-Article Text-103-1-10-20131210.pdf
- Nurhaeni., Pratiwi, D., Prismawiryanti. (2018). Modifikasi pati sukun (*Artocarpus altilis*) menggunakan anhidrida asam asetat dan aplikasinya pada pembuatan mie [Modified breadfruit (*Artocarpus altilis*) starch using acetic acid anhydride and its application for noodles production]. *KOVALEN*, 4(1): 33-40.
- Nurhidajah, N., N. Nurrahman. (2016). Efek hipoglikemik kecambah beras merah pada tikus yang diinduksi STZ-NA dengan parameter kadar insulin, indeks HOMA-IR dan HOMA β . (Hypoglycemic effect of red rice germ on insulin levels, HOMA-IR and HOMA β index of STZ-NA induced rats). *AGRITECH*, Vol. 36: 433-439. DOI: <http://dx.doi.org/10.22146/agritech.16767>



Nurhidajah dan Nurrahman. (2013). Evaluasi sifat fisik, kimia dan organoleptik kecambah beras merah. *Prosiding Seminar Nasional Menuju Masyarakat Madani dan Lestari*. 549-554. Direktorat Penelitian dan Pengabdian Masyarakat UII, Yogyakarta.

Oates, C. G. (1997). Towards an understanding of starch granule structure and hydrolysis. Review. *Trends Food Sci. Technol*, 8: 375-382.
[https://doi.org/10.1016/S0924-2244\(97\)01090-X](https://doi.org/10.1016/S0924-2244(97)01090-X)

Olaokun, O. O., McGaw, L. J., Janse van Rensburg, I., Eloff, J. N., & Naidoo, V. (2016). Antidiabetic activity of the ethyl acetate fraction of *Ficus lutea* (*Moraceae*) leaf extract: comparison of an in vitro assay with an in vivo obese mouse model. *BMC Complementary and Alternative Medicine*, 16(1), 1-12. DOI 10.1186/s12906-016-1087-z

Olayinka, O. O., Adebawale, K. O., & Olu-Owolabi, I. B. (2013). Physicochemical properties, morphological and X-ray pattern of chemically modified white sorghum starch.(Bicolor-Moench). *Journal of food science and technology*, 50(1), 70-77. DOI 10.1007/s13197-011-0233-3

Olivia Y., Teti Estiasi. (2016). Potensi senyawa bioaktif umbi-umbian lokal sebagai penurun kadar glukosa darah: Kajian Pustaka. *Jurnal Pangan dan Agroindustri*, 4 (1): 410-416.
<https://jpa.ub.ac.id/index.php/jpa/article/view/343/355>

Olu-Owolabi, B. I., Olayinka, O. O., Adegbemile, A. A., & Adebawale, K. O. (2014). Comparison of functional properties between native and chemically modified starches from acha (*Digitaria stapf*) grains. *Food and Nutrition Sciences*, 5:222-230. <http://dx.doi.org/10.4236/fns.2014.52027>

Ononamadu, C. J., Alhassan, A. J., Imam, A. A., Ibrahim, A., Ihegboro, G. O., Owolarafe, A. T., & Sule, M. S. (2019). In vitro and in vivo anti-diabetic and anti-oxidant activities of methanolic leaf extracts of *Ocimum canum*. *Caspian Journal of Internal Medicine*, 10(2), 162.
<http://dx.doi.org/10.22088/cjim.10.2.162>

Parera, E., Purwanto, R. H., & Permadi, D. B. (2022). How do the customary forest and protected forest management principles affect ambon city protected forest management between the government and indigenous peoples (case in the mount sirimau protection forest group, ambon city, maluku province). *Jurnal Manajemen Hutan Tropika*, 28(3), 254-254. DOI: 10.7226/jtfm.28.3.254



- Pigeolet, E., Corbisier, P., Houbion, A., Lambert, D., Michiels, C., Raes, M & Remacle, J. (1990). Glutathione peroxidase, superoxide dismutase, and catalase inactivation by peroxides and oxygen derived free radicals. *Mechanisms of ageing and development*, 51(3), 283-297. Elsevier Scientific Publishers Ireland Ltd.
- Pérez, E., Schultz, F. S., & de Delahaye, E. P. (2005). Characterization of some properties of starches isolated from *Xanthosoma sagittifolium* (tannia) and *Colocassia esculenta* (taro). *Carbohydrate polymers*, 60(2), 139-145. doi:10.1016/j.carbpol.2004.11.033
- Pérez, E. and Lares, M. (2005). Chemical composition, mineral profile, and functional properties of canna (*Canna edulis*) and arrowroot (*Maranta spp.*) Starches. *Plant Foods for Human Nutrition*, 60(3): 113–116. <https://doi.org/10.1007/s11130-005-6838-9>
- Pérez, S., & Bertoft, E. (2010). The molecular structures of starch components and their contribution to the architecture of starch granules: A comprehensive review. *Starch-Stärke*, 62(8), 389-420. DOI 10.1002/star.201000013
- Polnaya, F. J., Haryadi, & D.W, Marseno. (2008). Characteristics of hydroxypropylated and acetylated sago starches. *Sago Palm*, 16: 85-94.
- Polnaya, F. J., B.R.I. Puturuhi. (2010). Preparasi dan karakterisasi pati sagu asetil. *Buletin Penelitian Biam*, 6: 10-13.
- Polnaya, F. J., Hilda., Cynthia, G. C. L. (2020). Konsentrasi asam asetat mempengaruhi karakteristik fisikokimia pati sagu ihur terasetilasi. *J. teknol. Dan Industri Pangan*, 31(2); 180-187. DOI: 10.6066/jtip.2020.31.2.180
- Puddu, A., Sanguineti, R., Montecucco, F., & Viviani, G. L. (2014). Evidence for the gut microbiota short-chain fatty acids as key pathophysiological molecules improving diabetes. *Mediators of inflammation*, 2014.
- Raina, C., S. Singh, A. Bawa, and D. Saxena. (2006). Some characteristics of acetylated, cross-linked and dual modified Indian rice starches. *European Food Research and Technology*. 223: 561-570. <http://dx.doi.org/10.1007/s00217-005-0239-z>
- Radley, J. A. (1976). Starch production technology. *Applied science publisher*, Ltd. London.
- Rahmawati, W., Kusumastuti, Y. A., & Aryanti, N. (2012). Karakterisasi pati talas (*Colocasia esculenta* (L.) schott) sebagai alternatif sumber pati industri



di Indonesia. *Jurnal Teknologi Kimia dan Industri*, 1(1), 347-351.
<http://ejournal-s1.undip.ac.id/index.php/jtki>

Richana, N., & Sunarti, T. C. (2004). Karakterisasi sifat fisikokimia tepung umbi dan tepung pati dari umbi ganyong, suweg, ubikelapa dan gembili. *Jurnal pascapanen*, 1(1): 29-37.

Ridal, S. (2003). Karakterisasi sifat fisiko-kimia tepung dan pati talas (*Colocasia esculenta*) dan kimpul (*Xanthosoma sp.*) dan uji penerimaan alfa amilase terhadap patinya. Skripsi. Jurusan Teknologi Industri Pertanian IPB. Bogor

Riyanti E. (2018). Pati resisten pada beras : jenis, metode peningkatan, efek untuk kesehatan, dan aplikasinya *resistant starch on rice : types, enhancement methods, health effects, and its applications*. *Jurnal Pangan*. 26 (3): 11-15

Robertson, M. D., Currie, J. M., Morgan, L. M., Jewell, D. P., & Frayn, K. N. (2003). Prior short-term consumption of resistant starch enhances postprandial insulin sensitivity in healthy subjects. *Diabetologia*, 46(5), 659-665. <https://doi.org/10.1007/s00125-003-1081-0>

Sacks, D.B. (2001). *Carbohydrates*, in *Tietz Fundamentals of Clinical chemistry*, 5nd W.B. Saunders Company, USA

Sajilata, M. G., Singhal, R. S., & Kulkarni, P. R. (2006). Resistant starch—a review. *Comprehensive reviews in food science and food safety*, 5(1), 1-17. <https://doi.org/10.1111/j.1541-4337.2006.tb00076.x>

Sánchez-Rivera, M. M., Almanza-Benitez, S., Bello-Perez, L. A., Mendez-Monrealvo, G., Núñez-Santiago, M. C., Rodriguez-Ambriz, S. L., & Gutierrez-Meráz, F. (2013). Acetylation of banana (*Musa paradisiaca* L.) and corn (*Zea mays* L.) starches using a microwave heating procedure and iodine as catalyst: II. Rheological and structural studies. *Carbohydrate polymers*, 92(2), 1256-1261. <https://doi.org/10.1016/j.carbpol.2012.10.040>

Saputro, M. A., A. Kurniawan., dan D. S. Retnowati. (2012). Modifikasi pati talas dengan asetilasi menggunakan asam asetat. *Jurnal Teknologi Kimia dan Industri* 1: 258-263. <http://ejournal-s1.undip.ac.id/index.php/jtki>

Sarah A. Dainty.(2017). The effect of resistant starch bagels on glycemic response in adults at risk for type 2 diabetes. A Thesis presented to The University of Guelph, Ontario, Canada. https://atrium.lib.uoguelph.ca/xmlui/bitstream/handle/10214/8790/Dainty_Sarah_201505_MSc.pdf?sequence=7&isAllowed=y

Seib, P. A and K. Woo. (1999). Food grade starch resistant to alpha amilase and method of preparing the same. Kansas State University Research



Foundation. US Patent 5: 855-946.
<https://patents.google.com/patent/US5855946A/en>

Sha, X. S., Xiang, Z. J., Bin, L., Jing, L., Bin, Z., Jiao, Y. J. and Kun, S. R. (2012). Preparation and physical characteristics of resistant starch (type 4) in acetylated indica rice. *Food Chemistry*, 134(1): 149–154. <https://doi.org/10.1016/j.foodchem.2012.02.081>

Shah, A., Masoodi, F. A., Gani, A., & Ashwar, B. A. (2017). Physicochemical, rheological and structural characterization of acetylated oat starches. *LWT*, 80, 19-26. <http://dx.doi.org/10.1016%2Fj.lwt.2017.01.072>

Shanise L. M. El Halal., Colussi, R., Vania Z, P., Josiane B., Marjiana R., Neftali L,V,C., Alvaro,R, G, D & Elessandra da R, Z. (2015). Stucture, morphology and functionality of acetylated and oxidised barley starches. *Food Chemistry*, 168: 247-256. <https://doi.org/10.1016/j.foodchem.2014.07.046>

Sharma, A., Yadav, B. S., & Ritika. (2008). Resistant starch: physiological roles and food applications. *Food reviews international*, 24(2), 193-234.

Shi, Y. C., Cui, X., Birkett, A. M., & Thatcher, M. G. (2006). U.S. Patent No. 7,081,261. Washington, DC: U.S. Patent and Trademark Office.

Shi, Y. C., Cui, X., Birkett, A.M., & Thatcher, M G. (2006). Resistant starch Prepared by isoamylase debranching of low amylose st arch . US Pat ent Office, Pat No. 7081261. <https://www.freepatentsonline.com/7081261.pdf>

Shimelis, E. A., Meaza, M., & Rakshit, S. (2006). Physico-chemical properties, pasting behavior and functional characteristics of flours and starches from improved bean (*Phaseolus vulgaris L.*) varieties grown in East Africa. *Agricultural Engineering International: the CIGR Ejournal*. Manuscript FP 05 015. 8. <https://hdl.handle.net/1813/10533>

Shimotoyodome, A., Suzuki, J., Fukuoka, D., Tokimitsu, I., & Hase, T. (2010). RS4-type resistant starch prevents high-fat diet-induced obesity via increased hepatic fatty acid oxidation and decreased postprandial GIP in C57BL/6J mice. *American Journal of Physiology-Endocrinology and Metabolism*, 298(3), E652-E662. <https://doi.org/10.1152/ajpendo.00468.2009>

Shogren, R. L., & Biswas, A. (2010). Acetylation of starch with vinyl acetate in imidazolium ionic liquids and characterization of acetate distribution. *Carbohydrate Polymers*, 81(1), 149-151. <https://doi.org/10.1016/j.carbpol.2010.01.045>



Sichard Elfrits Mual. 2015. Analisis Sifat Fisikokimia Tepung keladi (*Xanthosoma sagittifolium*) Asal Maluku. Skripsi. Jurusan teknologi Hasil Pertanian. Universitas Pattimura. Ambon

Silva, G. M. D. S., Veloso, C. M., Santos, L. S., Melo Neto, B. A. D., Fontan, R. D. C. I., & Bonomo, R. C. F. (2020). Extraction and characterization of native starch obtained from the inhambu tuber. *Journal of food science and technology*, 57(5), 1830-1839. <https://doi.org/10.1007/s13197-019-04216-4>

Simsek, S., Ovando-Martínez, M., Whitney, K., & Bello-Pérez, L. A. (2012). Effect of acetylation, oxidation and annealing on physicochemical properties of bean starch. *Food Chemistry*, 134(4), 1796-1803. <https://doi.org/10.1016/j.foodchem.2012.03.078>

Singh, N., Singh, J., Kaur, L., Sodhi, N. S., & Gill, B. S. (2003). Morphological, thermal and rheological properties of starches from different botanical sources. *Food chemistry*, 81(2), 219-231. [https://doi.org/10.1016/S0308-8146\(02\)00416-8](https://doi.org/10.1016/S0308-8146(02)00416-8)

Singh, N., Chawla, D., & Singh, J. (2004). Influence of acetic anhydride on physicochemical, morphological and thermal properties of corn and potato starch. *Food Chemistry*, 86(4), 601-608. <https://doi.org/10.1016/j.foodchem.2003.10.008>

Sodhi, N. S., & Singh, N. (2005). Characteristics of acetylated starches prepared using starches separated from different rice cultivars. *Journal of Food Engineering*, 70(1), 117-127. <https://doi.org/10.1080/10942912.2010.483633>

Singh, H., Sodhi, N. S., & Singh, N. (2012). Structure and functional properties of acetylated sorghum starch. *International Journal of Food Properties*, 15(2), 312-325. <https://doi.org/10.1080/10942912.2010.483633>

Song, X., He, G., Ruan, H., & Chen, Q. (2006). Preparation and properties of octenyl succinic anhydride modified early indica rice starch. *Starch-Stärke*, 58(2), 109-117. <https://doi.org/10.1002/star.200500444>

Srichuwong, S., Sunarti, T. C., Mishima, T., Isono, N., & Hisamatsu, M. (2005). Starches from different botanical sources I: Contribution of amylopectin fine structure to thermal properties and enzyme digestibility. *Carbohydrate polymers*, 60(4), 529-538. <http://dx.doi.org/10.1016/j.carbpol.2005.03.004>

Sunarti. (2017). Serat pangan dalam penanganan sindrom metabolik. Gadjah Mada University Press. Yogyakarta.

Sutrisno K. 2009. Teknologi Modifikasi Pati. EbookPangan.com



- Tachibe, M., Kato, R., Sugano, S., Kishida, T., & Ebihara, K. (2009). Hydroxypropylated tapioca starch retards the development of insulin resistance in KKAY mice, a type 2 diabetes model, fed a high-fat diet. *Journal of food science*, 74(7), H232-H236. <https://doi.org/10.1111/j.1750-3841.2009.01276.x>
- Tan Kim H. (1998). Dasar - dasar kimia tanah. Gadjah Mada University Press. Yogyakarta
- Teja, A.W., I. P. Sindi, A. Ayucitra., and L. E. K. Setiawan. (2008). Karakteristik pati sagu dengan metode asetilasi dan cross-linking. *Jurnal Teknik Kimia Indonesia*. 7: 836-843. <https://www.aptekim.id/jtki/index.php/JTKI/article/viewFile/141/135>
- Tester, R. F., Morrison, W. R., & Schulman, A. H. (1993). Swelling and gelatinization of cereal starches. V. Risø mutants of Bomi and Carlsberg II barley cultivars. *Journal of Cereal Science*, 17(1), 1-9. <https://doi.org/10.1006/jcrs.1993.1001>
- Tijani, A. O., Ibitayo Omohimi, C., Sanni, L. O., & Oke, E. K. (2016). Physicochemical properties of food grade acetylated cocoyam (*Xanthosoma sagittifolium*) starches. *Croatian journal of food science and technology*, 8(2), 53-59. <http://dx.doi.org/10.17508/CJFST.2016.8.2.02>
- Topping, D. L., Clifton. P. (2001). Short chain fatty acid and human colonic function: role of resistant starch and non-starch polysaccharides. *Phys. Rev*, 81: 1031-1064. <https://doi.org/10.1152/physrev.2001.81.3.1031>
- Triwitono, P., Marsono, Y., Murdiati, A., & Marseno, D. W. (2017). Isolasi dan karakterisasi sifat pati kacang hijau (*Vigna radiata L.*) beberapa varietas lokal Indonesia. *Agritech*, 37(2), 192-198. <http://dx.doi.org/10.22146/agritech.10659>.
- Ulfa, M., Sakinah, E. N., Sudarmanto, Y., & Firdaus, J. (2020). Efek Pati Resisten Singkong (*Manihot esculenta Crantz*) terhadap Indeks HOMA-IR dan HOMA-B Tikus Model Diabetes. *Pustaka Kesehatan*, 7(1), 1-7. *Journal Pustaka Kesehatan*, p-ISSN: 2355-178x, e-ISSN: 2721-3218 7(1): 1-7. <https://doi.org/10.19184/pk.v7i1.17586>
- Valko, M., Leibfritz, D., Moncol, J., Cronin, M. T., Mazur, M., & Telser, J. (2007). Free radicals and antioxidants in normal physiological functions and human disease. *The international journal of biochemistry & cell biology*, 39(1), 44-84.



- Varavinit, S., Shobsngob, S., Varanyanond, W., Chinachoti, P., & Naivikul, O. (2003). Effect of amylose content on gelatinization, retrogradation and pasting properties of flours from different cultivars of Thai rice. *Starch-Stärke*, 55(9), 410-415. <https://doi.org/10.1002/star.200300185>.
- Villanueva-Peña Carrillo, M. L., Puente, J., Redondo, A., Clemente, F., & Valverde, I. (2001). Effect of GLP-1 treatment on GLUT2 and GLUT4 expression in type 1 and type 2 rat diabetic models. *Endocrine*, 15(2), 241-248. <https://doi.org/10.1385/ENDO:15:2:24>
- Wahjuningsih, S. B., Marsono, Y., Praseptiangga, D., & Haryanto, B. (2016). Resistant starch content and glycaemic index of sago (*Metroxylon* spp.) starch and red bean (*Phaseolus vulgaris*) based analogue rice. *Pak J Nutr*, 15(7), 667-672. . <https://dx.doi.org/10.3923/pjn.2016.667.672>
- Wahjuningsih, S. B., Untari, S., & Wijanarka, A. (2017). Hypoglycemic effect of analog rice made from modified cassava flour (Mocaf), arrowroot flour and kidney bean flour on STZ-NA induced diabetic rats. *Asian Journal of Clinical Nutrition*, 10(1), 8-15. <https://dx.doi.org/10.3923/ajcn.2018.8.15>
- Wahjuningsih, S. B., Haslina, H., & Putranto, A. T. (2020). Effect of Sago Analogue Rice and Red Bean Diet to the Improvement of β -cell Pancreas in Streptozotocin-Nicotinamide (STZ-NA) Induced Diabetic Rats. *Current Research in Nutrition and Food Science*, 8(2), 667. <http://dx.doi.org/10.12944/CRNFSJ.8.2.32>
- Waluszewski, K.N., Aparicio, M.A., Bello, L.A., Monroy, J.A. (2003). Changes of banana starch by chemical and physical modification. *Carbohydrate Polymers*, 52: 237–242. [https://doi.org/10.1016/S0144-8617\(02\)00270-9](https://doi.org/10.1016/S0144-8617(02)00270-9)
- Waterschoot, J., Gomand, S.V., Fierens, E. and Delcour, J.A. (2014). Production, structure, physicochemical and functional properties of maize, cassava, wheat, potato and rice starches. *Starch Stärk*, 66: 1–16. <https://doi.org/10.1002/star.201300238>
- Wattanachant, S., Muhammad, S. K. S., Mat Hashim, D., & Rahman, R. A. (2002). Suitability of sago starch as a base for dual-modification. *Songklanakarin J. Sci. Technol*, 24(3), 431-438.
- Weickert, M. O., and Mohlig, M. (2005). Impact of cereal fibre on glucose-regulating factors. *Diabetologia*, 48: 2343–2353. <https://doi.org/10.1007/s00125-005-1941-x>



Winarno, F. 2008. Kimia pangan dan gizi: Edisi Terbaru. Jakarta. *Gramedia Pustaka Utama*

Wronkowska, M., Juskiewicz, J., Zdunczyk, Z., Soral-Smietana, M., & Krupa-Kozak, U. (2011). Influence of chemically-modified potato starch (RS type 4) on the nutritional and physiological indices of rats. *Polish journal of food and nutrition sciences*, 61(2). <https://doi.org/10.2478/v10222-011-0015-7>

Wu, K. K., and Huan, Y. (2008). Streptozotocin-induced diabetic models in mice and rats. *Current Protocols in Pharmacology*, 5(4): 5.47.1-5.47.14. doi:10.1002/0471141755. ph0547s40. <https://doi.org/10.1002/0471141755.ph0547s40>

Wurzburg, O.B. (1989). Modified starches: properties and uses. CRC Press, Boca Raton, Florida.

Xu, Y., V. Miladinov., and M. A. Hanna. (2004). Synthesis and characterization of starch acetates with high substitution *Cereal Chem. J.* 81(6): 735–40. <http://dx.doi.org/10.1094/CCHEM.2004.81.6.735>.

Xu, Y. (2013) Resistant starch content, molecular structure and physicochemical properties of starches in Virginia-grown corn, potato and mungbean *J. Cereals Oilseed.* 4(1): 10–18. <http://dx.doi.org/10.5897/JCO2012.0097>

Xu, S., Sha, Zhang J., Xiang, Li Bin., Li Jing., Zhao Bin., Yan J. Jiao., Song R. Kun. (2012). Preparation and physical characteristics of resistant starch (type 4) in acetylated indica rice. *Food Chemistry*, 134: 149-154. DOI:[10.1016/J.FOODCHEM.2012.02.081](https://doi.org/10.1016/J.FOODCHEM.2012.02.081)

Yalçın T., Ayhan Al., Neslişah Rakıcıoğlu., (2017). The effects of meal glycemic load on blood glucose levels of adults with different body mass indexes. *Indian Journal of Endocrinology and Metabolism*, 21(1):71-75. <https://doi.org/10.4103/2230-8210.195995>

Zhou, Y., Meng, S., Chen, D., Zhu, X., & Yuan, H. (2014). Structure characterization and hypoglycemic effects of dual modified resistant starch from indica rice starch. *Carbohydrate polymers*, 103, 81-86. <https://doi.org/10.1016/j.carbpol.2013.12.020>

Zhou, Z., Wang, F., Ren, X., Wang, Y., & Blanchard, C. (2015). Resistant starch manipulated hyperglycemia/hyperlipidemia and related genes expression in diabetic rats. *International journal of biological macromolecules*, 75, 316-321. <https://doi.org/10.1016/j.ijbiomac.2015.01.052>

Zhou, F., Liu, Q., Zhang, H., Chen, Q., & Kong, B. (2016). Potato starch oxidation induced by sodium hypochlorite and its effect on functional



properties and digestibility. *International journal of biological macromolecules*, 84, 410-417. <https://doi.org/10.1016/j.ijbiomac.2015.12.050>

Zięba, T., Kapelko, M., & Szumny, A. (2013). Effect of preparation method on the properties of potato starch acetates with an equal degree of substitution. *Carbohydrate polymers*, 94(1), 193-198. .
<https://doi.org/10.1016/j.carbpol.2013.01.002>