UNIVERSITAS GADJAH MADA
PENGGUNAAN KOMBINASI UPPER LIP BITE TEST (ULBT), RATIO HEIGHT TO THYROMENTAL DISTANCE (RHTMD) DAN EXTENDED MALLAMPATI SCORE (EMS) SEBAGAI PREDIKTOR SULIT LARINGOSKOPI
DANUR ADI KUSUMA, pembimbing materi dr. Bowo Adiyanto Sp.An, M.Sc, KIC Departemen Anestesiologi dan Tera
Universitas Gadjah Mada, 2023 | Diunduh dari http://etd.repository.ugm.ac.id/

DAFTAR PUSTAKA

Alanzi, A. (2015) 'Intubations and airway management: An overview of Hassles through third millennium', *J Emerg Trauma Shock*, 8(2).

Amelis, G., Budipratama, D. and Oktaliansah, E. (2019) 'Perbandingan Angka Keberhasilan Intubasi dan Waktu Intubasi antara Menggunakan Bougie dan Purwarupa Camera–Bougie pada Maneken Simulasi Kesulitan Intubasi', *Jurnal Anestesi Perioperatif*, 9(7), pp. 18–24.

Apfelbaum, J. L., Hagberg, C. A., Connis, R. T., Abdelmalak, B. B., Agarkar, M., Dutton, R. P., & Tung, A. (2021). 2022 American Society of Anesthesiologists Practice Guidelines for Management of the Difficult Airway. *Anesthesiology*.

Basunia, S. R., Ghosh, S., Bhattacharya, S., Saha, I., Biswas, A., & Prasad, A. (2013). Comparison between different tests and their combination for prediction of difficult intubation: An analytical study. *Anesthesia, essays and researches*, *7*(1), 105.

Butterworth, J., Mackey, D. and Wasnick, J. (2018) *Morgan & Mikhail's Clinical Anesthesiology*. 6th edn. London: McGraw Hill.

Domi, R. 2009. A comparison of Wilson sum score and combination Mallampati, tiromental and sternomental distances for predicting difficult intubation. *Maced J Med Sci*, *2*(2), 141-4.

Erhenwerth, J., Eisenkraft, A. and Berry, J. (2013) *Anesthesia Equipment: Principles and Applications*. 5th edn. London: Saunders.

Faramarzi, E., Soleimanpour, H., Khan, Z. H., Mahmoodpoor, A., & Sanaie, S. (2018). Upper lip bite test for prediction of difficult airway: a systematic review. *Pakistan journal of medical sciences*, *34*(4), 1019.

George, R. B., McKeen, D., & Law, J. A. 2006. Laryngoscopic evaluation with the Airway Cam. *Canadian journal of anaesthesia = Journal canadien d'anesthesie*, *53*(5), pp. 512–515.

Goto, Y., Goto, T., Okamoto, H., Hagiwara, Y., Watase, H., Hasegawa, K., & Inoue, A. (2020). Factors associated with successful rescue intubation attempts in the emergency department: an analysis of multicenter prospective observational study in Japan. *Acute medicine & surgery*, *7*(1), e462.

Hagberg, C. (2018) *Hagberg and Benumof's Airway Management, Fourth Edition*. Elsevier.

Honarmand, A., Amoushahi, M. and Safavi, M. (2014) 'Prediction of difficult laryngoscopy: Extended mallampati score versus the MMT, ULBT and RHTMD', *Advanced Biomedical Research*, 3(1), p. 133. doi: 10.4103/2277-9175.133270.

PENGGUNAAN KOMBINASI UPPER LIP BITE TEST (ULBT), RATIO HEIGHT TO THYROMENTAL DISTANCE (RHTMD) DAN
EXTENDED MALLAMPATI SCORE (EMS) SEBAGAI PREDIKTOR SULIT LARINGOSKOPI
DANUR ADI KUSUMA, pembimbing materi dr. Bowo Adiyanto Sp.An, M.Sc, KIC Departemen Anestesiologi dan Tera
Universitas Gadjah Mada, 2023 | Diunduh dari http://etd.repository.ugm.ac.id/

Jagannatha, M. and Prabhudev, K. 2015. 'Ratio of Patients Height To Thyromental Distance (Rhtmd) Compared To Thyromental Distance for Prediction of Difficult Intubation', *Journal of Evidence Based Medicine and Healthcare*, 2(42), pp. 7462–7476. doi: 10.18410/jebmh/2015/1009.

Janssens, M., & Lamy, M. 2000. Airway Difficulty Score (ADS): a new score to predict difficulty in airway management. *European Journal of Anaesthesiology (EJA)*, *17*, 35.

Kaniyil, S., Anandan, K. and Thomas, S. (2018) 'Ratio of height to thyromental distance as a predictor of difficult laryngoscopy: A prospective observational study', *Journal of Anaesthesiology Clinical Pharmacology*, 34(4), p. 485. doi: 10.4103/joacp.JOACP_283_17.

Khan, Z. H., Kashfi, A., & Ebrahimkhani, E. 2003. A comparison of the upper lip bite test (a simple new technique) with modified Mallampati classification in predicting difficulty in endotracheal intubation: a prospective blinded study. *Anesthesia and analgesia*, *96*(2)

Krage, R., Van Rijn, C., Van Groeningen, D., Loer, S. A., Schwarte, L. A., & Schober, P. (2010). Cormack–Lehane classification revisited. *British journal of anaesthesia*, *105*(2), 220-227.

Krobbuaban, B., Diregpoke, S., Kumkeaw, S., & Tanomsat, M. (2005). The predictive value of the height ratio and thyromental distance: four predictive tests for difficult laryngoscopy. *Anesthesia & Analgesia*, *101*(5), 1542-1545.

Levitan, R. M., Ochroch, E. A., Kush, S., Shofer, F. S., & Hollander, J. E. 1998. Assessment of airway visualization: validation of the percentage of glottic opening (POGO) scale. *Academic emergency medicine : official journal of the Society for Academic Emergency Medicine*, *5*(9), pp. 919–923

Mallampati, S. R., Gatt, S. P., Gugino, L. D., Desai, S. P., Waraksa, B., Freiberger, D., & Liu, P. L. 1985. A clinical sign to predict difficult tracheal intubation: a prospective study. *Canadian Anaesthetists' Society journal*, *32*(4), pp. 429–434

Mashour, G. A., & Sandberg, W. S. (2006). Craniocervical extension improves the specificity and predictive value of the Mallampati airway evaluation. *Anesthesia & Analgesia*, *103*(5), 1256-1259.

Mononyane, R. (2018) 'Regional anaesthesia for awake intubation', *Southern African Journal of Anaesthesia and Analgesia*, 24(3), pp. S103–S105.

PENGGUNAAN KOMBINASI UPPER LIP BITE TEST (ULBT), RATIO HEIGHT TO THYROMENTAL DISTANCE (RHTMD) DAN
EXTENDED MALLAMPATI SCORE (EMS) SEBAGAI PREDIKTOR SULIT LARINGOSKOPI
DANUR ADI KUSUMA, pembimbing materi dr. Bowo Adiyanto Sp.An, M.Sc, KIC Departemen Anestesiologi dan Tera
Universitas Gadjah Mada, 2023 | Diunduh dari http://etd.repository.ugm.ac.id/

Munsy, A. K., Rehatta, N. M., Maulydia, M., Salinding, A., Utariani, A., Sylvaranto, T., & Hanindito, E. Perbandingan Visualisasi Laring dan Glotis pada Maneken Intubasi Sulit menggunakan Video Laryngoscope C-MAC dan VL-Scope. *JAI (Jurnal Anestesiologi Indonesia)*, *10*(3), 143-152.

Nuckton, T. J., Glidden, D. V., Browner, W. S., & Claman, D. M. (2006). Physical examination: Mallampati score as an independent predictor of obstructive sleep apnea. *Sleep*, *29*(7), 903-908.

Oates, J. D. L., Macleod, A. D., Oates, P. D., Pearsall, F. J., Howie, J. C., & Murray, G. D. (1991). Comparison of two methods for predicting difficult intubation. *British Journal of Anaesthesia*, *66*(3), 305-309.

Oria, M. S., Halimi, S. A., Negin, F., & Asady, A. (2022). Predisposing Factors of Difficult Tracheal Intubation Among Adult Patients in Aliabad Teaching Hospital in Kabul, Afghanistan–A Prospective Observational Study. *International Journal of General Medicine*, *15*, 1161.

Panjiar, P., Bhat, KM., Yousuf, I., Kochhar, A., Ralli, T., 2021. Study comparing different airway assessment tests in predicting difficult laryngoscopy: A prospective study in geriatric patients. Indian J Anaesth, 65(4), pp. 309-315

Parillo, J. and Dellinger, P. (2019) *Critical Care Medicine: Principles of Diagnosis and Management in the Adult*. 5th edn. London: Elsevier.

Patel J., Patel D., Vaidya V., Chauhan J., et al. 2020. Comparison of the Upper Lip Bite Test (ULBT) with the Ratio of Height to Thyromental Distance (RHTMD) for the Prediction of Difficult Laryngoscopy in Apparently Normal Patients. Indian Journal of Anesthesia and Analgesia.

Philip, S., and Nizar, S., 2016. Prediction of difficult laryngoscopy in patients undergoing endotracheal intubation: A comparative study of various airway assessment tests. *Astrocyte* 3 pp. 90-95

Safavi, M., Honarmand, A., & Amoushahi, M. (2014). Prediction of difficult laryngoscopy: Extended mallampati score versus the MMT, ULBT and RHTMD. *Advanced biomedical research*, p133.

Savva D. 1994. Prediction of difficult tracheal intubation. *British journal of anaesthesia*, *73*(2), pp. 149–153.

Simatupang, R. 2014. Korelasi Upper Lip Bite Test (ULBT) Dengan Mallampati Sebagai Prediktor Kesulitan Intubasi Di Rumah Sakit Sanglah Denpasar. Tesis. Denpasar. Universitas

PENGGUNAAN KOMBINASI UPPER LIP BITE TEST (ULBT), RATIO HEIGHT TO THYROMENTAL DISTANCE (RHTMD) DAN
EXTENDED MALLAMPATI SCORE (EMS) SEBAGAI PREDIKTOR SULIT LARINGOSKOPI
DANUR ADI KUSUMA, pembimbing materi dr. Bowo Adiyanto Sp.An, M.Sc, KIC Departemen Anestesiologi dan Tera
Universitas Gadjah Mada, 2023 | Diunduh dari http://etd.repository.ugm.ac.id/

Udayana.

Standring, S. (2020) *Gray's Anatomy : The Anatomical Basis of Clinical Practice*. 42nd edn. San Fransisco: Elsevier.

Swasono, G., Suwarman and Kadarsah, R. (2017) 'Perbandingan antara Uji Mallampati Modifikasi dan Mallampati Ekstensi Sebagai Prediktor Kesulitan Intubasi Endotrakeal di RSUP Dr. Hasan Sadikin Bandung', *Jurnal Anestesi*, 5(3), pp. 163–70.

Tayyeb, M., Faisal, S., Abdullah, A. M., Ahmad, N., & Arsalan, M. (2020). Prediction of Difficult Intubation with the Modified Mallampati and Upper Lip Bite Test: A Cross-Sectional Analysis. *J Biomed Sci*, *9*(2), 5.

Thankamony, D. H. (2019) 'Diagnostic Value of Upper Lip Bite Test combined with Hyomental Distance Ratio in assessing the airway', *Journal of Current Medical Research and Opinion*, 02(12), pp. 383–387. doi: 10.15520/jcmro.v2i12.241.

Tompodung, S., Tambajong, H. F., & Lalenoh, D. (2013). MULA KERJA ROKURONIUM. *JURNAL BIOMEDIK: JBM*, *5*(1).

Wardhana, Y. K. (2009). Penggunaan kriteria mallampati, jarak tiromental dan gigit bibir atas sebagai prediktor kombinasi pada kesulitan intubasi endotrakea. Tesis. Yogyakarta. Universitas Gadjah Mada