



- [1] S.-H. Kong, S.-K. Loh, R. T. Bachmann, S. A. Rahim, and J. Salimon, "Biochar from oil palm biomass: A review of its potential and challenges," *Renew. Sustain. Energy Rev.*, vol. 39, pp. 729–739, Nov. 2014, doi: 10.1016/j.rser.2014.07.107.
- [2] K. Y. Phang and S. W. Lau, "A Survey on the Usage of Biomass Wastes from Palm Oil Mills on Sustainable Development of Oil Palm Plantations in Sarawak," *IOP Conf. Ser. Mater. Sci. Eng.*, vol. 206, no. 1, 2017, doi: 10.1088/1757-899X/206/1/012091.
- [3] M. N. F. Norrrahim, M. A. A. Farid, A. Abdullahi Lawal, T. A. T. Yasim-Anuar, M. Hafif Samsudin, and A. Aiman Zulkifli, "Emerging technologies for value-added use of oil palm biomass," *Environ. Sci. Adv.*, vol. 1, no. 3, pp. 259–275, 2022, doi: 10.1039/D2VA00029F.
- [4] R. Nabila *et al.*, "Oil palm biomass in Indonesia: Thermochemical upgrading and its utilization," *Renew. Sustain. Energy Rev.*, vol. 176, p. 113193, Apr. 2023, doi: 10.1016/j.rser.2023.113193.
- [5] M. A. Sukiran *et al.*, "Experimental and modeling study of the torrefaction of empty fruit bunches as a potential fuel for palm oil mill boilers," *Biomass Bioenergy*, vol. 136, 2020, doi: 10.1016/j.biombioe.2020.105530.
- [6] S. Saelor, P. Kongjan, and S. O-Thong, "Biogas Production from Anaerobic Co-digestion of Palm Oil Mill Effluent and Empty Fruit Bunches," *Energy Procedia*, vol. 138, pp. 717–722, Oct. 2017, doi: 10.1016/j.egypro.2017.10.206.
- [7] P. Wadchasit, W. Suksong, S. O-Thong, and K. Nuithitikul, "Development of a novel reactor for simultaneous production of biogas from oil-palm empty fruit bunches (EFB) and palm oil mill effluents (POME)," *J. Environ. Chem. Eng.*, vol. 9, no. 3, p. 105209, Jun. 2021, doi: 10.1016/j.jece.2021.105209.
- [8] S. Kaniapan, S. Hassan, H. Ya, K. Patma Nesan, and M. Azeem, "The Utilisation of Palm Oil and Oil Palm Residues and the Related Challenges as a Sustainable Alternative in Biofuel, Bioenergy, and Transportation Sector: A Review," *Sustainability*, vol. 13, no. 6, Art. no. 6, Jan. 2021, doi: 10.3390/su13063110.
- [9] S. Novianti, A. Nurdiawati, I. N. Zaini, P. Prawisudha, H. Sumida, and K. Yoshikawa, "Low-potassium Fuel Production from Empty Fruit Bunches by Hydrothermal Treatment Processing and Water Leaching," *Energy Procedia*, vol. 75, pp. 584–589, Aug. 2015, doi: 10.1016/j.egypro.2015.07.460.
- [10] A. Nurdiawati, S. Novianti, I. N. Zaini, H. Sumida, and K. Yoshikawa, "Production of Low-Potassium Solid Fuel from Empty Fruit Bunches (EFB) by Employing Hydrothermal Treatment and Water Washing Process," *J. Jpn. Inst. Energy*, vol. 94, no. 8, pp. 775–780, 2015, doi: 10.3775/jie.94.775.
- [11] A. Haryanto, D. A. Iryani, U. Hasanudin, M. Telaumbanua, S. Triyono, and W. Hidayat, "Biomass Fuel from Oil Palm Empty Fruit Bunch Pellet: Potential and Challenges," *Procedia Environ. Sci. Eng. Manag.*, vol. 8, no. 1, Art. no. 1, 2021, Accessed: Mar. 07, 2023. [Online]. Available: <http://www.procedia-esem.eu/index.html>
- [12] Agency for Natural Resources and Energy, "Feed-in Tariff Scheme for Renewable Energy," Oct. 2011. [https://www.meti.go.jp/english/policy/energy\\_environment/renewable/pdf/summary201209.pdf](https://www.meti.go.jp/english/policy/energy_environment/renewable/pdf/summary201209.pdf)



- [13] “The 83rd meeting of the Procurement Price Calculation Committee (in Japanese),” *Ministry of Economy, Trading and Industry*, Jan. 17, 2023. [https://www.meti.go.jp/shingikai/santeii/pdf/083\\_01\\_00.pdf](https://www.meti.go.jp/shingikai/santeii/pdf/083_01_00.pdf) (accessed Mar. 07, 2023).
- [14] S. S. Harsono *et al.*, “Energy balances, greenhouse gas emissions and economics of biochar production from palm oil empty fruit bunches,” *Resour. Conserv. Recycl.*, vol. 77, pp. 108–115, Aug. 2013, doi: 10.1016/j.resconrec.2013.04.005.
- [15] Y. Krishnan *et al.*, “Co-composting of palm empty fruit bunch and palm oil mill effluent: Microbial diversity and potential mitigation of greenhouse gas emission,” *J. Clean. Prod.*, vol. 146, pp. 94–100, Mar. 2017, doi: 10.1016/j.jclepro.2016.08.118.
- [16] Z. L. Chew, E. H. Tan, S. A. L. Palaniandy, K. S. Woon, and Z. X. Phuang, “An integrated life-cycle greenhouse gas protocol accounting on oil palm trunk and empty fruit bunch biofuel production,” *Sci. Total Environ.*, vol. 856, p. 159007, Jan. 2023, doi: 10.1016/j.scitotenv.2022.159007.
- [17] H. Fujita, K. Nakano, T. Chaisomphob, and E. Hambali, “Greenhouse Gas Emission of Electricity Generation and Field Abandonment Scenarios Using Empty Fruit Bunches at a Palm Oil Mill, Surat Thani Province, Thailand,” *J. Jpn. Inst. Energy*, vol. 98, no. 5, pp. 119–123, 2019, doi: 10.3775/jie.98.119.
- [18] I. Sato, T. Aikawa, C. S. Goh, and C. Kayo, “Life-cycle greenhouse gas emissions in power generation using palm kernel shell,” *GCB Bioenergy*, vol. 14, no. 7, pp. 875–892, 2022, doi: 10.1111/gcbb.12950.
- [19] D. Gamaralalage *et al.*, “Life Cycle Assessment of International Biomass Utilization: A Case Study of Malaysian Palm Kernel Shells for Biomass Power Generation in Japan,” *Waste Biomass Valorization*, vol. 13, no. 5, pp. 2717–2733, May 2022, doi: 10.1007/s12649-021-01643-3.
- [20] International Organization for Standardization, “ISO 14067:2018 Greenhouse gases — Carbon footprint of products — Requirements and guidelines for quantification,” *ISO*, Aug. 2018. <https://www.iso.org/standard/71206.html> (accessed Nov. 24, 2022).
- [21] S. Yusoff, “Renewable energy from palm oil – innovation on effective utilization of waste,” *J. Clean. Prod.*, vol. 14, no. 1, pp. 87–93, Jan. 2006, doi: 10.1016/j.jclepro.2004.07.005.
- [22] M. F. Awalludin, O. Sulaiman, R. Hashim, and W. N. A. W. Nadhari, “An overview of the oil palm industry in Malaysia and its waste utilization through thermochemical conversion, specifically via liquefaction,” *Renew. Sustain. Energy Rev.*, vol. 50, pp. 1469–1484, Oct. 2015, doi: 10.1016/j.rser.2015.05.085.
- [23] N. Rahman, T. B. Bruun, K. E. Giller, J. Magid, G. W. J. van de Ven, and A. de Neergaard, “Soil greenhouse gas emissions from inorganic fertilizers and recycled oil palm waste products from Indonesian oil palm plantations,” *GCB Bioenergy*, vol. 11, no. 9, pp. 1056–1074, 2019, doi: 10.1111/gcbb.12618.
- [24] TNO innovation for life, “Phyllis2 - ECN Phyllis classification-Empty Fruit Bunches palm oil (#2932).” <https://phyllis.nl/Browse/Standard/ECN-Phyllis#2932> (accessed Mar. 04, 2023).
- [25] M. Rizki, S. Jeanly, N. Nanda, G. Phillipe Gunawan, and W. Retono, “Indonesia Zero Emissions Application (EMISI): Methodology for Calculating Individual Emissions from Goods Deliveries, and Marine and Aviation Transport,” Dec. 2021, Accessed: Mar. 03, 2023. [Online]. Available: <https://wri-indonesia.org/en/publications/indonesia-zero-emissions-application-emisi-methodology-calculating-individual-0>
- [26] “AR5 Climate Change 2013: The Physical Science Basis — IPCC.” <https://www.ipcc.ch/report/ar5/wg1/> (accessed Mar. 04, 2023).



- [27] F. Cherubini, G. P. Peters, T. Berntsen, A. H. Strømman, and E. Hertwich, “CO<sub>2</sub> emissions from biomass combustion for bioenergy: atmospheric decay and contribution to global warming,” *GCB Bioenergy*, vol. 3, no. 5, pp. 413–426, Oct. 2011, doi: 10.1111/J.1757-1707.2011.01102.X.
- [28] K. Saswattecha, M. Cuevas Romero, L. Hein, W. Jawjit, and C. Kroeze, “Non-CO<sub>2</sub> greenhouse gas emissions from palm oil production in Thailand,” *J. Integr. Environ. Sci.*, vol. 12, no. sup1, pp. 67–85, Dec. 2015, doi: 10.1080/1943815X.2015.1110184.
- [29] IPCC, “2019 Refinement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories Volume 5 Waste,” IPCC, 2019. [Online]. Available: <https://www.ipcc-nggip.iges.or.jp/public/2019rf/vol5.html>
- [30] Joint Research Centre (European Commission) *et al.*, *Definition of input data to assess GHG default emissions from biofuels in EU legislation: version 1c July 2017*. LU: Publications Office of the European Union, 2017. Accessed: Mar. 05, 2023. [Online]. Available: <https://data.europa.eu/doi/10.2790/658143>
- [31] H.-H. Tao, J. L. Snaddon, E. M. Slade, L. Henneron, J.-P. Caliman, and K. J. Willis, “Application of oil palm empty fruit bunch effects on soil biota and functions: A case study in Sumatra, Indonesia,” *Agric. Ecosyst. Environ.*, vol. 256, pp. 105–113, Mar. 2018, doi: 10.1016/j.agee.2017.12.012.
- [32] H.-H. Tao *et al.*, “Long-term crop residue application maintains oil palm yield and temporal stability of production,” *Agron. Sustain. Dev.*, vol. 37, no. 4, 2017, doi: 10.1007/s13593-017-0439-5.
- [33] L. M. Noirot *et al.*, “Impacts of empty fruit bunch applications on soil organic carbon in an industrial oil palm plantation,” *J. Environ. Manage.*, vol. 317, p. 115373, Sep. 2022, doi: 10.1016/j.jenvman.2022.115373.
- [34] K. Rudolf, N. Hennings, M. A. Dippold, E. Edison, and M. Wollni, “Improving economic and environmental outcomes in oil palm smallholdings: The relationship between mulching, soil properties and yields,” *Agric. Syst.*, vol. 193, p. 103242, Oct. 2021, doi: 10.1016/j.agry.2021.103242.
- [35] “Japan - Countries & Regions,” *IEA*, 2021. <https://www.iea.org/countries/japan> (accessed Mar. 05, 2023).
- [36] P. Moura Costa and C. Wilson, “An equivalence factor between CO<sub>2</sub> avoided emissions and sequestration – description and applications in forestry,” *Mitig. Adapt. Strategy. Glob. Change* 2000 51, vol. 5, no. 1, pp. 51–60, 2000, doi: 10.1023/A:1009697625521.
- [37] F. Joos *et al.*, “Carbon dioxide and climate impulse response functions for the computation of greenhouse gas metrics: A multi-model analysis,” *Atmospheric Chem. Phys.*, vol. 13, no. 5, pp. 2793–2825, 2013, doi: 10.5194/ACP-13-2793-2013.
- [38] W. Chen, Y. Geng, J. Hong, D. Yang, and X. Ma, “Life cycle assessment of potash fertilizer production in China,” *Resour. Conserv. Recycl.*, vol. 138, pp. 238–245, Nov. 2018, doi: 10.1016/j.resconrec.2018.07.028.
- [39] “Google Map,” 2022. <https://www.google.com/maps> (accessed Dec. 12, 2022).
- [40] “SEA-DISTANCES.ORG - Distances,” *SEA-DISTANCE.org*. <https://sea-distances.org/> (accessed Mar. 06, 2023).
- [41] E. Imamura, M. Iuchi, and B. Shigeru, *Comprehensive Assessment of Life Cycle CO<sub>2</sub> Emissions from Power Generation Technologies in Japan*. Central Research Institute of Electric Power Industry, 2016.



- [42] A. Sodri and F. E. Septriana, “Biogas Power Generation from Palm Oil Mill Effluent (POME): Techno-Economic and Environmental Impact Evaluation,” *Energies*, vol. 15, no. 19, Art. no. 19, Jan. 2022, doi: 10.3390/en15197265.