

## DAFTAR PUSTAKA

- Aiman W, Rayi A. Low Grade Gliomas. [Updated 2021 Feb 11]. In: *StatPearls* [Internet]. Treasure Island (FL): StatPearls Publishing; 2021 Jan
- Agustini, Sheila ; Aninditha, Tiara ; Soeridewi, Lyna ; Ranakusuma TA. Penilaian kualitas hidup pada pasien tumor otak. *Neurona*. 2011;28(2).
- Alexiou G, Vartholomatos E, Voulgaris S, Zagorianakou P. Prognostic significance of neutrophil-to-lymphocyte ratio in glioblastoma. *Neuroimmunol Neuroinflammation*. 2014;1(3):131.
- Ashwath KG, Aggarwal A, Praneeth K, Singla N, Gupta K. Neutrophil-To-lymphocyte ratio: Can it be used as an adjunct tool to predict histopathological grade of brain tumor? *J Neurosci Rural Pract*. 2019;10(4):648–52.
- Baehaqi R, Kholis F. Hubungan Antara Jumlah Leukosit Dengan Skor Karnofsky Pada Pasien Kanker Paru. *J Kedokt Diponegoro*. 2012;1(1):139105.
- Bambury RM, Teo MY, Power DG, Yusuf A, Murray S, Battley JE, et al. The association of pre-treatment neutrophil to lymphocyte ratio with overall survival in patients with glioblastoma multiforme. *J Neurooncol*. 2013;114(1):149–54.
- Banerjee P, Leu K, Harris RJ, Cloughesy TF, Lai A, Nghiemphu PL, et al. Association between lesion location and language function in adult glioma using voxel-based lesion-symptom mapping. *NeuroImage Clin* [Internet]. 2015;9:617–24. Available from: <http://dx.doi.org/10.1016/j.nicl.2015.10.010>
- Bao Y, Yang M, Jin C, Hou S, Shi B, Shi J, et al. Preoperative Hematologic Inflammatory Markers as Prognostic Factors in Patients with Glioma. *World Neurosurg* [Internet]. 2018;119:e710–6. Available from: <https://doi.org/10.1016/j.wneu.2018.07.252>
- Bose R, Narang KS, Bhangale D, Kedia R, Sharma V, Jha AN. Survival trends in glioma: Experience at a tertiary care centre. *Neurol India*. 2017;65:1295-1301. Available from: <https://www.neurologyindia.com/text.asp?2017/65/6/1295/217968>
- Brenner A, Friger M, Geffen DB, Kaisman-Elbaz T, Lavrenkov K. The prognostic value of the pretreatment neutrophil/lymphocyte ratio in patients with glioblastoma multiforme brain tumors: A retrospective cohort study of patients treated with combined modality surgery, radiation therapy, and temozolomide chemotherapy. *Oncol*. 2019;97(5):255–63.
- Cagney DN, Alexander BM. The cost and value of glioblastoma therapy. *Expert Rev Anticancer Ther* [Internet]. 2017;17(8):657–9. Available from: <https://doi.org/10.1080/14737140.2017.1351355>
- Chang SM, Nelson S, Vandenberg S, Cha S, Prados M, Butowski N, et al. Integration of preoperative anatomic and metabolic physiologic imaging of newly diagnosed glioma. *J Neurooncol*. 2009;92(3 SPEC. ISS.):401–15.
- Dolan RD, Lim J, McSorley ST, Horgan PG, McMillan DC. The role of the systemic inflammatory response in predicting outcomes in patients with operable cancer: Systematic review and meta-analysis. *Sci Rep*. 2017;7(1):1–31.

- Eisenstat DD, Pollack IF, Demers A, Sapp M V., Lambert P, Weisfeld-Adams JD, et al. Impact of tumor location and pathological discordance on survival of children with midline high-grade gliomas treated on Children's Cancer Group high-grade glioma study CCG-945. *J Neurooncol.* 2015;121(3):573–81.
- Febrian, Dananjoyo, Malueka, Faktor faktor yang memengaruhi status performa pasien pada pasien glioma derajat tinggi. *Jaknews* 2021.
- Fallo INY, Sidharta RA, Gunawan LS. Korelasi antara Neutrophyl Lymphocyte Ratio dengan Stadium Kanker pada Pasien Kanker Payudara. *Biomedika.* 2018;11(2):63–9.
- Forget, P., Khalifa, C., Defour, J.P., Latinne, D., Van Pel, M.C. & De Kock, M. 2017. What is the normal value of the neutrophil-to-lymphocyte ratio? *BMC Res. Notes.*, 10:12. DOI 10.1186/s13104-016-2335-5.
- Forst, D.A, Brian V. Nahed, Jay S. Loeffler TTB. Low-grade gliomas. *The Oncologist.* 2014;19:403–13.
- Figuroa JM, Semonche A, Magoon S, Shah A, Luther E, Eichberg D, et al. The role of neutrophil-to-lymphocyte ratio in predicting overall survival in patients undergoing laser interstitial thermal therapy for glioblastoma. *J Clin Neurosci* [Internet]. 2020;72(XXXX):108–13. Available from: <https://doi.org/10.1016/j.jocn.2019.12.057>
- Forst DA, Quain KM, Poisson K, Eusebio J, Jackson VA, Greer JA, et al. Molecular Pathology and Classification -. 2017;6(June 2015):2017.
- Galvão RP, Zong H. Inflammation and Gliomagenesis: Bi-Directional Communication at Early and Late Stages of Tumor Progression. *Curr Pathobiol Rep.* 2013;1(1):19–28.
- Gladson CL, Prayson RA, Liu WM. The pathobiology of glioma tumors. *Annu Rev Pathol Mech Dis.* 2010;5:33–50.
- Gomes LT, Morato-Conceição YT, Gambati AVM, Maciel-Pereira CM, Fontes CJF. Diagnostic value of neutrophil-to-lymphocyte ratio in patients with leprosy reactions. *Heliyon.* 2020;6(2).
- Gregory AD, Houghton AM. Tumor-associated neutrophils: new targets for cancer therapy. *Cancer Res.* 2011 Apr 01;71(7):2411–2416.
- Gunawan PY. Hubungan Karakteristik Biologi Molekuler , Derajat Glioma , dan Lokasi Tumor dengan Perbaikan Status Fungsional pada Pasien Glioma. 2020;0–121. <http://repository.unhas.ac.id/id/eprint/3308/>
- Gunawan PY, Islam A, July J, Patellongi I, Nasrum M, Aninditha T. Karnofsky Performance Scale and Neurological Assessment of Neuro-Oncology Scale as Early Predictor in Glioma. *Asian Pacific J Cancer Prev.* 2020;21(11):3387–92.
- Guyton, A.C. & Hall, J.E. 2006. Textbook of Medical Physiology, 429-438. Ed 11. Pennsylvania: Elsevier Inc.
- Han S, Liu Y, Li Q, Li Z, Hou H, Wu A. Pre-treatment neutrophil-to-lymphocyte ratio is associated with neutrophil and T-cell infiltration and predicts clinical outcome in patients with glioblastoma. *BMC Cancer* [Internet]. 2015;15(1):1–10. Available from: <http://dx.doi.org/10.1186/s12885-015-1629-7>
- Hardiany NS, Mulyawan W, Wanandi SI. Correlation between oxidative stress and tumor grade in glioma cells from patients in Jakarta. *Med J Indones.* 2012;21(3):122–7.
- Heidel E. Youden Index: Determine the overall benefit of using a diagnostic test. 2022. Available from: <https://www.scalestatistics.com/youden-index.html>

- John F. de Groot. High-grade Gliomas. *Neuro-oncology* p. 332-344 April 2015, Vol.21, No.2 doi: 10.1212/01.CON.0000464173.58262.d9
- Kelly WJ, Giles AJ, Gilbert M. T lymphocyte-targeted immune checkpoint modulation in glioma. *J Immunother Cancer*. 2020;8(1):1–18.
- Kemenkes. Panduan Nasional Pelayanan Kesehatan Tumor Otak. 2017;92.
- Kemerdere R, Akgun MY, Toklu S, Alizada O, Tanriverdi T. Preoperative systemic inflammatory markers in low- and high-grade gliomas: A retrospective analysis of 171 patients. *Heliyon* [Internet]. 2019;5(5):e01681.
- Laila, M. N., Andriani, R., Sofyan, H. R., & Aninditha, T. (2019). Faktor-faktor yang Memengaruhi Status Performa Pasien Tumor Intrakranial di RSUPN dr. Cipto Mangunkusumo. *Majalah Kedokteran Neurosains Perhimpunan Dokter Spesialis Saraf Indonesia*, 37(1). <https://doi.org/10.52386/neurona.v37i1.91>
- Larjavaara S, Mäntylä R, Salminen T, Haapasalo H, Raitanen J, Jääskeläinen J, et al. Incidence of gliomas by anatomic location. *Neuro Oncol*. 2007;9(3):319–25.
- Lee J, Park SH, Kim YZ. Prognostic Evaluation of Neurological Assessment of the Neuro-Oncology Scale in Glioblastoma Patients. *Brain Tumor Res Treat*. 2018;6(1):22.
- Leick, M., Azcutia, V., Newton, G. & Luscinskas, F.W. 2014. Leukocyte Recruitment in Inflammation: Basic Concepts and New Mechanistic Insights Based on New Models and Microscopic Imaging Technologies. *Cell Tis-sue Res.*, 355(3):647-56. DOI:10.1007/s00441-014-1809-9.
- Liu J, Liu Y, Xiang P, Pu L, Xiong H, Li C, et al. Neutrophil-to-lymphocyte ratio predicts critical illness patients with 2019 coronavirus disease in the early stage. *J Transl Med*. 2020;18(1).
- Malueka RG, Dwianingsih EK, Bayuangga HF, Argo IW, Donurizki AD, Shaleh S, et al. Association of IDH1 mutation with survival of glioma patients in Indonesia. *J Neurol Sci*. 2019;405:101.
- Malueka RG, Yudiyanta Y, Asmedi A. Peran Penanda Molekuler Pada Terapi Glioma. *Majalah Kedokteran Neurosains PERDOSSI*. 2017;35(1):36–44.
- McCrea HJ, Bander ED, Venn RA, Reiner AS, Iorgulescu JB, Puchi LA, et al. Sex, age, anatomic location, and extent of resection influence outcomes in children with high-grade glioma. *Neurosurgery*. 2015;77(3):443–52.
- Marinari E, Allard M, Gustave R, Widmer V, Philippin G, Merkler D, et al. Inflammation and lymphocyte infiltration are associated with shorter survival in patients with high-grade glioma. *Oncoimmunology* [Internet]. 2020;9(1).
- Maru GB, Gandhi K, Ramchandani A, Kumar G. Inflammatory signaling pathways associated with Skin Cancer. *Inflammation and Cancer*. 2014. 443–448 p.
- Massara M, Persico P, Bonavita O, Poeta VM, Locati M, Simonelli M, et al. Neutrophils in gliomas. *Front Immunol*. 2017;8(OCT):1–7.
- Ohgaki H, Kleihues P. The definition of primary and secondary glioblastoma. *Clin Cancer Res*. 2013;19(4):764–72.
- Ostrom QT, Gittleman H, Fulop J, Liu M, Blanda R, Kromer C, et al. CBTRUS statistical Report: primary brain and central nervous system tumors diagnosed in the United States in 2008-2012. *Neuro Oncol*. 2015;17:iv1–62.

- Ostrom QT, Patil N, Cioffi G, Waite K, Kruchko C, Barnholtz-Sloan JS. CBTRUS statistical report: Primary brain and other central nervous system tumors diagnosed in the United States in 2013-2017. *Neuro Oncol*. 2020;22(Supplement\_1):IV1–96.
- Panciroli C, Estival A, Lucente G, Velarde JM, Garcia R, Vila L, et al. ECOG or Karnofsky Performance Status to Assess Functionality in Glioblastoma Patients Among Different Observers. *J Mol Biomark Diagn*. 2017;01(S2).
- Oslobanu A, Florian SI. Anatomic locations in high grade glioma. *Rom Neurosurg*. 2015;29(3):271–7.
- Park SA, Chung SH, Lee Y. Factors influencing the quality of life of patients with advanced cancer. *Appl Nurs Res* [Internet]. 2017;33:108–12. Available from: <http://dx.doi.org/10.1016/j.apnr.2016.10.002>
- Péus D, Newcomb N, Hofer S. Appraisal of the Karnofsky Performance Status and proposal of a simple algorithmic system for its evaluation. *BMC Med Inform Decis Mak* [Internet]. 2013;13(1):1. Available from: BMC Medical Informatics and Decision Making
- Pratiknya, A.W. 2011. Dasar-Dasar Metodologi Penelitian Kedokteran dan Kesehatan. Jakarta: PT. Raja Grafindo Persada.
- Preusser M, Haberler C, Hainfellner JA. Malignant glioma: Neuropathology and neurobiology. *Wiener Medizinische Wochenschrift*. 2006;156(11–12): 332–7.
- Rasmussen BK, Hansen S, Laursen RJ, Kosteljanetz M, Schultz H, Nørgård BM, et al. Epidemiology of glioma: clinical characteristics, symptoms, and predictors of glioma patients grade I–IV in the the Danish Neuro-Oncology Registry. *J Neurooncol* [Internet]. 2017;135(3):571–9. Available from: <http://dx.doi.org/10.1007/s11060-017-2607-5>
- Ritarwan K, Nasution IK, Erwin I, Nerdy N. Correlation of leukocyte subtypes, neutrophil-to-lymphocyte ratio, and functional outcome in brain metastasis. *Open Access Maced J Med Sci*. 2018;6(12):2333–6.
- Sadighi ZS, Curtis E, Zabrowski J, Billups C, Gajjar A, Khan R, et al. Neurologic impairments from pediatric low-grade glioma by tumor location and timing of diagnosis. *Pediatr Blood Cancer*. 2018;65(8):1–8.
- Sagberg LM, Iversen DH, Fyllingen EH, Jakola AS, Reinertsen I, Solheim O. Brain atlas for assessing the impact of tumor location on perioperative quality of life in patients with high-grade glioma: A prospective population-based cohort study. *NeuroImage Clin* [Internet]. 2019;21(January):101658.
- Sastroasmoro, S., Ismael, S. 2018. Dasar-Dasar Metodologi Penelitian Klinis. Edisi ke-5 cetakan ke-3. Jakarta: *Sagung Seto*.
- Schag CC, Heinrich RL, Ganz PA. Karnofsky performance status revisited: Reliability, validity, and guidelines. *J Clin Oncol*. 1984;2(3):187–93.
- Shaul, Merav & Fridlender, Zvi. (2017). Neutrophils as active regulators of the immune system in the tumor microenvironment. *Journal of Leukocyte Biology*. 102. jlb.5MR1216-508R. 10.1189/jlb.5MR1216-508R.
- Spena G, D'Agata F, Panciani PP, Buglione di Monale M, Fontanella MM. Supratentorial gliomas in eloquent areas: which parameters can predict functional outcome and extent of resection?. *PLoS One*. 2013;8(12):e80916. Published 2013 Dec 5.
- Suzuki a, Tsutomi Y, Shimizu M, Matsuzawa a. Karnofsky Performance Status Scale Definitions Rating. *Oxford Textb Palliat Med* [Internet]. 1994;6:2. Available from: <http://www.hospicepatients.org/karnofsky.html>

- Tang Q, Lian Y, Yu J, Wang Y, Shi Z, Chen L. Anatomic mapping of molecular subtypes in diffuse glioma. *BMC Neurol.* 2017;17(1):1–8.
- Theresia E, Malueka RG, Pranacipta S, Kameswari B, Dananjoyo K, Asmedi A, et al. Association between Ki-67 labeling index and histopathological grading of Glioma in Indonesian population. *Asian Pacific J Cancer Prev.* 2020;21(4):1063–8.
- Topkan E, Selek U, Ozdemir Y, Yildirim BA, Guler OC, Ciner F, et al. Prognostic value of the Glasgow Prognostic Score for glioblastoma multiforme patients treated with radiotherapy and temozolomide. *J Neurooncol* [Internet]. 2018;139(2):411–9. Available from: <http://dx.doi.org/10.1007/s11060-018-2879-4>
- Tortora, G.J. & Derrickson, B. 2009. *Principles of Anatomy and Physiology*, 699-701. Ed 12. Hoboken: *John Wiley & Sons Inc.*
- Vergani F, Martino J, Gozé C, Rigau V, Duffau H. World health organization grade II gliomas and subventricular zone: Anatomic, genetic, and clinical considerations. *Neurosurgery.* 2011;68(5):1293–8.
- Wang CH, Lin CY, Chen JS, Ho CL, Rau KM, Tsai JT, et al. Karnofsky performance status as a predictive factor for cancer-related fatigue treatment with astragalus polysaccharides (PG2) injection-A double blind, multi-center, randomized phase IV study. *Cancers (Basel).* 2019;11(2).
- Wang J, Xiao W, Chen W, Hu Y. Prognostic significance of preoperative neutrophil-to-lymphocyte ratio and platelet-to-lymphocyte ratio in patients with glioma. *EXCLI J.* 2018;17:505–12.
- Wang PF, Meng Z, Song HW, Yao K, Duan ZJ, Yu CJ, et al. Preoperative changes in hematological markers and predictors of glioma grade and survival. *Front Pharmacol.* 2018;9(AUG):1–7.
- Wang ZL, Zhang CB, Liu YQ, Wang Z, Jiang T. Peripheral blood test provides a practical method for glioma evaluation and prognosis prediction. *CNS Neurosci Ther.* 2019;25(8):876–83.
- Weiji Weng, Xu Chen, Shaohui Gong, Liemei Guo & Xiaohua Zhang (2018):Preoperative neutrophil–lymphocyte ratio correlated with glioma grading and glioblastoma survival, *Neurological Research*. <https://doi.org/10.1080/01616412.2018.1497271>
- Weller M, Wick W, Aldape K, Brada M, Berger M, Pfister SM, et al. Glioma. *Nat Rev Dis Prim.* 2015;1(July).
- Weng W, Chen X, Gong S, Guo L, Zhang X. Preoperative neutrophil–lymphocyte ratio correlated with glioma grading and glioblastoma survival. *Neurol Res* [Internet]. 2018;40(11):917–22.
- Wright, H.L., Moots, R.J., Bucknall, R.C. & Edwards, S.W. 2010. Neutrophil function in inflammation and inflammatory diseases. *Rheumatology*, 49: 1618-31. DOI:10.1093/rheumatology/keq045.
- Xu W, Wang D, Zheng X, Ou Q, Huang L. Sex-dependent association of preoperative hematologic markers with glioma grade and progression. *J Neurooncol* [Internet]. 2018;137(2):279–87. Available from: <http://dx.doi.org/10.1007/s11060-017-2714-3>
- Zadora P, Dabrowski W, Czarko K, Smolen A, Kotlinska-Hasiec E, Wiorowski K, et al. Preoperative neutrophil-lymphocyte count ratio helps predict the grade of glial tumor—a pilot study. *Neurol Neurochir Pol.* 2015;49(1):41–4.



- Zha C, Meng X, Li L, Mi S, Qian D, Li Z, et al. Neutrophil extracellular traps mediate the crosstalk between glioma progression and the tumor microenvironment via the HMGB1/RAGE/IL-8 axis. *Cancer Biol Med.* 2020;17(1):154–68.
- Zhang H, Fan F, Yu Y, Wang Z, Liu F, Dai Z, et al. Clinical characterization, genetic profiling, and immune infiltration of TOX in diffuse gliomas. *J Transl Med* [Internet]. 2020;18(1):1–16. Available from: <https://doi.org/10.1186/s12967-020-02460-3>
- Zhang J, Yao L, Peng S, Fang Y, Tang R, Liu J. Correlation between glioma location and preoperative seizures: a systematic review and meta-analysis. *Neurosurg Rev.* 2019;42(3):603–18.
- Zhang J, Cai L, Song Y, Shan B, He M, Ren Q, et al. Prognostic role of neutrophil lymphocyte ratio in patients with spontaneous intracerebral hemorrhage. *Oncotarget.* 2017;8(44):77752–60.
- Zheng SH, Huang JL, Chen M, Wang BL, Ou QS, Huang SY. Diagnostic value of preoperative inflammatory markers in patients with glioma: A multicenter cohort study. *J Neurosurg.* 2018;129(3):583–92.
- Zou Y, Zhang W, Huang C, Zhu Y. Clinical significance of neutrophil to lymphocyte ratio and platelet to lymphocyte ratio in acute cerebral hemorrhage with gastrointestinal hemorrhage, and logistic regression analysis of risk factors. *Exp Ther Med.* 2019;1533–8.