

DAFTAR PUSTAKA

- Aydın, M.M. and Akçalı, K.C. (2018). Liver fibrosis. *The Turkish Journal of Gastroenterology*, 29(1), pp.14–21.
- Boscarelli, A. (2019). Kasai Hepatoportoenterostomy for Biliary Atresia in Children: Technical Notes and Details of Perioperative Therapy. *International Journal of Pediatrics and Child Health*, 7(1), pp.24–27.
- Burns, J. and Davenport, M. (2020). Adjuvant treatments for biliary atresia. *Translational Pediatrics*, 9(3), pp.253–265.
- Chardot, C. (2006). Biliary atresia. *Orphanet Journal of Rare Diseases*, 1(1).
- Chin, J., Powell, L.W., Ramm, L.E., Hartel, G.F., Olynyk, J.K. and Ramm, G.A. (2020). Utility of Serum Biomarker Indices for Staging of Hepatic Fibrosis Before and After Venesection in Patients With Hemochromatosis Caused by Variants in HFE. *Clinical Gastroenterology and Hepatology*, 19(7).
- Daniel Kalman Podolsky (2016). *Yamada's textbook of gastroenterology*. Chichester, West Sussex: John Wiley & Sons Inc.
- Davenport, M., Muntean, A. and Hadzic, N. (2021). Biliary Atresia: Clinical Phenotypes and Aetiological Heterogeneity. *Journal of Clinical Medicine*, 10(23), p.5675.
- Degtyareva, A., Razumovskiy, A., Kulikova, N., Ratnikov, S., Filippova, E., Gordeeva, E., Albegova, M., Rebrikov, D. and Puchkova, A. (2020). Long-

- Term Effects of Kasai Portoenterostomy for Biliary Atresia Treatment in Russia. *Diagnostics*, 10(9), p.686.
- Feldman, A.G. and Mack, C.L. (2012). Biliary atresia: cellular dynamics and immune dysregulation. *Seminars in Pediatric Surgery*, 21(3), pp.192–200.
- Friedman, M. (2020). *Sleisenger and Fordtran's Gastrointestinal and Liver Disease- 2 Volume Set : pathophysiology, diagnosis, management*. S.L.: Elsevier - Health Science.
- Haafiz, A.B. (2010). Liver fibrosis in biliary atresia. *Expert Review of Gastroenterology & Hepatology*, 4(3), pp.335–343.
doi:<https://doi.org/10.1586/egh.10.29>.
- Hukkinen, M., Lohi, J., Heikkilä, P., Kivisaari, R., Jahnukainen, T., Jalanko, H., and Pakarinen, M. P. (2019). Noninvasive Evaluation of Liver Fibrosis and Portal Hypertension After Successful Portoenterostomy for Biliary Atresia. *Hepatology Communications*, 3(3), 382–391.
- Jancelewicz, T., Barmherzig, R., Chung, C.T.-S. ., Ling, S.C., Kamath, B.M., Ng, V.L., Amaral, J., O'Connor, C., Fecteau, A. and Langer, J.C. (2015). A screening algorithm for the efficient exclusion of biliary atresia in infants with cholestatic jaundice. *Journal of Pediatric Surgery*, 50(3), pp.363–370.
- Jiang, J., Wan, R., He, S., Wu, Y., Shen, Z., Chen, G., Sun, S., Yan, W. and Zheng, S. (2021). Epidemiological characteristics and risk factors of biliary atresia: a case–control study. *BMJ Open*, [online] 11(12), p.e049354.

- Kim, S.Y., Seok, J.Y., Han, S.J. and Koh, H. (2010). Assessment of Liver Fibrosis and Cirrhosis by Aspartate Aminotransferase-to-Platelet Ratio Index in Children With Biliary Atresia. *Journal of Pediatric Gastroenterology & Nutrition*, 51(2), pp.198–202.
- Lala, V. and Minter, D.A. (2019). *Liver Function Tests*. [online] Nih.gov. Available at: <https://www.ncbi.nlm.nih.gov/books/NBK482489/>.
- Lin, Z.-H., Xin, Y.-N., Dong, Q.-J., Wang, Q., Jiang, X.-J., Zhan, S.-H., Sun, Y. and Xuan, S.-Y. (2011). Performance of the aspartate aminotransferase-to-platelet ratio index for the staging of hepatitis C-related fibrosis: An updated meta-analysis. *Hepatology*, 53(3), pp.726–736.
- Loeza-del-Castillo, A., Paz-Pineda, F., Oviedo-Cárdenas, E., Sánchez-Ávila, F. and Vargas-Voráčková, F. (2008). AST to platelet ratio index (APRI) for the noninvasive evaluation of liver fibrosis. *Annals of Hepatology*, 7(4), pp.350–357.
- Lurie, Y. (2015). Non-invasive diagnosis of liver fibrosis and cirrhosis. *World Journal of Gastroenterology*, [online] 21(41), p.11567.
- Mcperson, R.A. and Pincus, M.R. (2017). *Henry's Clinical Diagnosis and Management by Laboratory Methods*. 23rd ed. St. Louis, Missouri: Elsevier.
- Mohamed Sira, M. (2017). Utility of Noninvasive Serum Biomarkers of Liver Fibrosis in Infants with Biliary Atresia. *Advanced Research in Gastroenterology & Hepatology*, 3(2).

Neto, B., Borges-Dias, M., Trindade, E., Estevão-Costa, J. and Campos, J.M. (2017).

Biliary Atresia - Clinical Series. *GE - Portuguese Journal of Gastroenterology*, [online] 25(2), pp.68–73.
doi:<https://doi.org/10.1159/000480708>.

Nomden, M., Beljaars, L., Verkade, H.J., Hulscher, J.B.F. and Olinga, P. (2020).

Current Concepts of Biliary Atresia and Matrix Metalloproteinase-7: A Review of Literature. *Frontiers in Medicine*, 7.

Price, K.B. and Temples, H.S. (2022). Neonatal Jaundice: Early Identification of

Biliary Atresia. *The Journal for Nurse Practitioners*.

Qisthi, S.A., Saragih, D.S.P., Sutowo, D.W., Sirait, D.N., Imelda, P., Kencana,

S.M.S., Makhmudi, A. and Gunadi (2020). Prognostic Factors for Survival of Patients with Biliary Atresia Following Kasai Surgery. *The Kobe journal of medical sciences*, [online] 66(2), pp.E56–E60.

Redkar, R., Karkera, P.J., Raj, V., Bangar, A., Hathiramani, V. and Krishnan, J.

(2017). Outcome of biliary atresia after Kasai's portoenterostomy: A 15-year experience. *Indian Pediatrics*, 54(4), pp.291–294. doi:10.1007/s13312-017-1091-5.

Sastiono, S., Maheranny, M. and Oswari, H. (2016). Factors Affecting the Success

Rate of Kasai Portoenterostomy. *The New Ropanasuri Journal of Surgery*, 1(1), pp.27–30.

Shao, L.-N., Zhang, S.-T., Wang, N., Yu, W.-J., Chen, M., Xiao, N., Duan, Y., Pan,

L.-Z., Song, W.-Q., Xia, Y.-X., Zhang, L., Qi, N., Liu, M. and Zhou, S.-H.

- (2020). Platelet indices significantly correlate with liver fibrosis in HCV-infected patients. *PLOS ONE*, 15(1), p.e0227544.
doi:<https://doi.org/10.1371/journal.pone.0227544>.
- Shen, W.-J., Chen, G., Wang, M. and Zheng, S. (2018). Liver fibrosis in biliary atresia. *World Journal of Pediatrics*, 15(2), pp.117–123.
- Siddiqui, A.I. and Ahmad, T. (2020). *Biliary Atresia*. [online] PubMed.
- Suk, K.T. (2015). Staging of liver fibrosis or cirrhosis: The role of hepatic venous pressure gradient measurement. *World Journal of Hepatology*, 7(3), p.607.
- Vij, M. and Rela, M. (2020). Biliary atresia: pathology, etiology and pathogenesis. *Future Science OA*, p.FSO466.
- Wang, K.S. (2015). Newborn Screening for Biliary Atresia. *PEDIATRICS*, 136(6), pp.e1663–e1669.
- Wildhaber, B.E. (2012). Biliary atresia: 50 years after the first kasai. *ISRN surgery*, [online] 2012, p.132089.
- Williams, N.S., C J K Bulstrode, P Ronan O'connell, Bailey, H. and Love (2018). *Bailey & Love's short practice of surgery*. Boca Raton: Crc Press.
- Yang, L.-Y., Fu, J., Peng, X.-F., Pang, S.-Y., Gao, K.-K., Chen, Z.-R., He, L.-J., Wen, Z., Wang, H., Li, L., Wang, F.-H., Yu, J.-K., Xu, Y., Gong, S.-T., Xia, H.-M. and Liu, H.-Y. (2015). Validation of aspartate aminotransferase to platelet ratio for diagnosis of liver fibrosis and prediction of postoperative prognosis in infants with biliary atresia. *World Journal of Gastroenterology*, 21(19), pp.5893–5900.

Yerina, S.E. and Ekong, U.D. (2021). Biliary Atresia/Neonatal Cholestasis. *Pediatric*

Clinics of North America, 68(6), pp.1333–1341.

Zani, A., Quaglia, A., Hadzić, N., Zuckerman, M. and Davenport, M. (2015).

Cytomegalovirus-associated biliary atresia: An aetiological and prognostic

subgroup. *Journal of Pediatric Surgery*, 50(10), pp.1739–1745.