

DAFTAR PUSTAKA

- American College of Obstetricians And Gynecologists (2016) ‘ACOG PB 171 management of preterm labor’, *Journal of Pediatrics*, 128(4), pp. 155–164.
- Angum, F. *et al.* (2020) ‘The Prevalence of Autoimmune Disorders in Women: A Narrative Review’, *Cureus*, 12(5). doi: 10.7759/cureus.8094.
- Asai, A., Miethke, A. and Bezerra, J. A. (2015) ‘Pathogenesis of biliary atresia: defining biology to understand clinical phenotypes’, *Nature reviews. Gastroenterology & hepatology*, 12(6), p. 342. doi: 10.1038/NRGASTRO.2015.74.
- Bezerra, J. A. *et al.* (2018) ‘Biliary Atresia: Clinical and Research Challenges for the Twenty-First Century’, *Hepatology*, 68(3), pp. 1163–1173. doi: 10.1002/hep.29905.
- Cavallo, L. *et al.* (2022) ‘The Epidemiology of Biliary Atresia: Exploring the Role of Developmental Factors on Birth Prevalence’, *The Journal of Pediatrics*, 246, pp. 89-94.e2. doi: 10.1016/J.JPIDS.2022.03.038.
- Chiu, C. Y. *et al.* (2013) ‘Biliary atresia in preterm infants in Taiwan: A nationwide survey’, *Journal of Pediatrics*, 163(1), pp. 1–5. doi: 10.1016/j.jpids.2012.12.085.
- Chung, P. H. Y., Zheng, S. and Tam, P. K. H. (2020) ‘Biliary atresia: East versus west’, *Seminars in Pediatric Surgery*, 29(4). doi: 10.1016/j.sempedsurg.2020.150950.
- Dumont-Lagacé, M., St-Pierre, C. and Perreault, C. (2015) ‘Sex hormones have pervasive effects on thymic epithelial cells’, *Scientific Reports*, 5, pp. 1–12. doi:

10.1038/srep12895.

Durkin, N., Deheragoda, M. and Davenport, M. (2017) 'Prematurity and biliary atresia: a 30-year observational study', *Pediatric Surgery International*, 33(12), pp. 1355–1361. doi: 10.1007/s00383-017-4193-1.

Fischler, B., Haglund, B. and Hjern, A. (2002) 'A population-based study on the incidence and possible pre- and perinatal etiologic risk factors of biliary atresia', *Journal of Pediatrics*, 141(2), pp. 217–222. doi: 10.1067/mpd.2002.126001.

Gunadi *et al.* (2018) 'Liver transplant score for prediction of biliary atresia patients' survival following Kasai procedure', *BMC Research Notes*, 11(1), pp. 1–5. doi: 10.1186/s13104-018-3498-z.

Hartley, J. L., Davenport, M. and Kelly, D. A. (2009) 'Biliary atresia', *Lancet (London, England)*, 374(9702), pp. 1704–1713. doi: 10.1016/S0140-6736(09)60946-6.

Invernizzi, P. *et al.* (2009) 'Female predominance and X chromosome defects in autoimmune diseases', *Journal of Autoimmunity*, 33(1), pp. 12–16. doi: 10.1016/j.jaut.2009.03.005.

Jiang, J. *et al.* (2021) 'Epidemiological characteristics and risk factors of biliary atresia: A case-control study', *BMJ Open*, 11(12), pp. 1–9. doi: 10.1136/bmjopen-2021-049354.

Karri, P. S. *et al.* (2022) 'Biliary Atresia Screening in India—Strategies and Challenges in Implementation', *Indian Journal of Pediatrics*, 89(2), pp. 133–140. doi: 10.1007/S12098-021-03862-X/TABLES/4.

Lakshminarayanan, B. and Davenport, M. (2016) 'Biliary atresia: A comprehensive

- review', *Journal of Autoimmunity*, 73, pp. 1–9. doi: 10.1016/j.jaut.2016.06.005.
- Lee, K. J. *et al.* (2017) 'Epidemiology of biliary atresia in Korea', *Journal of Korean Medical Science*, 32(4), pp. 656–660. doi: 10.3346/jkms.2017.32.4.656.
- Livesey, E. *et al.* (2009) 'Epidemiology of biliary atresia in England and Wales (1999-2006)', *Archives of Disease in Childhood: Fetal and Neonatal Edition*, 94(6), pp. 451–456. doi: 10.1136/adc.2009.159780.
- Mack, C. L. and Sokol, R. J. (2005) 'Unraveling the Pathogenesis and Etiology of Biliary Atresia', *Pediatric Research*, 57(5), pp. 87R-94R. doi: 10.1203/01.PDR.0000159569.57354.47.
- Medicine, M. (2013) 'ACOG Committee opinion No 579: Definition of Term Pregnancy', *Obstetrics and Gynecology*, 122(5), pp. 1139–1140.
- Narkewicz, M. R. (2001) 'Biliary atresia: An update on our understanding of the disorder', *Current Opinion in Pediatrics*, 13(5), pp. 435–440. doi: 10.1097/00008480-200110000-00009.
- Qisthi, S. A. *et al.* (2020) 'Prognostic factors for survival of patients with biliary atresia following kasai surgery', *Kobe Journal of Medical Sciences*, 66(2), pp. E56–E60. Available at: <https://pubmed.ncbi.nlm.nih.gov/33024065/> (Accessed: 8 November 2022).
- Ramanathan, K. *et al.* (2020) 'Preterm birth1: Epidemiology and causes of preterm birth', *Lancet* 200(January), pp. 75–84.
- Raqib, R. *et al.* (2007) 'Low birth weight is associated with altered immune function in rural Bangladeshi children: A birth cohort study', *American Journal of Clinical Nutrition*, 85(3), pp. 845–852. doi: 10.1093/ajcn/85.3.845.

- Siddiqui, A. and Ahmad, I. (2022) *Biliary Atresia*, StatPearls Publishing. Treasure Island (FL): StatPearls Publishing. Available at: <https://www.ncbi.nlm.nih.gov/books/NBK537262/>.
- Sinha, C. K. and Davenport, M. (2008) 'Biliary atresia', *Journal of Indian Association of Pediatric Surgeons*, 13(2), p. 49. doi: 10.4103/0971-9261.43015.
- Sokol, R. J. and Mack, C. (2001) 'Etiopathogenesis of biliary atresia', *Seminars in Liver Disease*, 21(4), pp. 517–524. doi: 10.1055/s-2001-19032.
- The, N. S. *et al.* (2007) 'Risk factors for isolated biliary atresia, National Birth Defects Prevention Study, 1997–2002', *American Journal of Medical Genetics Part A*, 143(19), pp. 2274–2284.
- Tsimis, M. E. *et al.* (2015) 'Prematurity: present and future', *Minerva ginecologica*, 67(1), pp. 35–46.
- Uemura, M. *et al.* (2013) 'Sox17 haploinsufficiency results in perinatal biliary atresia and hepatitis in C57BL/6 background mice', *Development (Cambridge)*, 140(3), pp. 639–648. doi: 10.1242/dev.086702.
- Vij, M. and Rela, M. (2020) 'Biliary atresia: pathology, etiology and pathogenesis', *Future Sci OA*, 6(5), p. FSO466. Available at: <https://doi.org/10.2144/fsoa-2019-0153>.
- Wada, H. *et al.* (2007) 'Insignificant seasonal and geographical variation in incidence of biliary atresia in Japan: a regional survey of over 20 years', *Journal of Pediatric Surgery*, 42(12), pp. 2090–2092. doi: 10.1016/j.jpedsurg.2007.08.035.
- Wang, K. S. (2015) 'Newborn Screening for Biliary Atresia', *Pediatrics*, 136(6), p.

e1663. doi: 10.1542/PEDS.2015-3570.

Wang, L. *et al.* (2018) 'Early differential diagnosis methods of biliary atresia: a meta-analysis', *Pediatric Surgery International*, 34(4), pp. 363–380. doi: 10.1007/s00383-018-4229-1.

WHO (2016) 'International Statistical Classification of Diseases and Related Health Problems, 10th Revision ICD-10: Tabular List', *World Health Organization*, 1, pp. 2, 33–95. Available at: <http://www.who.int/classifications/icd/icdonlineversions/en/>.

Xu, Y. *et al.* (2012) 'The perinatal infection of cytomegalovirus is an important etiology for biliary atresia in China', *Clinical Pediatrics*, 51(2), pp. 109–113. doi: 10.1177/0009922811406264.

Zani, A. *et al.* (2015) 'Cytomegalovirus-associated biliary atresia: An aetiological and prognostic subgroup', *Journal of Pediatric Surgery*, 50(10), pp. 1739–1745. doi: 10.1016/j.jpedsurg.2015.03.001.