

DAFTAR PUSTAKA

- Abukari, A. (2019). Influence of rice husk biochar on water holding capacity of soil in the savannah ecological zone of Ghana. *Turkish Journal of Agriculture - Food Science and Technology*, 7(6), 888–891.
- Ahmed, A., Kurian, J., & Raghavan, V. (2016). Biochar influences on agricultural soils, crop production, and the environment: A review. *Environmental Reviews*, 24(4), 495–502.
- Alexis, M. A., Rasse, D. P., Rumpel, C., Bardoux, G., Péchot, N., Schmalzer, P., Drake, A., & Mariotti, B. (2007). Fire impact on C and N losses and charcoal production in a scrub oak ecosystem. *Biogeochemistry Volume*, 82, 201–216.
- Ali, A. Y. A., Zhou, G., Elsiddig, A. M., Zhu, G., Meng, T., Jiao, X., Ahmed, I., Ibrahim Salih, E. G., & Hussien Ibrahim, M. E. (2022). Effects of jasmonic acid in foliar spray and an humic acid amendment to saline soils on forage sorghum plants' growth and antioxidant defense system. *PeerJ*, 10, 1–25.
- Ali, B. (2017). Salicylic acid induced antioxidant system enhances the tolerance to aluminium in mung bean (*Vigna radiata* L. Wilczek) plants. *Indian Journal of Plant Physiology*, 22(2), 178–189.
- Amri, E. (2014). The role of selected plant families with dietary Ethnomedicinal species used as anticancer. *Journal of Medicinal Plants Studies*, 2(1), 28–39. www.plantsjournal.com
- Anjum, Shakeel A., Ashraf, U., Tanveer, M., Khan, I., Hussain, S., Shahzad, B., Zohaib, A., Abbas, F., Saleem, M. F., Ali, I., & Wang, L. C. (2017). Drought induced changes in growth, osmolyte accumulation and antioxidant metabolism of three maize hybrids. *Frontiers in Plant Science*, 8(february), 1–12.
- Anjum, Shakeel, A., Xie, X., Wang, L., Saleem, M. F. M, C., & Lei, W. (2011). Morphological, physiological and biochemical responses of plants to drought stress. *African Journal of Agricultural Research*, 6(9), 2026–2032.
- Anyanwu, I. N., Alo, M. N., Onyekwere, A. M., Crosse, J. D., Nworie, O., & Chamba, E. B. (2018). Influence of biochar aged in acidic soil on ecosystem engineers and two tropical agricultural plants. *Ecotoxicology and Environmental Safety*, 153, 116–126.
- Arnon, D. I. (1949). Cooper enzyme in isolated chloroplast. Polyphenoloxidase in *Beta vulgaris*. *Plant Physiology*, 24(1), 1–15.
- Asai, H., Samson, B. K., Stephan, H. M., Songyikhangsuthor, K., Homma, K., Kiyono, Y., Inoue, Y., Shiraiwa, T., & Horie, T. (2009). Biochar amendment techniques for upland rice production in Northern Laos. 1. Soil physical properties, leaf SPAD and grain yield. *Field Crops Research*, 111(1–2), 81–84.

- Asra, R., Samarlina, R. A., & Silalahi, M. (2020). *Hormon Tumbuhan* (I. Jatmoko (ed.). Jakarta: UKI Press.
- Awad, Y. M., Lee, S. S., Kim, K. H., Ok, Y. S., & Kuzyakov, Y. (2018). Carbon and nitrogen mineralization and enzyme activities in soil aggregate-size classes: Effects of biochar, oyster shells, and polymers. *Chemosphere*, 198, 40–48.
- Badar, R., & Qureshi, S. A. (2014). Composted rice husk improves the growth and biochemical parameters of sunflower plants. *Journal of Botany*, 2014, 1–7.
- Balittanah (2009). *Analisis Kimia Tanah, Tanaman, Air, dan Pupuk*. Bogor: Balai Penelitian Tanah.
- Bandurska, H., & Stroiński, A. (2005). The effect of salicylic acid on barley response to water deficit. *Acta Physiol Plant*, 27, 379–386.
- Bashir, A., Rizwan, M., Zia ur Rehman, M., Zubair, M., Riaz, M., Qayyum, M. F., Alharby, H. F., Bamagoos, A. A., & Ali, S. (2020). Application of co-composted farm manure and biochar increased the wheat growth and decreased cadmium accumulation in plants under different water regimes. *Chemosphere*, 246, 1–10.
- Bates, L. S. (1973). Rapid Determination of Free Proline for Water-Stress Studies. *Plant and Soil*, 39, 205–207.
- Ben Ammar, H., Arena, D., Treccarichi, S., Di Bella, M. C., Marghali, S., Ficcadenti, N., Lo Scalzo, R., & Branca, F. (2023). The effect of water stress on the glucosinolate content and profile: a comparative study on roots and leaves of *Brassica oleracea* L. Crops. *Agronomy*, 13, 1–19.
- Bhat, R., & Vyas, D. (2019). Myrosinase: insights on structural, catalytic, regulatory, and environmental interactions. *Critical Reviews in Biotechnology*, 39(4), 508–523.
- Blanco-Canqui, H. (2017). Biochar and Soil Physical Properties. *Soil Science Society of America Journal*, 81(4), 678–771.
- Borsani, O., Valpuesta, V., & Botella, M. A. (2001). Evidence for a role of salicylic acid in the oxidative damage generated by NaCl and osmotic stress in *Arabidopsis* seedlings. *Plant Physiology*, 126(3), 1024–1030.
- Buckingham, A., & Whittingham, J. (2008). *Grow Vegetables*. United States: DK Publishing.
- Buraniya, S., Basant, N. K. G., Dadrwal, K., & Kakrallya, B. L. (2020). Study the Effect of PEG 6000 on *Brassica* genotypes germination and growth parameters under drought condition. *International Journal of Current Microbiology and Applied Sciences*, 11, 3840–3847.
- Butnan, S., Deenik, J. L., Toomsan, B., Antal, M. J., & Vityakon, P. (2015). Biochar characteristics and application rates affecting corn growth and properties of

- soils contrasting in texture and mineralogy. *Geoderma*, 237–238, 105–116.
- Campbell, N. A., & Reece, J. B. (2012). *Biology Jilid 1* (8th ed.). Jakarta: Penerbit Erlangga.
- Campos, P., Knicker, H., López, R., & De la Rosa, J. M. (2021). Application of biochar produced from crop residues on trace elements contaminated soils: Effects on soil properties, enzymatic activities and brassica rapa growth. *Agronomy*, 11(1394), 1–18.
- Cao, Y., & Pawłowski, A. (2013). Life cycle assessment of two emerging sewage sludge-to-energy systems: Evaluating energy and greenhouse gas emissions implications. *Artur*, 127, 81–91.
- Carvalho, M. T. M., Madari, B. E., Bastiaans, L., van Oort, P. A. J., Leal, W. G. O., Heinemann, A. B., da Silva, M. A. S., Maia, A. H. N., Parsons, D., & Meinke, H. (2016). Properties of a clay soil from 1.5 to 3.5 years after biochar application and the impact on rice yield. *Geoderma*, 276, 7–18.
- Casson, S. A., Chilley, P. M., Topping, J. F., Evans, I. M., Souter, M. A., & Lindsey, K. (2002). The POLARIS gene of *Arabidopsis* encodes a predicted peptide required for correct root growth and leaf vascular patterning. *Plant Cell*, 14(8), 1705–1721.
- Chai, C., & Subudhi, P. K. (2016). Comprehensive analysis and expression profiling of the OsLAX and OsABCB auxin transporter gene families in rice (*Oryza sativa*) under phytohormone stimuli and abiotic stresses. *Frontiers in Plant Science*, 7(593), 1–13.
- Chen, Z., Zheng, Z., Huang, J., Lai, Z., & Fan, B. (2009). Biosynthesis of salicylic acid in plants. *Plant Signaling and Behavior*, 4(6), 493–496.
- Cheng, C. H., Lehmann, J., Thies, J. E., Burton, S. D., & Engelhard, M. H. (2006). Oxidation of black carbon by biotic and abiotic processes. *Organic Geochemistry*, 37(11), 1477–1488.
- Chhajed, S., Mostafa, I., He, Y., Abou-Hashem, M., El-Domiaty, M., & Chen, S. (2020). Glucosinolate biosynthesis and the glucosinolate–myrosinase system in plant defense. *Agronomy*, 10 11, 1–25.
- Chhaya, Yadav, B., Jogawat, A., Gnanasekaran, P., Kumari, P., Lakra, N., Lal, S. K., Pawar, J., & Narayan, O. P. (2021). An overview of recent advancement in phytohormones-mediated stress management and drought tolerance in crop plants. *Plant Gene*, 25(100264), 1–12.
- Chilley, P. M., Casson, S. A., Tarkowski, P., Hawkins, N., Wang, K. L. C., Hussey, P. J., Beale, M., Ecker, J. R., Sandberg, G. K., & Lindsey, K. (2006). The POLARIS peptide of *Arabidopsis* regulates auxin transport root growth via effects on ethylene signaling. *Plant Cell*, 18(11), 3058–3072.
- Chini, A., Grant, J. J., Seki, M., Shinozaki, K., & Loake, G. J. (2004). Drought

tolerance established by enhanced expression of the CC-NBS-LRR gene, ADR1, requires salicylic acid, EDS1 and ABI1. *Plant Journal*, 38, 810–822.

Chowdhury, P. (2022). *Glucosinolates and Its Role in Mitigating Abiotic and Biotic Stress in Brassicaceae*. IntechOpen. <https://doi.org/10.5772/intechopen.102367>

Chun, J. H., Kim, N. H., Seo, M. S., Jin, M., Park, S. U., Arasu, M. V., Kim, S. J., & Al-Dhabi, N. A. (2018). Molecular characterization of glucosinolates and carotenoid biosynthetic genes in Chinese cabbage (*Brassica rapa* L. ssp. *pekinensis*). *Saudi Journal of Biological Sciences*, 25(1), 71–82.

Claoston, N., Samsuri, A. W., Ahmad Husni, M. H., & Mohd Amran, M. S. (2014). Effects of pyrolysis temperature on the physicochemical properties of empty fruit bunch and rice husk biochars. *Waste Management and Research*, 32(4), 331–339.

Cocetta, G., Mishra, S., Raffaelli, A., & Ferrante, A. (2018). Effect of heat root stress and high salinity on glucosinolates metabolism in wild rocket. *Journal of Plant Physiology*, 231(August), 261–270.

Colebrook, E. H., Thomas, S. G., Phillips, A. L., & Hedden, P. (2014). The role of gibberellin signalling in plant responses to abiotic stress. *Journal of Experimental Biology*, 217(1), 67–75.

Comas, L. H., Becker, S. R., Cruz, V. M. V., Byrne, P. F., & Dierig, D. A. (2013). Root traits contributing to plant productivity under drought. *Frontiers in Plant Science*, 4(442), 1–16.

Connolly, E. L., Sim, M., Travica, N., Marx, W., Beasy, G., Lynch, G. S., Bondonno, C. P., Lewis, J. R., Hodgson, J. M., & Blekkenhorst, L. C. (2021). Glucosinolates from cruciferous vegetables and their potential role in chronic disease: investigating the preclinical and clinical evidence. *Frontiers in Pharmacology*, 12, 1–12.

Coulombe, J., Villeneuve, S., Lamy, P., Yelle, S., Bélec, C., & Tremblay, N. (1998). Evaluation of soil and petiole sap nitrate quick tests for broccoli in Québec. *Acta Horti*, 506, 147–152.

Cutler, S. R., Rodriguez, P. L., Finkelstein, R. R., & Abrams, S. R. (2010). Absciscic acid: Emergence of a core signaling network. *Annual Review of Plant Biology*, 61, 651–679.

Dai, Z., Zhang, X., Tang, C., Muhammad, N., Wu, J., Brookes, P. C., & Xu, J. (2017). Potential role of biochars in decreasing soil acidification - A critical review. *Science of the Total Environment*, 581–582, 601–611.

Davies, Peter J. (2010). *Plant Hormones Biosynthesis, Signal Transduction, Action! Revised 3rd Edition*. Dordrecht: Springer.

de Ollas, C., Arbona, V., & Gómez-Cadenas, A. (2015). Jasmonoyl isoleucine

- accumulation is needed for abscisic acid build-up in roots of *Arabidopsis* under water stress conditions. *Plant Cell and Environment*, 38(10), 2157–2170.
- de Ollas, C., & Dodd, I. C. (2016). Physiological impacts of ABA–JA interactions under water-limitation. *Plant Molecular Biology*, 91(6), 641–650.
- Delker, C., Stenzel, I., B, H., Miersch, O., Feussner, I., & Wasternack, C. (2006). Jasmonate biosynthesis in *Arabidopsis thaliana*—enzymes, products, regulation. *Plant Biol*, 8, 297–306.
- DeLuca, T., MacKenzie, M., & Gundale, M. (2009). Biochar effects on soil nutrient transformations. In S. Lehman, J., Joseph (Ed.), *Biochar for Environmental Management* (pp. 251–270).
- Dempsey, D. A., Vlot, A. C., Wildermuth, M. C., & Klessig, D. F. (2011). Salicylic Acid Biosynthesis and Metabolism. *Arabidopsis Book*, 9, 1–24.
- Di Martino, C., Pizzuto, R., Pallotta, M. L., De Santis, A., & Passarella, S. (2006). Mitochondrial transport in proline catabolism in plants: The existence of two separate translocators in mitochondria isolated from durum wheat seedlings. *Planta*, 223(6), 1123–1133.
- Dobra, J., Motyka, V., Dobrev, P., Malbeck, J., Prasil, I. T., Haisel, D., Gaudinova, A., Havlova, M., Gubis, J., & Vankova, R. (2010). Comparison of hormonal responses to heat, drought and combined stress in tobacco plants with elevated proline content. *Journal of Plant Physiology*, 167(16), 1360–1370.
- Dominguez-Perles, R., Martinez-Ballesta, M. C., Riquelme, F., Carvajal, M., Garcia-Viguera, C., & Moreno, D. A. (2011). Novel varieties of broccoli for optimal bioactive components under saline stress. *Journal of the Science of Food and Agriculture*, 91(9), 1638–1647.
- Dong, D., Feng, Q., McGrouther, K., Yang, M., Wang, H., & Wu, W. (2015). Effects of biochar amendment on rice growth and nitrogen retention in a waterlogged paddy field. *Journal of Soils and Sediments*, 15(1), 153–162.
- dos Santos, I. C., De Almeida, A. A. F., Anher, D., Da Conceição, A. S., Pirovani, C. P., Pires, J. L., Valle, R. R., & Baligar, V. C. (2014). Molecular, physiological and biochemical responses of *Theobroma cacao* L. genotypes to soil water deficit. *PLoS ONE*, 9(12), 1–31.
- Downie, A., Crosky, A., & Munroe, P. (2009). Physical Properties of Biochar. In Johannes Lehmann & S. Josep (Eds.), *Biochar for Environmental Management: Science and Technology* (pp. 13–32).
- Egamberdieva, D., Wirth, S., Behrendt, U., Abd-Allah, E. F., & Berg, G. (2016). Biochar treatment resulted in a combined effect on soybean growth promotion and a shift in plant growth promoting rhizobacteria. *Frontiers in Microbiology*, 7(209), 1–11.
- Eom, S. H., Baek, S. A., Kim, J. K., & Hyun, T. K. (2018). Transcriptome analysis

in chinese cabbage (*Brassica rapa* ssp. *pekinensis*) provides the role of glucosinolate metabolism in response to drought stress. *Molecules*, 23(5).

- Fahey, J. W. (2015). Brassica: Characteristics and Properties. In *Encyclopedia of Food and Health* (3rd ed.). Amsterdam: Elsevier Ltd.
- Fang, Y., & Xiong, L. (2015). General mechanisms of drought response and their application in drought resistance improvement in plants. *Cellular and Molecular Life Sciences*, 72(4), 673–689.
- Farhangi-Abri, S., & Torabian, S. (2018). Biochar increased plant growth-promoting hormones and helped to alleviate salt stress in common bean seedlings. *Journal of Plant Growth Regulation*, 37(2), 591–601.
- Farooq, M., Hussain, M., Wahid, A., & Siddique, K. H. M. (2012). Drought Stress in Plants: An Overview. In R. Aroca (Ed.), *Plant Responses to Drought Stress: From Morphological to Molecular Features* (pp. 1–33). Heidelberg: Springer Berlin.
- Fichman, Y., Gerdes, S. Y., Kovács, H., Szabados, L., Zilberstein, A., & Csonka, L. N. (2015). Evolution of proline biosynthesis: Enzymology, bioinformatics, genetics, and transcriptional regulation. *Biological Reviews*, 90(4), 1065–1099.
- French, E., & Iyer-Pascuzzi, A. S. (2018). A role for the gibberellin pathway in biochar-mediated growth promotion. *Scientific Reports*, 8(5389), 1–10.
- Gagneul, D., Ainouche, A., Duhazé, C., Lugan, R., Larher, F. R., & Bouchereau, A. (2007). A reassessment of the function of the so-called compatible solutes in the halophytic plumbaginaceae *Limonium latifolium*. *Plant Physiology*, 144(3), 1598–1611.
- García-Ibañez, P., Sánchez-garcía, M., Sánchez-monedero, M. A., Luz, M., & Moreno, D. A. (2020). Olive tree pruning derived biochar increases glucosinolate concentrations in broccoli. *Scientia Horticulturae*, 267(109329), 109329.
- Geilfus, C.-M., Hasler, K., Witzel, K., Gerendás, J., & Mühling, K. H. (2016). Interactive effects of genotype and N/S-supply on glucosinolates and glucosinolate breakdown products in Chinese cabbage (*Brassica rapa* L. Ssp. *pekinensis*). *Journal of Applied Botany and Food Quality*, 89, 279–286.
- Ghars, M. A., Richard, L., Lefebvre-De Vos, D., Leprince, A. S., Parre, E., Bordenave, M., Abdelly, C., & Savouré, A. (2012). Phospholipases C and D modulate proline accumulation in *Thellungiella halophila*/salsuginea differently according to the severity of salt or hyperosmotic stress. *Plant and Cell Physiology*, 53(1), 183–192.
- Gomez, & Gomez. (2010). *Prosedur Statistika Untuk Penelitian Pertanian*. Jakarta: UI Press.

- Graber, E. R., Harel, Y. M., Kolton, M., Cytryn, E., Silber, A., David, D. R., Tsechansky, L., Borenshtein, M., & Elad, Y. (2010). Biochar impact on development and productivity of pepper and tomato grown in fertigated soilless media. *Plant and Soil*, 337(1), 481–496.
- Gujjar, R. S., & Supaibulwatana, K. (2019). The mode of cytokinin functions assisting plant adaptations to osmotic stresses. *Plants*, 8(542), 1–18.
- Gul, S., & Whalen, J. K. (2016). Biochemical cycling of nitrogen and phosphorus in biochar-amended soils. *Soil Biology and Biochemistry*, 103, 1–15.
- Gundale, M. J., & DeLuca, T. H. (2006). Temperature and source material influence ecological attributes of ponderosa pine and Douglas-fir charcoal. *Forest Ecology and Management*, 231(1–3), 86–93.
- Ha, S., Vankova, R., Yamaguchi-Shinozaki, K., Shinozaki, K., & Tran, L. S. P. (2012). Cytokinins: Metabolism and function in plant adaptation to environmental stresses. *Trends in Plant Science*, 17(3), 172–179.
- Hafeez, Y., Iqbal, S., Jabeen, K., Shahzad, S., Jahan, S., & Rasul, F. (2017). Effect of biochar application on seed germination and seedling growth of *Glycine max* (L.) merr. Under drought stress. *Pakistan Journal of Botany*, 49(Special Issue), 7–13.
- Halkier, B. A., & Gershenzon, J. (2006). Biology and biochemistry of glucosinolates. *Annual Review of Plant Biology*, 57, 303–333.
- Haryanto, E. (2007). *Sawi dan Selada*. Jakarta: Penebar Swadaya.
- Hasni, I., Yaakoubi, H., Hamdani, S., Tajmir-Riahi, H. A., & Carpentier, R. (2015). Mechanism of interaction of Al^{3+} with the proteins composition of photosystem II. *PLoS ONE*, 10(3), 1–23.
- Hayat, S., Hayat, Q., Alyemeni, M. N., Wani, A. S., Pichtel, J., & Ahmad, A. (2012). Role of proline under changing environments: A review. *Plant Signaling and Behavior*, 7(11), 1–11.
- Herath, H. M. S. K., Camps-Arbestain, M., Hedley, M. J., Kirschbaum, M. U. F., Wang, T., & van Hale, R. (2015). Experimental evidence for sequestering C with biochar by avoidance of CO₂ emissions from original feedstock and protection of native soil organic matter. *GCB Bioenergy*, 7(3), 512–526.
- Homagain, K., Shahi, C., Luckai, N., & Sharma, M. (2015). Life cycle environmental impact assessment of biochar-based bioenergy production and utilization in Northwestern Ontario, Canada. *Journal of Forestry Research*, 26(4), 799–809.
- Hopkins, W. G., & Hüner, N. P. A. (2009). *Introduction to plant physiology* (4th ed.). USA: John Wiley & Sons.
- Hou, Z. ., Du, G. ., & Sun, Z. . (2004). Preliminary Report of Returning Corn Straw into Soil on Soil Fertility. Rain Fed Crops. *Rain Fed Crops*, 24, 164–167.

- Hsiao, T. (1973). *Plant Responses to Water Stress. Annual Review of Plant Physiology*, 24(1), 519–570.
- Huang, B. (2000). Waterlogging Responses and Interaction with Temperature, Salinity, and Nutrients. In R. E. Wilkinson (Ed.), *Plant–Environmental Interaction* (pp. 263–281). Marcel Dekker Inc.
- Ilyas, M., Nisar, M., Khan, N., Hazrat, A., Khan, A. H., Hayat, K., Fahad, S., Khan, A., & Ullah, A. (2021). Drought Tolerance Strategies in Plants: A Mechanistic Approach. *Journal of Plant Growth Regulation*, 40, 926–944.
- IPCC Climate Change. (2007). *Climate change 2007: the physical science basis, vol 1009*. Cambridge: Cambridge University Press.
- Ippolito, M. P., Fasciano, C., d’Aquino, L., & Tommasi, F. (2011). Responses of antioxidant systems to lanthanum nitrate treatments in tomato plants during drought stress. *Plant Biosystems*, 145(1), 248–252.
- Irwan, A. W., & Wicaksono, F. Y. (2017). Comparations of Soybean ’s Leaf Area Measurement using Gravimetry , Regression , and Scanning. *Jurnal Kultivasi*, 16(3), 425–429.
- Ismail, Hamayun, M., Hussain, A., Iqbal, A., Khan, S. A., & Lee, I. J. (2020). *Aspergillus niger* boosted heat stress tolerance in sunflower and soybean via regulating their metabolic and antioxidant system. *Journal of Plant Interactions*, 15(1), 223–232.
- Jaiswal, A. K., Alkan, N., Elad, Y., Sela, N., Philosoph, A. M., Graber, E. R., & Frenkel, O. (2020). Molecular insights into biochar-mediated plant growth promotion and systemic resistance in tomato against fusarium crown and root rot disease. *Scientific Reports*, 10(13934).
- Jaiswal, A. K., Elad, Y., Paudel, I., Graber, E. R., Cytryn, E., & Frenkel, O. (2017). Linking the belowground microbial composition, diversity and activity to soilborne disease suppression and growth promotion of tomato amended with biochar. *Scientific Reports*, 7(44382), 1–17.
- Jaleel, C. A., Manivannan, P., Wahid, A., Farooq, M., Al-Juburi, H. J., Somasundaram, R., & Panneerselvam, R. (2009). Drought stress in plants: A review on morphological characteristics and pigments composition. *International Journal of Agriculture and Biology*, 11(1), 100–105.
- Javaid, T., Farooq, M. A., Akhtar, J., Saqib, Z. A., & Anwar-ul-Haq, M. (2019). Silicon nutrition improves growth of salt-stressed wheat by modulating flows and partitioning of Na⁺, Cl⁻ and mineral ions. *Plant Physiology and Biochemistry*, 141(January), 291–299.
- Jones Jr., J. B. (2012). *Plant Nutrition and Soil Fertility Manual*. Boca Raton: CRC Press.
- Jung, H., Lee, D. K., Choi, Y. Do, & Kim, J. K. (2015). OsIAA6, a member of the

rice Aux/IAA gene family, is involved in drought tolerance and tiller outgrowth. *Plant Science*, 236, 304–312.

Kalb, T., & Chang, L. (2005). *Suggested Cultural Practices for Headig Chinese Cabbage*. AVRDC - The World Vegetable Centre.

Kang, G. Z., Li, G. Z., Liu, G. Q., Xu, W., Peng, X. Q., Wang, C. Y., Zhu, Y. J., & Guo, T. C. (2013). Exogenous salicylic acid enhances wheat drought tolerance by influence on the expression of genes related to ascorbate-glutathione cycle. *Biologia Plantarum*, 57(4), 718–724.

Karam, D. S., Nagabovanalli, P., Sundara Rajoo, K., Fauziah Ishak, C., Abdu, A., Rosli, Z., Melissa Muharam, F., & Zulperi, D. (2022). An overview on the preparation of rice husk biochar, factors affecting its properties, and its agriculture application. *Journal of the Saudi Society of Agricultural Sciences*, 21(3), 149–159.

Kartika, Fadilah, L. N., & Lakitan, B. (2021). Growth responses and yield of cauliflower (*Brassica oleracea* var. *botrytis* L.) to the delayed transplanting and drought stress. *E3S Web of Conferences*, 306, 1–9.

Kavitha, B., Reddy, P. V. L., Kim, B., Lee, S. S., Pandey, S. K., & Kim, K. H. (2018). Benefits and limitations of biochar amendment in agricultural soils: A review. *Journal of Environmental Management*, 227, 146–154.

Ke, Q., Wang, Z., Ji, C. Y., Jeong, J. C., Lee, H. S., Li, H., Xu, B., Deng, X., & Kwak, S. S. (2015). Transgenic poplar expressing Arabidopsis YUCCA6 exhibits auxin-overproduction phenotypes and increased tolerance to abiotic stress. *Plant Physiology and Biochemistry*, 94, 19–27.

Khorram, M. S., Fatemi, A., Khan, M. A., Kiefer, R., & Jafarnia, S. (2018). Potential risk of weed outbreak by increasing biochar's application rates in slow-growth legume, lentil (*Lens culinaris* Medik.). *Journal of the Science of Food and Agriculture*, 98(6), 2080–2088.

Kido, É. A., Ferreira-Neto, J. R. C., da Silva, M. D., Santos, V. E. P., da Silva Filho, J. L. B., & Benko-Iseppon, A. M. (2019). smoprotectant-Related Genes in Plants Under Abiotic Stress: Expression Dynamics, In Silico Genome Mapping, and Biotechnology. In M. Hossain, V. Kumar, D. Burritt, M. Fujita, & P. Mäkelä (Eds.), *Osmoprotectant-Mediated Abiotic Stress Tolerance in Plants* (pp. 1–40). Springer, Cham.

Kim, H., Seomun, S., Yoon, Y., & Jang, G. (2021). Jasmonic acid in plant abiotic stress tolerance and interaction with abscisic acid. *Agronomy*, 11(9), 1–16.

Kim, J. I., Baek, D., Park, H. C., Chun, H. J., Oh, D. H., Lee, M. K., Cha, J. Y., Kim, W. Y., Kim, M. C., Chung, W. S., Bohnert, H. J., Lee, S. Y., Bressan, R. A., Lee, S. W., & Yun, D. J. (2013a). Overexpression of arabidopsis YUCCA6 in potato results in high-auxin developmental phenotypes and enhance. *Molecular Plant*, 6(2), 337–349.

- Kim, J., Jung, Y., Song, B., Bong, Y. S., Ryu, D. H., Lee, K. S., & Hwang, G. S. (2013b). Discrimination of cabbage (*Brassica rapa* ssp. *pekinensis*) cultivars grown in different geographical areas using ¹H NMR-based metabolomics. *Food Chemistry*, 137(1–4), 68–75.
- Kim, K. H., & Chung, S.-O. (2018). *Comparison of Plant Growth and Glucosinolates of Chinese Cabbage and Kale Crops under Three Cultivation Conditions*. 43(1), 30–36.
- Kim, Y.-Y., Oh, S. H., Pang, W., Li, X., Ji, S. J., Son, E., Han, S., Park, S., Soh, E., Kim, H., & Lim, Y. P. (2017). A review of the scientific names of chinese cabbage according to the international codes of nomenclature. *Horticultural Science and Technology*, 35(2), 165–169.
- Kolton, M., Graber, E. R., Tsehansky, L., Elad, Y., & Cytryn, E. (2016). Biochar-stimulated plant performance is strongly linked to microbial diversity and metabolic potential in the rhizosphere. *New Phytologist*, 213(3), 1393–1404.
- Kookana, R. S., Sarmah, A. K., Zwieten, L. V., Krull, E. V., & Singh, B. (2011). Chapter three - Biochar Application to Soil: Agronomic and Environmental Benefits and Unintended Consequences. In D. L. Sparks (Ed.), *Advances in Agronomy* (pp. 103–143). United States: Academic Press.
- Kramell, R., Miersch, O., Atzorn, R., Parthier, B., & Wasternack, C. (2000). Octadecanoid-derived alteration of gene expression and the “oxylipin signature” in stressed barley leaves. Implications for different signaling pathways. *Plant Physiology*, 123, 177–187.
- Ku, K. M., & Juvik, J. A. (2013). Environmental stress and methyl jasmonatemediated changes in flavonoid concentrations and antioxidant activity in broccoli florets and kale leaf tissues. *HortScience*, 48, 996–1002.
- Kulkarni, M. S., GovindMirgal, P., Bodhale, P. P., & Tande, S. N. (2014). Effect of rice husk ash on properties of concrete. *Journal of Civil Engineering and Environmental Technology*, 1(1), 26–29.
- Kumar, S., Sangwan, P., V, D. R. M., & Bidra, S. (2013). Utilization of rice husk and their ash : A Review. *Journal of Chemical and Environmental Sciences*, 1(5), 126–129.
- Kusumaningrum, I., Hastuti, R. B., & Haryanti, S. (2007). Pengaruh perasan *Sargassum crassifolium* dengan konsentrasi yang berbeda terhadap pertumbuhan tanaman kedelai (*Glycine max* (L) Merrill). *Buletin Anatomi Dan Fisiologi*, XV(2), 17–13.
- Kusumawati, K., Muhartini, S., & Rogomulyo, R. (2015). Pengaruh konsentrasi dan frekuensi pemberian limbah tahu terhadap pertumbuhan dan hasil bayam (*Amaranthus tricolor* L.). *Vegetalika*, 4(2), 48–62.
- Kuzyakov, Y., Bogomolova, I., & Glaser, B. (2014). Biochar stability in soil: Decomposition during eight years and transformation as assessed by

- compound-specific ^{14}C analysis. *Soil Biology and Biochemistry*, 70, 229–236.
- Kwapinski, W., Byrne, C. M. P., Kryachko, E., Wolfram, P., Adley, C., Leahy, J. J., Novotny, E. H., & Hayes, M. H. B. (2010). Biochar from biomass and waste. *Waste and Biomass Valorization*, 1(2), 177–189.
- La, V. H., Lee, B. R., Zhang, Q., Park, S. H., Islam, M. T., & Kim, T. H. (2019). Salicylic acid improves drought-stress tolerance by regulating the redox status and proline metabolism in *Brassica rapa*. *Horticulture Environment and Biotechnology*, 60(1), 31–40.
- Labola, Y. A., & Puspita, D. (2017). Peran Antioksidan Karotenoid Penangkal Radikal Bebas Penyebab Berbagai Penyakit. *Farmasetika.Com (Online)*, 2(2), 12.
- Laxa, M., Liebthal, M., Telman, W., Chibani, K., & Dietz, K. J. (2019). The role of the plant antioxidant system in drought tolerance. *Antioxidants*, 8(94), 1–31.
- Lebrun, M., Miard, F., Nandillon, R., Scippa, G. S., Bourgerie, S., & Morabito, D. (2019). Biochar effect associated with compost and iron to promote Pb and As soil stabilization and *Salix viminalis* L. growth. *Chemosphere*, 222, 810–822.
- Lee, J., Nam, J., Park, H. C., Na, G., Miura, K., Jin, J. B., Yoo, C. Y., Baek, D., Kim, D. H., Jeong, J. C., Kim, D., Lee, S. Y., Salt, D. E., Mengiste, T., Gong, Q., Ma, S., Bohnert, H. J., Kwak, S. S., Bressan, R. A., Yun, D. J. (2007). Salicylic acid-mediated innate immunity in *Arabidopsis* is regulated by SIZ1 SUMO E3 ligase. *Plant Journal*, 49(1), 79–90.
- Lee, M. K., Chun, J. H., Byeon, D. H., Chung, S. O., Park, S. U., Park, S., Arasu, M. V., Al-Dhabi, N. A., Lim, Y. P., & Kim, S. J. (2014). Variation of glucosinolates in 62 varieties of Chinese cabbage (*Brassica rapa* L. ssp. *pekinensis*) and their antioxidant activity. *Lwt*, 58(1), 93–101.
- Lehmann, Johannes. (2007). A handful of carbon. *Nature*, 447, 143–144.
- Lehmann, Johannes, da Silva Jr., J. P., Steiner, C., Nehls, T., Zech, W., & Glaser, B. (2003). Nutrient availability and leaching in an archaeological Anthrosol and a Ferralsol of the Central Amazon basin: fertilizer, manure and charcoal amendments. *Plant and Soil* 249:, 249, 343–357.
- Lehmann, Jörg, Atzorn, R., Brückner, C., Reinbothe, S., Leopold, J., Wasternack, C., & Parthier, B. (1995). Accumulation of jasmonate, abscisic acid, specific transcripts and proteins in osmotically stressed barley leaf segments. *Planta*, 197, 156–162.
- Leng, L., Huang, H., Li, H., Li, J., & Zhou, W. (2019). Biochar stability assessment methods: A review. *Science of the Total Environment*, 647, 210–222.
- Lewis, D. R., Ramirez, M. V., Miller, N. D., Vallabhaneni, P., Keith Ray, W., Helm, R. F., Winkel, B. S. J., & Muday, G. K. (2011). Auxin and ethylene induce flavonol accumulation through distinct transcriptional networks. *Plant*

Physiology, 156(1), 144–164.

- Li, J., Sima, W., Ouyang, B., Wang, T., Ziaf, K., Luo, Z., Liu, L., Li, H., Chen, M., Huang, Y., Feng, Y., Hao, Y., & Ye, Z. (2012). Tomato SIDREB gene restricts leaf expansion and internode elongation by downregulating key genes for gibberellin biosynthesis. *Journal of Experimental Botany*, 63(18), 6407–6420.
- Li, W., Herrera-Estrella, L., & Tran, L. S. P. (2016). The Yin-Yang of Cytokinin Homeostasis and drought acclimation/adaptation. *Trends in Plant Science*, 21(7), 548–550.
- Li, Y., Zhao, H., Duan, B., Korpelainen, H., & Li, C. (2011). Effect of drought and ABA on growth, photosynthesis and antioxidant system of *Cotinus coggygia* seedlings under two different light conditions. *Environmental and Experimental Botany*, 71(1), 107–113.
- Liao, F., Yang, L., Qiang, L., Li, Y.-R., Yang, L.-T., Anas, M., & Huang, D.-L. (2018). Characteristics and inorganic N holding ability of biochar derived from the pyrolysis of agricultural and forestal residues in the southern China. *Journal of Analytical and Applied Pyrolysis*, 134, 544–551.
- Liu, D, Pei, Z. F., Neem, M. S., Ming, D. ., Liu, B., Khan, F., & Zhou, W. J. (2011). 5-aminolevulinic acid activates antioxidative defence system and seedling growth in *Brassica napus* L. under water-deficit stress. *J. Agronomy & Crop Science*, 197, 284–295.
- Liu, Di, Ding, Z., Ali, E. F., Kheir, A. M. S., Eissa, M. A., & Ibrahim, O. H. M. (2021). Biochar and compost enhance soil quality and growth of roselle (*Hibiscus sabdariffa* L.) under saline conditions. *Scientific Reports*, 11(8739), 1–11.
- Liu, J., Mehdi, S., Topping, J., Tarkowski, P., & Lindsey, K. (2010). Modelling and experimental analysis of hormonal crosstalk in *Arabidopsis*. *Molecular Systems Biology*, 6(373), 1–13.
- Liu, P., Xu, Z. S., Pan-Pan, L., Hu, D., Chen, M., Li, L. C., & Ma, Y. Z. (2013). A wheat PI4K gene whose product possesses threonine autophosphorylation activity confers tolerance to drought and salt in *Arabidopsis*. *Journal of Experimental Botany*, 64(10), 2915–2927.
- Lo, S. F., Ho, T. H. D., Liu, Y. L., Jiang, M. J., Hsieh, K. T., Chen, K. T., Yu, L. C., Lee, M. H., Chen, C. Y., Huang, T. P., Kojima, M., Sakakibara, H., Chen, L. J., & Yu, S. M. (2017). Ectopic expression of specific GA2 oxidase mutants promotes yield and stress tolerance in rice. *Plant Biotechnology Journal*, 15(7), 850–864.
- López-Berenguer, C., Martínez-Ballesta, M. C., García-Viguera, C., & Carvajal, M. (2008). Leaf water balance mediated by aquaporins under salt stress and associated glucosinolate synthesis in broccoli. *Plant Sci.*, 174, 321–328.
- López-Pérez, L., Martínez-Ballesta, M. C., Maurel, C., & Carvajal, M. (2009).

- Changes in plasma membrane lipids, aquaporins and proton pump of broccoli roots, as an adaptation mechanism to salinity. *Phytochemistry*, 70, 492–500.
- Lugtenberg, B., & Kamilova, F. (2009). Plant-growth-promoting rhizobacteria. *Annual Review of Microbiology*, 63, 541–556.
- Luo, L. J. (2010). Breeding for water-saving and drought-resistance rice (WDR) in China. *Journal of Experimental Botany*, 61(13), 3509–3517.
- Lynch, J. P. (2018). Rightsizing root phenotypes for drought resistance. *Journal of Experimental Botany*, 69(13), 3279–3292.
- Maftu'ah, E., & Nursyamsi, D. (2015). *Potensi berbagai bahan organik rawa sebagai sumber biochar. 1*, 776–781.
- Mahajan, S., & Tuteja, N. (2005). Cold, salinity and drought stresses: An overview. *Archives of Biochemistry and Biophysics*, 444(2), 139–158.
- Malik, Z., Afzal, S., Dawood, M., Abbasi, G. H., Khan, M. I., Kamran, M., Zhran, M., Hayat, M. T., Aslam, M. N., & Rafay, M. (2022). Exogenous melatonin mitigates chromium toxicity in maize seedlings by modulating antioxidant system and suppresses chromium uptake and oxidative stress. *Environmental Geochemistry and Health*, 44(5), 1451–1469.
- Manavalan, L. P., Guttikonda, S. K., Phan Tran, L. S., & Nguyen, H. T. (2009). Physiological and molecular approaches to improve drought resistance in soybean. *Plant and Cell Physiology*, 50(7), 1260–1276.
- Manivannan, P., Jaleel, C. A., Somasundaram, R., & Panneerselvam, R. (2008). Osmoregulation and antioxidant metabolism in drought-stressed *Helianthus annuus* under triadimefon drenching. *Comptes Rendus - Biologies*, 331(6), 418–425.
- (2017). Alleviation of Adverse Effect of drought stress on soybean (*Glycine max*. L.) by using poultry litter biochar. *Bangladesh Agronomy Journal*, 19(2), 61–69.
- Mano, Y., & Nemoto, K. (2012). The pathway of auxin biosynthesis in plants. *Journal of Experimental Botany*, 63(8), 2853–2872.
- Mansoor, S., Kour, N., Manhas, S., Zahid, S., Wani, O. A., Sharma, V., Wijaya, L., Alyemeni, M. N., Alsahli, A. A., El-Serehy, H. A., Paray, B. A., & Ahmad, P. (2021). Biochar as a tool for effective management of drought and heavy metal toxicity. *Chemosphere*, 271, 1–10.
- Martínez-Ballesta, M. del C., Moreno, D. A., & Carvajal, M. (2013). The physiological importance of glucosinolates on plant response to abiotic stress in *Brassica*. *International Journal of Molecular Sciences*, 14(6), 11607–11625.
- Martínez-Gómez, Á., Poveda, J., & Escobar, C. (2022). Overview of the use of biochar from main cereals to stimulate plant growth. *Frontiers in Plant*

Science, 13(August), 1–17.

- Martins, M., Sousa, B., Lopes, J., Soares, C., Machado, J., Carvalho, S., Fidalgo, F., & Teixeira, J. (2020). *Diclofenac shifts the role of root glutamine synthetase and glutamate dehydrogenase for maintaining nitrogen assimilation and proline production at the expense of shoot carbon reserves in Solanum lycopersicum L.* 27, 29130–29142.
- Maxwell, B. B., & Kieber, J. J. (2010). Cytokinin Signal Transduction. In P.J Davies (Ed.), *Plant Hormones*. Dordrecht: Springer.
- Mehmood, M., Pérez-Llorca, M., Casadesús, A., Farrakh, S., & Munné-Bosch, S. (2021). Leaf size modulation by cytokinins in sesame plants. *Plant Physiology and Biochemistry*, 167(September), 763–770.
- Misra, B. B., Acharya, B. R., Granot, D., Assmann, S. M., & Chen, S. (2015). The guard cell metabolome: Functions in stomatal movement and global food security. *Frontiers in Plant Science*, 6, 1–13.
- Miura, K., Okamoto, H., Okuma, E., Shiba, H., Kamada, H., Hasegawa, P. M., & Murata, Y. (2013). SIZ1 deficiency causes reduced stomatal aperture and enhanced drought tolerance via controlling salicylic acid-induced accumulation of reactive oxygen species in Arabidopsis. *Plant Journal*, 73(1), 91–104.
- Montoya, D., Fernández, J. A., Franco, J. A., & del Carmen Martínez Ballesta, M. (2022). Enriched-biochar application increases broccoli nutritional and phytochemical content without detrimental effect on yield. *Journal of the Science of Food and Agriculture*, 102(15), 7353–7362.
- Munemasa, S., Hauser, F., Park, J., Waadt, R., Brandt, B., & Schroeder, J. I. (2015). Mechanisms of abscisic acid-mediated control of stomatal aperture. *Current Opinion in Plant Biology*, 28, 154–162.
- Ndekei, A., Gitita, M., Njomo, N., & Mbui, D. (2021). Synthesis and characterization of rice husk biochar and its application in the adsorption studies of lead and copper. *International Research Journal of Pure and Applied Chemistry*, 22(4), 36–50.
- Niether, W., Glawe, A., Pfohl, K., Adamtey, N., Schneider, M., Karlovsky, P., & Pawelzik, E. (2020). The effect of short-term vs. long-term soil moisture stress on the physiological response of three cocoa (*Theobroma cacao* L.) cultivars. *Plant Growth Regulation*, 92(2), 295–306.
- Nir, I., Moshelion, M., & Weiss, D. (2014). The Arabidopsis GIBBERELLIN METHYL TRANSFERASE 1 suppresses gibberellin activity, reduces whole-plant transpiration and promotes drought tolerance in transgenic tomato. *Plant, Cell and Environment*, 37, 113–123.
- Nishiyama, R., Watanabe, Y., Fujita, Y., Le, D. T., Kojima, M., Werner, T., Vankova, R., Yamaguchi-Shinozaki, K., Shinozaki, K., Kakimoto, T.,

- Sakakibara, H., Schmülling, T., & Tran, L. S. P. (2011). Analysis of cytokinin mutants and regulation of cytokinin metabolic genes reveals important regulatory roles of cytokinins in drought, salt and abscisic acid responses, and abscisic acid biosynthesis. *Plant Cell*, 23(6), 2169–2183.
- Nishiyama, R., Watanabe, Y., Leyva-Gonzalez, M. A., Van Ha, C., Fujita, Y., Tanaka, M., Seki, M., Yamaguchi-Shinozaki, K., Shinozaki, K., Herrera-Estrella, L., & Tran, L. S. P. (2013). Arabidopsis AHP2, AHP3, and AHP5 histidine phosphotransfer proteins function as redundant negative regulators of drought stress response. *Proceedings of the National Academy of Sciences of the United States of America*, 110(12), 4840–4845.
- Nurjanaty, N., Linda, R., & Mukarlina, M. (2019). Pengaruh cekaman air dan pemberian pupuk daun terhadap pertumbuhan tanaman sawi (*Brassica juncea* L.). *Jurnal Protobiont*, 8(3), 6–11.
- Nwajiaku, I. M., Olanrewaju, J. S., Sato, K., Tokunari, T., Kitano, S., & Masunaga, T. (2018). Change in nutrient composition of biochar from rice husk and sugarcane bagasse at varying pyrolytic temperatures. *International Journal of Recycling of Organic Waste in Agriculture*, 7(4), 269–276.
- Pan, G.-X., Zhang, A.-F., Zou, J.-W., Li, L.-Q., Zhang, X.-H., & Zheng, J.-W. (2010). Biochar from agro-byproducts used as amendment to croplands: An option for low carbon agriculture. *Journal of Ecology and Rural Environment*, 26(4), 394–400.
- Patanè, C., Scordia, D., Testa, G., & Cosentino, S. . (2016). Physiological screening for drought tolerance in Mediterranean long-storage tomato. *Plant Sci.*, 249, 25–34.
- Pavlović, I., Petřík, I., Tarkowská, D., Lepeduš, H., Vujčić Bok, V., Radić Brkanac, S., Novák, O., & Salopek-Sondi, B. (2018). Correlations between Phytohormones and drought tolerance in selected brassica crops: Chinese cabbage, white cabbage and kale. *International Journal of Molecular Sciences*, 19(2866), 1–23.
- Pieterse, C. M. J., Leon-Reyes, A., Van Der Ent, S., & Van Wees, S. C. M. (2009). Networking by small-molecule hormones in plant immunity. *Nature Chemical Biology*, 5(5), 308–316.
- Podda, A., Pollastri, S., Bartolini, P., Pisuttu, C., Pellegrini, E., Nali, C., Cencetti, G., Michelozzi, M., Frassinetti, S., Giorgetti, L., Fineschi, S., Del Carratore, R., & Maserti, B. (2019). Drought stress modulates secondary metabolites in *Brassica oleracea* L. convar. *acephala* (DC) Alef, var. *sabellica* L. *Journal of the Science of Food and Agriculture*, 99(12), 5533–5540.
- Powlson, D. S., Whitmore, A. P., & Goulding, K. W. T. (2011). Soil carbon sequestration to mitigate climate change: a critical re-examination to identify the true and the false. *European Journal of Soil Science*, 62(1), 42–55.

- Prerostova, S., Dobrev, P. I., Gaudinova, A., Hosek, P., Soudek, P., Knirsch, V., & Vankova, R. (2017). Hormonal dynamics during salt stress responses of salt-sensitive *Arabidopsis thaliana* and salt-tolerant *Thellungiella salsuginea*. *Plant Science*, 264, 188–198.
- Prieto, M. A., López, C. J., & Simal-Gandara, J. (2019). Glucosinolates: Molecular structure, breakdown, genetic, bioavailability, properties and healthy and adverse effects. *Advances in Food and Nutrition Research*, 90, 305–350.
- Qin, H. Z. (2012). *Study on Properties of Biochar Made from Household Waste*. Nanjing: Nanjing Agricultural University.
- Rao, G. N. (2007). *Statistics for Agricultural Sciences* (Second). Hyderabad: BS Publications.
- Raza, A., Hafeez, M. B., Zahra, N., Shaukat, K., Umbreen, S., Tabassum, J., Charagh, S., Khan, R. S. A., & Hasanuzzaman, M. (2020). The Plant Family Brassicaceae: Introduction, Biology, And Importance. In M. Hasanuzzaman (Ed.), *The Plant Family Brassicaceae: Biology and Physiological Responses to Environmental Stresses* (pp. 1–43). Singapore: Springer.
- Riemann, M., Dhakarey, R., Hazman, M., Miro, B., Kohli, A., & Nick, P. (2015). Exploring jasmonates in the hormonal network of drought and salinity responses. *Frontiers in Plant Science*, 6(1077), 1–16.
- Rivero, R. M., Kojima, M., Gepstein, A., Sakakibara, H., Mittler, R., Gepstein, S., & Blumwald, E. (2007). Delayed leaf senescence induces extreme drought tolerance in a flowering plant. *Proceedings of the National Academy of Sciences of the United States of America*, 104(49), 19631–19636.
- Rosa, E. A. S., Heaney, R. K., Fenwick, G. R., & Portas, C. A. . (1997). Glucosinolates in crop plants. *Hortic. Rev.*, 19, 99–215.
- Rouphael, Y., Cardarelli, Mariateresa Schwarz, D., Franken, P., & Colla, G. (2012). Effects of Drought on Nutrient Uptake and Assimilation in Vegetable Crops. In R. Aorca (Ed.), *Plant Responses to Drought Stress: From Morphological to Molecular Features* (pp. 171–195).
- Rowe, J. H., Topping, J. F., Liu, J., & Lindsey, K. (2016). Absciscic acid regulates root growth under osmotic stress conditions via an interacting hormonal network with cytokinin, ethylene and auxin. *New Phytologist*, 211(1), 225–239.
- Safari, S., Nazari, F., Vafaei, Y., & Teixeira da Silva, J. A. (2022). Impact of rice husk biochar on drought stress tolerance in perennial ryegrass (*Lolium perenne* L.). *Journal of Plant Growth Regulation*, 42(2), 1-17.
- Sahebi, M., Hanafi, M. M., Rafii, M. Y., Mahmud, T. M. M., Azizi, P., Osman, M., Abiri, R., Taheri, S., Kalhori, N., Shabanimofrad, M., Miah, G., & Atabaki, N. (2018). Improvement of drought tolerance in rice (*Oryza sativa* L.): genetics, genomic tools, and the WRKY gene family. *BioMed Research International*,

2018.

- Sahin, U., Ekinici, M., Ors, S., Turan, M., Yildiz, S., & Yildirim, E. (2018). Effects of individual and combined effects of salinity and drought on physiological, nutritional and biochemical properties of cabbage (*Brassica oleracea* var. *capitata*). *Scientia Horticulturae*, 240(June), 196–204.
- Saini, S., Sharma, I., Kaur, N., & Pati, P. K. (2013). Auxin: A master regulator in plant root development. *Plant Cell Reports*, 32(6), 741–757.
- Salehin, M., Li, B., Tang, M., Katz, E., Song, L., Ecker, J. R., Kliebenstein, D. J., & Estelle, M. (2019). Auxin-sensitive Aux/IAA proteins mediate drought tolerance in *Arabidopsis* by regulating glucosinolate levels. *Nature Communications*, 10(1), 1–9.
- Salvi, P., Manna, M., Kaur, H., Thakur, T., Gandass, N., Bhatt, D., & Muthamilarasan, M. (2021). Phytohormone signaling and crosstalk in regulating drought stress response in plants. *Plant Cell Reports*, 40(8), 1305–1329.
- Sánchez-Rodríguez, E., del Mar Rubio-Wilhelmi, M., Cervilla, L. M., Blasco, B., Rios, J. J., Leyva, R., Romero, L., & Ruiz, J. M. (2010). Study of the ionome and uptake fluxes in cherry tomato plants under moderate water stress conditions. *Plant and Soil*, 335(1), 339–347.
- Schreiner, M., Beyene, B., Krumben, A., & Stutzel, H. (2009). Ontogenetic changes of 2-propenyl and 3-indolylmethyl glucosinolates in *Brassica carinata* leaves as affected by water supply. *J. Agric. Food Chem.*, 57, 7259–7263.
- Seo, P. J., & Park, C. M. (2010). MYB96-mediated abscisic acid signals induce pathogen resistance response by promoting salicylic acid biosynthesis in *Arabidopsis*. *New Phytologist*, 186(2), 471–483.
- Seo, P. J., Xiang, F., Qiao, M., Park, J. Y., Lee, Y. N., Kim, S. G., Lee, Y. H., Park, W. J., & Park, C. M. (2009). The MYB96 transcription factor mediates abscisic acid signaling during drought stress response in *Arabidopsis*. *Plant Physiology*, 151(1), 275–289.
- Sharma, A., Kumar, V., Shahzad, B., Ramakrishnan, M., Singh Sidhu, G. P., Bali, A. S., Handa, N., Kapoor, D., Yadav, P., Khanna, K., Bakshi, P., Rehman, A., Kohli, S. K., Khan, E. A., Parihar, R. D., Yuan, H., Thukral, A. K., Bhardwaj, R., & Zheng, B. (2020). Photosynthetic response of plants under different abiotic stresses: A Review. *Journal of Plant Growth Regulation*, 39(2), 509–531.
- Shashi, M., Mannan, M., Islam, M., & Rahman, M. (2018). Impact of rice husk biochar on growth, water relations and yield of maize (*Zea mays* L.) under Drought Condition. *The Agriculturists*, 16(2), 93–101.
- Shawon, R. A., Kang, B. S., Kim, H. C., Lee, S. G., Kim, S. K., Lee, H. J., Bae, J. H., & Ku, Y. G. (2018). Changes in free amino acid, carotenoid, and proline

content in chinese cabbage (*Brassica rapa* subsp. *pekinensis*) in response to drought stress. *Korean Journal of Plant Resources*, 31(6), 622–633.

- Shawon, R. A., Kang, B. S., Lee, S. G., Kim, S. K., Ju Lee, H., Katrich, E., Gorinstein, S., & Ku, Y. G. (2020). Influence of drought stress on bioactive compounds, antioxidant enzymes and glucosinolate contents of Chinese cabbage (*Brassica rapa*). *Food Chemistry*, 308(October 2019), 125657.
- Shi, W., Ju, Y., Bian, R., Li, L., Joseph, S., Mitchell, D. R. G., Munroe, P., Taherymoosavi, S., & Pan, G. (2020). Biochar bound urea boosts plant growth and reduces nitrogen leaching. *Science of the Total Environment*, 701, 134424.
- Singh, B. (2018). Rice husk ash. In *Waste and Supplementary Cementitious Materials in Concrete: Characterisation, Properties and Applications*. USA: Elsevier Ltd.
- Singh, V. P., Prasad, S. M., Munné-Bosch, S., & Müller, M. (2017). Editorial: Phytohormones and the regulation of stress tolerance in plants: Current status and future directions. *Frontiers in Plant Science*, 8(October), 1–2.
- Siregar, B. (2017). Analisa kadar C-organik dan perbandingan C/N tanah di lahan tambak Kelurahan Sicanang Kecamatan Medan Belawan. *Jurnal Warta*, 53(1), 1–14.
- Soengas, P., Rodríguez, V. M., Velasco, P., & Cartea, M. E. (2018). Effect of temperature stress on antioxidant defenses in *Brassica oleracea*. *ACS Omega*, 3(5), 5237–5243.
- Southavong, S., Ismail, M., Preston, T., Saud, H., & Ismail, R. (2018). Effects of pyrolysis temperature and residence time on rice straw-derived biochar for soil application. *International Journal of Plant & Soil Science*, 23(3), 1–11.
- Stepanova, A. N., Yun, J., Likhacheva, A. V., & Alonso, J. M. (2007). Multilevel interactions between ethylene and auxin in *Arabidopsis* roots. *Plant Cell*, 19, 2169–2185.
- Subbarao, G. V., Nam, N. H., Chauhan, Y. S., & Johansen, C. (2000). Osmotic adjustment, water relations and carbohydrate remobilization in pigeonpea under water deficits. *Journal of Plant Physiology*, 157(6), 651–659.
- Sun, J., Yang, J., & Yu, T. (2010). Effects of drought stress simulated by PEG on glucosinolates content in *Arabidopsis thaliana*. *Agricultural Science & Technology-Hunan*, 11(3), 20–24.
- Taiz, L., Zeiger, E., Moller, I. M., & Murphy, A. (2015). *Plant Physiology and Development* (6th ed.). Sunderland: Sinauer Associates, Inc.
- Tan, Z., Lin, C. S. K., Ji, X., & Rainey, T. J. (2017). Returning biochar to fields: A review. *Applied Soil Ecology*, 116, 1–11.
- Telaumbanua, M., Purwantana, B., Sutiarso, L., & Falah, M. A. F. (2016). Studi pola pertumbuhan tanaman sawi (*Brassica rapa* var. *parachinensis* L.)

- hidroponik di dalam greenhouse terkontrol. *AGRITECH*, 36(1), 104–110.
- Thảo, V. T. M., Khánh, N. M., Nguyễn, N. T. H., Anh, T. T., Niệm, P. T. Á., & Đức, N. T. (2021). Ảnh Hưởng Của Nhiệt Độ Nhiệt Phân Đến Tính Chất Hóa Lý Của Than Sinh Học Từ Trấu. *Tạp Chí Khoa Học Đại Học Mở Thành Phố Hồ Chí Minh - Kỹ Thuật Và Công Nghệ*, 16(1), 126–141.
- Tian, J., Wang, J., Dippold, M., Gao, Y., Blagodatskaya, E., & Kuzyakov, Y. (2016). Biochar affects soil organic matter cycling and microbial functions but does not alter microbial community structure in a paddy soil. *Science of The Total Environment*, 556, 88–97.
- Ty, C., Sina, V., Borin, K., & Preston, T. R. (2013). Effect of different levels of biochar on the yield and nutritive value of celery cabbage (*Brassica chinensis* var), Chinese cabbage (*Brassica pekinensis*), Mustard green (*Brassica juncea*) and water spinach (*Ipomoea aquatica*). *Livestock Research for Rural Development*, 25(1).
- Ullah, A., Manghwar, H., Shaban, M., Khan, A. H., Akbar, A., Ali, U., Ali, E., & Fahad, S. (2018). Phytohormones enhanced drought tolerance in plants: a coping strategy. *Environmental Science and Pollution Research*, 25(33), 33103–33118.
- van Zwieten, L., Kimber, S., Morris, S., Chan, K. Y., Downie, A., Rust, J., Joseph, S., & Cowie, A. (2010). Effects of biochar from slow pyrolysis of papermill waste on agronomic performance and soil fertility. *Plant and Soil*, 327(1), 235–246.
- Verdiana, M. A., Sebayang, H. T., & Sumarni, T. (2016). Pengaruh Berbagai dosis biochar sekam padi dan pupuk NPK Terhadap pertumbuhan dan hasil tanaman jagung (*Zea mays* L.). *Jurnal Produksi Tanaman*, 4, 611–616.
- Vieira, F. R., Romero Luna, C. M., Arce, G. L. A. F., & Ávila, I. (2020). Optimization of slow pyrolysis process parameters using a fixed bed reactor for biochar yield from rice husk. *Biomass and Bioenergy*, 132(105412), 1–11.
- Viger, M., Hancock, R. D., Miglietta, F., & Taylor, G. (2015). More plant growth but less plant defence? First global gene expression data for plants grown in soil amended with biochar. *GCB Bioenergy*, 7(4), 658–672.
- Viger, M., Smith, H. K., Cohen, D., Dewoody, J., Trewin, H., Steenackers, M., Bastien, C., & Taylor, G. (2016). Adaptive mechanisms and genomic plasticity for drought tolerance identified in European black poplar (*Populus nigra* L.). *Tree Physiology*, 36(7), 909–928.
- Vithanage, M., Rajapaksha, A. U., Zhang, M., Thiele-Bruhn, S., Lee, S. S., & Ok, Y. S. (2015). Acid-activated biochar increased sulfamethazine retention in soils. *Environmental Science and Pollution Research*, 22(3), 2175–2186.
- Vogel, J. P., Woeste, K. E., Theologis, A., & Kieber, J. J. (1998). Recessive and dominant mutations in the ethylene biosynthetic gene ACS5 of Arabidopsis

confer cytokinin insensitivity and ethylene overproduction, respectively. *Proceedings of Notes SI Construction of Hormonal Crosstalk Networks under Osmotic Stress Conditions. the National Academy of Sciences*, 4766–4771.

- Wang, C., Yang, A., Yin, H., & Zhang, J. (2008). Influence of water stress on endogenous hormone contents and cell damage of maize seedlings. *Journal of Integrative Plant Biology*, 50(4), 427–434.
- Wang, D., Jiang, P., Zhang, H., & Yuan, W. (2020a). Biochar production and applications in agro and forestry systems: A review. *Science of the Total Environment*, 723.
- Wang, J., Song, L., Gong, X., Xu, J., & Li, M. (2020b). Functions of jasmonic acid in plant regulation and response to abiotic stress. *International Journal of Molecular Sciences*, 21(4).
- Wang, Y., Wu, F., Bai, J., & He, Y. (2014). BrpSPL9 (*Brassica rapa* ssp. *pekinensis* SPL9) controls the earliness of heading time in Chinese cabbage. *Plant Biotechnology Journal*, 12(3), 312–321.
- Warnock, D. D., Lehmann, J., Kuypers, T. W., & Rillig, M. C. (2007). Mycorrhizal responses to biochar in soil - Concepts and mechanisms. *Plant and Soil*, 300, 9–20.
- Wasternack, C. (2007). Jasmonates: an update on biosynthesis, signal transduction and action in plant stress response, growth and development. *Ann Bot*, 100(4), 681–697.
- Werner, T., Nehnevajova, E., Köllmer, I., Novák, O., Stmad, M., Krämer, U., & Schmülling, T. (2010). Root-specific reduction of cytokinin causes enhanced root growth, drought tolerance, and leaf mineral enrichment in *Arabidopsis* and tobacco. *Plant Cell*, 22(12),
- Wolters, H., & Jürgens, G. (2009). Survival of the flexible: Hormonal growth control and adaptation in plant development. *Nature Reviews Genetics*, 10(5), 305–317.
- Wu, Y., Zhang, D., Chu, J. Y., Boyle, P., Wang, Y., Brindle, I. D., De Luca, V., & Després, C. (2012). The *Arabidopsis* NPR1 protein is a receptor for the plant defense hormone salicylic acid. *Cell Reports*, 1(6), 639–647.
- Xiong, S., Wang, Y., Chen, Y., Gao, M., Zhao, Y., & Wu, L. (2022). Effects of drought stress and rehydration on physiological and biochemical properties of four oak species in China. *Plants*, 11(5), 1–17.
- Xu, C., & Leskovar, D. I. (2014). Growth, Physiology and Yield Responses of Cabbage to Deficit Irrigation. *Horticultural Science*, 41(3), 138–146.
- Xu, Y., Burgess, P., Zhang, X., & Huang, B. (2016). Enhancing cytokinin synthesis by overexpressing ipt alleviated drought inhibition of root growth through activating ROS-scavenging systems in *Agrostis stolonifera*. *J. Exp. Bot.*, 67,

1979–1992.

- Xu, Z., Zhou, G., & Shimizu, H. (2010). Plant responses to drought and rewatering. *Plant Signaling and Behavior*, 5(6), 649–654.
- Yang, K., Jiang, Y., Wang, J., Cai, X., Wen, Z., Qiu, Z., & Qiao, G. (2022). Tobacco straw biochar improved the growth of Chinese cherry (*Prunus pseudocerasus*) via altering plant physiology and shifting the rhizosphere bacterial community. *Scientia Horticulturae*, 303(111244), 1–13.
- Yildirim, E., Ekinci, M., & Turan, M. (2021). Impact of biochar in mitigating the negative effect of drought stress on cabbage seedlings. *Journal of Soil Science and Plant Nutrition*, 21(3), 2297–2309.
- Yoo, G., Kim, H., Chen, J., & Kim, Y. (2014). Effects of biochar addition on nitrogen leaching and soil structure following fertilizer application to rice paddy soil. *Soil Science Society of America Journal*, 78(3), 852–860.
- Yu-Xue, L., Wei, L., Wei-Xiang, W., Zhe-Ke, Z., & Ying-Xu, C. (2009). Environmental behavior and effect of biomass-derived black carbon in soil: A review. *Chinese Journal of Applied Ecology*, 20(4), 977–982.
- Yuan, J.-H., Xu, R.-K., & Zhang, H. (2011). The forms of alkalis in the biochar produced from crop residues at different temperatures. *Bioresource Technology*, 102, 3488–3497.
- Zaghdoud, C., Alcaraz-López, C., Mota-Cadenas, C., Martnez-Ballesta, M. D. C., Moreno, D. A., Ferchichi, A., & Carvajal, M. (2012). Differential responses of two broccolis (*Brassica oleracea* L. var *italica*) cultivars to salinity and nutritional quality improvement. *The Scientific World Journal*, 2012, 1–12.
- Zhang, X., Wang, D., Jiang, C. C., & Peng, S. A. (2013). Research progress of amelioration effects of biochars on acid soils. *Hubei Agric Sci*, 52, 997–1000.
- Zhang, Xuekun, Lu, G., Long, W., Zou, X., Li, F., & Nishio, T. (2014). Recent progress in drought and salt tolerance studies in *Brassica* crops. *Breeding Science*, 64, 60–73.
- Zhang, Y., Li, Y., Hassan, M. J., Li, Z., & Peng, Y. (2020). Indole-3-acetic acid improves drought tolerance of white clover via activating auxin, abscisic acid and jasmonic acid related genes and inhibiting senescence genes. *BMC Plant Biology*, 20(150), 1–12.
- Zwack, P. J., & Rashotte, A. M. (2015). Interactions between cytokinin signalling and abiotic stress responses. *Journal of Experimental Botany*, 66(16), 4863–4871.