

DAFTAR PUSTAKA

- Aguera, A. & Brophy, D., 2011. Use of sagittal otolith shape analysis to discriminate Northeast Atlantic and Western Mediterranean stocks of Atlantic Saury, *Scomberesox saurus saurus* (Walbaum). Fisheries research, 110(3): 465-471.
- Asia, Zaini, M. & Ngatimin, S. N. A., 2018. Mengenal ikan cakalang dan potensinya sebagai komoditi andalan di Indonesia. Yogyakarta: Leutikaprio.
- Avigliano, E., Domanico, A., Sanchez, S. & Volpedo, A. V., 2017. Otolith elemental fingerprint and scale and otolith morphometry in *Prochilodus lineatus* provide identification of natal nurseries. Fisheries research, 186(1): 1-10.
- Bani, A., Poursaeid, S. & Tuset, V. M., 2013. comparative morphology of the sagittal otolith in three species of South Caspian Gobies. Journal of fish biology, 82: 1321-1332.
- Baweleng, S., Manginsela, F. B. & Sangari, J. R. R., 2018. Studi otolith ikan layang, *Decapterus akaadsi* Abe 1958 dari Perairan Teluk Amurang. Ilmiah platax, 6(2): 66-76.
- Campana, S. E. & Casselman, J. M., 1993. Stock discrimination using otolith shape-analysis. Canadian journal of fisheries and aquatic sciences, 50: 1062-1083.
- Campana, S. E. & Neilson, J. D., 1985. Microstructure of fish otoliths. Canadian Journal of fisheries and aquatic sciences, 42: 1014-1032.
- Cardinale, M., Doering-Arjes, P., Kastowsky, M. & Mosegaard, H., 2004. Effect of sex, stock, and environment on the shape of known-age atlantic cod (*Gadus morhua*) otoliths. Canadian journal of fisheries and aquatic science, 61: 158-167.
- Faizah, R. & Sadiyah, L., 2020. Some biology aspects of indian scad (*Decapterus ruselli* Rupell, 1928) in Pemangkat Fisheries Port West Kalimantan. IOP conference series, 429: 1-10.
- Global Biodiversity Information Facility, 2023. *Katsuwonus pelamis* (Linnaeus, 1758). <<https://www.gbif.org/species/2374191>>. Diakses pada 28 Juni 2023.
- Kementerian Kelautan dan Perikanan, 2022. Rilis data kelautan dan perikanan triwulan i tahun 2022, Jakarta: Pusat Data, Statistik dan Informasi Sekretariat Jenderal Kementerian Kelautan dan Perikanan.

- Kusmini, I. I., Radona, D. & Putri, F. P., 2018. Pola pertumbuhan dan faktor kondisi benih ikan tengadak (*Barbonymus schwanenfeldii*) pada wadah pemeliharaan yang berbeda. LIMNOTEK perairan darat tropis di Indonesia, 25(1): 1-9.
- Mahe, K., Ider D, Massaro A, Hamed O, Jurado-Ruzafa A, Goncalves P, Anastaspoulou A, Jadaud A, Mytilineou C, Elleboode R, Ramdane Z, Bacha M, Amara R, de Pontual H, & Ernande B., 2018. directional bilateral asymmetry in otolith morphology may affect fish stock discrimination based on otolith shape analysis. ICES Journal of marine science, 76(1): 232-243.
- Manginsela, F. B., Mamuaya, G. E., Rompas, R. M. & Lumingas, L. J. L., 2020. The size and the shape of sagittal otolith of redbell scud, *Decapterus kurroides* Bleeker 1855 from Kema Bay, North Minahasa Regency, North Sulawesi, Indonesia. Omni-akuatika, 16(3): 99-110.
- Manginsela, F. B., Rompas, R. M., Mamuaya, G. E. & Lumingas, L. J. L., 2020. Otolith size and shape indeks of mackerel scud *Decapterus macarellus* (Cuvier, 1833) from Manado Bay and Kema Bay, North Sulawesi, Indonesia. AACI bioflux, 13(3): 1723-1734.
- Matsumoto, W. M., Skillman, R. A. & Dizon, A. E., 1984. Synopsis of biological data on skipjack tuna *Katsuwonus pelamis*. NOAA Technical Report NMFS Circular 451. U.S. Department of Commerce.
- Mogea, D., Lumingas, L. J. L. & Mamuaya, G. E., 2019. Analisis biometri otolith (sagitta) untuk pemisahan stok ikan cakalang *Katsuwonus pelamis* (Linnaeus, 1758) yang didaratkan di Pelabuhan Perikanan Pantai Tumumpa (Manado), Sulawesi Utara. Jurnal ilmiah platax, 7(2): 401-412.
- Moore, B. R., Parker, S. J. & Pinkerton, M. H., 2022. Otolith shape as a tool for species identification of the grenadiers *Macrourus caml* and *M. whitsoni*. Fisheries research, 253: 1-11.
- Mourniaty, A. Z. A., Jabbar, M. A., Suyasa, I. N. & Wujdi, A., 2020. Hubungan morfometrik otolith dengan ukuran ikan layang deles (*Decapterus macrosoma* Bleeker, 1851) di Perairan Bali Selatan. BAWAL, 12(3): 103-107.
- Nursan, M., Y. & Agus, S. B., 2022. Distribusi daerah penangkapan ikan cakalang (*katsuwonus pelamis*) menggunakan alat tangkap purse seine di WPP 573. Jurnal pengelolaan perikanan tropis , 06(01): 8-16.
- Ozpacak, M., Saygin, S. & Polat, N., 2019. Otolith shape analysis of bluefish, *Pomatomus saltatrix* (Linnaeus, 1766) in the Black Sea Region (Samsun, Turkey). Acta aquatica turcica, 15(4): 507-516.

- Ponton, D., 2006. Is geometric morphometrics efficient for comparing otolith shape of different fish species?. *Journal of morphology*, 267: 750-757.
- Sadighzadeh, Z., Tuset, V. M., Valinassab, T. Dadpour, M. R & Lombarte, A., 2012. Comparison of different otolith shape descriptors and morphometrics for the identification of closely related species of *Lutjanus* spp. from The Persian Gulf. *Marine biology research*, 8(9): 802-814.
- Salam, A., 2017. Keberlanjutan perikanan tangkap ikan cakalang (*Katsuwonus pelamis*). Yogyakarta: Zahir Publishing.
- Secor, D. H., John, M. D. & Elisabeth, H. L., 1992. Otolith removal and preparation for microchemical examination. *Canadian special publication fisheries aquatic science*, 19-57.
- Simbolon, D., 2011. Bioekologi dan dinamika daerah penangkapan ikan. Bogor: IPB.
- Sinaga, M., Eddlwani, Windarti & Aslah, N., 2021. Growth Circle patterns in the otolith of the gray fish (*Ostochilus melanopleurus*) From The Siak. *Asian journal of akuatik sciences*, 4(2): 144-153.
- Suharti, S. R., 2002. Menentukan umur ikan melalui mikrostruktur otolith. *Oseana*, XXVII: 1-8.
- Syahailatua, A., 2003. Lebih jauh tentang otolith. *Oseana*, XXVIII(1): 7-18.
- Taliawo, R., Manginsela, F. B. & Bataragoa, N. E., 2018. Morfometrik otolith ikan selar (*Selar crumenophthalmus*) dari Teluk Kema. *Jurnal ilmiah platax*, 6(1): 100-106.
- Tanabe, T., Kayama, S., Ogura, M. & Tanaka, S., 2003. Daily increment formation in otoliths of juvenile skipjack tuna *Katsuwonus pelamis*. *Fisheries science*, 69: 731-737.
- Tuli, M., 2018. Sumbedaya ikan cakalang. Gorontalo: Ideas Publishing.
- Viera, A. R., Neves, A., Paira, R. B., Gordo, L. S., 2014. Otolith shape analysis as a tool for stock discrimination of forkbeard (*Phycis phycis*) in the Northeast Atlantic. *Hydrobiologia*, 728(1): 103-110.
- Wujdi, A., Agustina, M. & Jatmiko, I., 2018. Indeks bentuk otolith ikan cakalang, *Katsuwonus pelamis* (Linnaeus, 1758) dari Samudera Hindia. *Jurnal Iktiologi Indonesia*, 18(2): 151-163.
- Wujdi, A., Prihatiningsih & Suwarso, 2016. Karakteristik morfologi dan hubungan morfometrik otolith dengan ukuran ikan lemuru (*Sardinella lemuru* Bleeker, 1853) di Selat Bali. *BAWAL*, 8(3): 159-172.

- Wujdi, A., Setyadji, B. & Nugroho, S. C., 2017. Identifikasi struktur stok ikan cakalang (*Katsuwonus pelamis* Linnaeus, 1758) di Samudera Hindia (WPP NRI 573) Menggunakan Analisis Bentuk Otolith. Jurnal penelitian perikanan indonesia, 23(2): 77-88.
- Yedier, S., 2021. *Otolith* shape analysis and relationship between total length and otolith dimensiens of european barracuda *Sphyraena sphyraena* in the Mediterranean Sea. Iranian journal of fisheries sciences, 20(4): 1080-1096.
- Zengin, M., Saygin, S. & Polat, N., 2015. Otolith shape analyses and dimensions od the anchovy *Engraulis encrasicolus* L. in the Black and Marmara Seas. Sains malaysia, 44(5): 657-662.
- Zengin, M., Saygin, S. & Polat, N., 2017. Relationship between otolith size and total length of bluefish, *Pomatomus saltatrix* (Linnaeus,1766), in Black Sea (Turkey). North-Western journal of zoology, 13(1): 169-171.
- Zischke, M. T., Litherland, L., Tilyard B. R., Stratford, N. J., Jones, E. L., Wang, You-Gan., 2016. Otolith morphology of four mackerel species (*Scombemorus* spp.) in Australia: Species Differentiation and Prediction for Fisheries Monitoring and Assessment. Fisheries research, 176: 39-47.