

DAFTAR PUSTAKA

- Agustin, V. 2022. Aktivitas antioksidan hidrolisat protein dari hasil samping ikan gabus (*Channa striata*) yang diproses secara enzimatis. Fakultas Pertanian. Universitas Gadjah Mada. Tesis.
- Agustina, S., Aidha, N. N., & Oktarina, E. 2018. Ekstraksi antioksidan *spirulina sp.* dengan menggunakan metode ultrasonikasi dan aplikasinya untuk krim kosmetik. *Jurnal Kimia dan Kemasan*. 40(2): 105-16.
- Aidah, S. N. 2020. Panduan praktis usaha budidaya belut bagi pemula. Penerbit KBM Indonesia. Bojonegoro.
- Alemán, A., Pérez-Santín, E., Bordenave-Juchereau, S., Arnaudin, I., Gómez-Guillén M.C. & Montero, P. 2011. Squid gelatin hydrolysates with antihypertensive, anticancer and antioxidant activity. *Food Res Int* 44:1044–1051. doi: 10.1007/s13197-014-1581-6.
- Al Umam, Richard. 2023. Pengaruh konsentrasi papain pada pembuatan hidrolisat protein insang dan arborescent lele terhadap aktivitas antibakterinya pada bakteri pembentuk histamin. Fakultas Pertanian. Universitas Gadjah Mada. Skripsi.
- Andarwulan, N., Wijaya, C. H., & Cahyono, D. T. 1996. Aktivitas antioksidan dari daun sirih (*Piper betle L.*). *Teknologi dan Industri Pangan*. 7(1): 29-30.
- Annisa, S., Darmanto, Y. S., & Amalia, U. 2017. Pengaruh perbedaan spesies ikan terhadap hidrolisat protein ikan dengan penambahan enzim papain (*The effect of various fish species on fish protein hydrolysate with the addition of papain enzyme*). *Indonesian Journal of Fisheries Science and Technology*. 13(1): 24-30.
- AOAC. 2005. Analysis of the association of official analytical chemist. Official Methods of Analysis of the Association of Official Analytical Chemist 16th Ed. Washington DC.
- Aspevik, T., Totland, C., Lea, P., & Oterhals, Å. 2016. Sensory and surface-active properties of protein hydrolysates based on atlantic salmon (*Salmo salar*) by-products. *Process Biochemistry*. 51(8): 1006-1014.
- Aspmo SI, Horn SJ & Eijsink VGH. 2005. Use of hydrolysates from atlantic cod (*gadus morhua l.*) viscera as complex nitrogen source for lactic acid bacteria. *FEMS Microbiol Lett*. 248 : 65-68.
- Astiana, I. 2012. Perubahan komposisi asam amino dan mineral belut sawah (*monopterus albus*) akibat proses penggorengan. Skripsi. Fakultas Perikanan dan Ilmu Kelautan. IPB. Bogor.

- Astiana, I. 2012. Perubahan komposisi asam amino dan mineral belut sawah (*monopterus albus*) akibat proses penggorengan. *Jurnal Pengolahan Hasil Perikanan Indonesia*. 18(3): 230-239.
- Awwaly, K. U. A. 2017. *protein pangan hasil ternak dan aplikasinya*. UB Press. Malang.
- Baehaki, A., Lestari, S. D., & Romadhoni, A. R. 2015. Hidrolisis protein ikan patin menggunakan enzim papain dan aktivitas antioksidan hidrolisatnya. *Jurnal Pengolahan Hasil Perikanan Indonesia*. 18(3): 230-239.
- Balti, R., Jridi, M., Sila, A., Souissi, N., Nedjar-Arroume, N., Guillochon, D., & Nasri, M. 2011. extraction and functional properties of gelatin from the skin of cuttlefish (*Sepia officinalis*) using smooth hound crude acid protease-aided process. *Food Hydrocolloids*. 25(5): 943-950.
- Barzana, E., and Garcia-Garibay, N., 1994. Production of Fish protein concentrate, in martin, a.m. (ed.) *fisheries processing, biotechnology applications*. London Chapman & Hall. 221pp.
- Bernadeta, P. A. 2012. Penentuan kondisi optimum hidrolisat protein dari limbah ikan ekor kuning (*Caesio cuning*) berdasarkan karakteristik organoleptik. *jurnal kimia khatulistiwa*. 1(1).
- Bhat, M. S., Ray, S., & Datta, P. M. 2018. A new assemblage of freshwater sharks (*Chondrichthyes: Elasmobranchii*) from the upper triassic of India. *Geobios*. 51(4): 269-283.
- Binsan, W., Benjakul, S., Visessanguan, W., Roytrakul, S., Tanaka, M., & Kishimura, H. 2008. Antioxidative activity of mungoong, an extract paste from the cephalothorax of white shrimp (*Litopenaeus Vannamei*). *Food Chemistry*. 106(1): 185-193.
- Boligon, A. A., Machado, M. M., & Athayde, M. L. 2014. Technical evaluation of antioxidant activity. *Med. Chem*. 4(7): 517-522.
- Cahyaningati, K. 2019. Aktivitas antioksidan hasil hidrolisis enzimatis protein ikan wader (*Rasbora jacobsoni*) menggunakan kombinasi calotropin dan papain. *Teknologi Hasil Pertanian Fakultas Teknologi Pertanian Universitas Jember*. Tesis.
- Chandrasekara, A., & Shahidi, F. 2011. Determination of antioxidant activity in free and hydrolyzed fractions of millet grains and characterization of their phenolic profiles by HPLC-DAD-ESI-MSn. *Journal of Functional Foods*. 3(3): 144-158.
- Chandrasekara, N., & Shahidi, F. 2011. Effect of roasting on phenolic content and antioxidant activities of whole cashew nuts, kernels, and testa. *Journal of Agricultural and Food Chemistry*. 59(9): 5006-5014.
- Cheng, M., Qin, Z., Hu, J., Liu, Q., Wei, T., Li, W., & Liu, B. 2020. Facile and rapid one-step extraction of carboxylated cellulose nanocrystals by H₂SO₄/HNO₃ mixed acid hydrolysis. *Carbohydrate Polymers*. 231: 115701.

- Cheung, R. C. F., Ng, T. B., & Wong, J. H. 2015. Marine peptidas: bioactivities and applications. *Marine drugs* 13(7): 4006-4043.
- Elavarasan, K., dan B. A. Shamasundar. 2015. Effect of oven drying and freeze drying on the antioxidant and functional properties of protein hydrolysates derived from freshwater fish (*Cirrhinus mrigala*) using papain enzyme. *Journal of Food Science and Technology*. 53(2): 1303–1311.
- Elsson, M., Wijanarko, A., Hermansyah, H., & Sahlan, M. 2019. Michaelis-menten parameters characterization of commercial papain enzyme “paya”. IOP Conference Series: Earth and Environmental Science. 217(1): 012037. IOP Publishing.
- Farzaneh, V., Gominho, J., Pereira, H., & Carvalho, I. S. 2018. Screening of the antioxidant and enzyme inhibition potentials of portuguese *Pimpinella anisum L.* seeds by GC-MS. *Food Analytical Methods*. 11: 2645-2656.
- Gbogouri, G. A., Linder, M., Fanni, J., & Parmentier, M. 2004. Influence of hydrolysis degree on the functional properties of salmon byproducts hydrolysates. *Journal of Food Science*. 69(8): C615-C622.
- Guan, H., Diao, X., Jiang, F., Han, J., & Kong, B. 2018. The enzymatic hydrolysis of soy protein isolate by *Corolase PP* under high hydrostatic pressure and its effect on bioactivity and characteristics of hydrolysates. *Food Chemistry*. 245: 89-96.
- Hafifani, W., Puspita, I. D., & Putra, M. M. 2021. The effect of hydrolysis duration on the antibacterial activity of swamp eel head protein hydrolysate produced by papain against histamine-producing bacteria. In *E3S Web of Conferences*. 322: 04005.
- Harnedy, P. A., Parthasarathy, V., McLaughlin, C. M., O’Keeffe, M. B., Allsopp, P. J., McSorley, E. M., & FitzGerald, R. J. 2018. Atlantic salmon (*Salmo salar*) co-product-derived protein hydrolysates: a source of antidiabetic peptidas. *Food Research International*. 106: 598-606.
- Hidayati, A., & Santoso, J. 2019. Aktivitas antioksidan hidrolisat protein miofibril belut (*Synbranchus bengalensis*) yang dihidrolisis dengan enzim papain. *Jurnal Teknologi Industri Pertanian*. 29(3).
- Hihat, S., Remini, H., & Madani, K. 2017. Effect of oven and microwave drying on phenolic compounds and antioxidant capacity of coriander leaves. *International Food Research Journal*. 24(2).
- Himaya, S. W. A., Ngo, D. H., Ryu, B., & Kim, S. K. 2012. An active peptida purified from gastrointestinal enzyme hydrolysate of pacific cod skin gelatin attenuates angiotensin-1 converting enzyme (ACE) activity and cellular oxidative stress. *Food Chemistry*. 132(4): 1872-1882.
- Hoyle, N. T., & Merritt, J. H. 1994. Quality of fish protein hydrolysates from herring (*Clupea harengus*). *Journal of Food Science*. 59(1): 76-79.

- Hustiany, R. 2005. Karakteristik produk olahan kerupuk dan surimi dari daging ikan patin (*Pangasius sutchi*) hasil budidaya sebagai sumber protein hewani. *Media Gizi dan Keluarga*, 29(2), 66-74.
- Jamil, B., Abbasi, R., Abbasi, S., Imran, M., Khan, S. U., Ihsan, A. & Imran, M. 2016. encapsulation of cardamom essential oil in chitosan nano-composites: in-vitro efficacy on antibiotic-resistant bacterial pathogens and cytotoxicity studies. *Frontiers in Microbiology*. 7: 1580.
- Jellouli, K., Balti, R., Bougatef, A., Hmidet, N., Barkia, A., & Nasri, M. 2011. Chemical composition and characteristics of skin gelatin from grey triggerfish (*Balistes capriscus*). *LWT-Food Science and Technology*. 44(9): 1965-1970.
- Kantun, W. K., Malik, A. A., & Harianti, H. 2015. Feasibility of solid waste tuna loin of yellowfin tuna (*Thunnus albacares*) raw materials for the product diversification. *Jurnal Pengolahan Hasil Perikanan Indonesia*. 18(3).
- Khanh, N., & Ngan, H. 2010. current practices of rice field eel (*Monopterus albus*) culture in Vietnam. *Aquaculture Asia Mag*. 15: 26-29.
- Kiay, N., Suryanto, E., & Mamahit, L. 2011. Efek lama perendaman ekstrak kalamansi (*Citrus microcarpa*) terhadap aktivitas antioksidan tepung pisang goroho (*Musa spp.*). *Chemistry Progress*. 4(1).
- Kielkopf, C. L., Bauer, W., & Urbatsch, I. L. 2020. Bradford assay for determining protein concentration. *Cold Spring Harbor Protocols*. 2020(4): pdb-prot102269.
- Kiewiet, M. B., Faas, M. M., & De Vos, P. 2018. immunomodulatory protein hydrolysates and their application. *Nutrients*. 10(7): 904.
- Klomkloa, S., Benjakul, S., & Kishimura, H. 2013. Functional properties and antioxidative activity of protein hydrolysates from toothed ponyfish muscle treated with viscera extract from hybrid catfish. *International Journal of Food Science & Technology*. 48(7), 1483-1489.
- Kristinsson, H. G., & Rasco, B. A. 2000. Biochemical and functional properties of atlantic salmon (*Salmo salar*) muscle proteins hydrolyzed with various alkaline proteases. *Journal of Agricultural And Food Chemistry*. 48(3): 657-666.
- Kumar, V., & Shamasundar, B. A. 2014. Antioxidant and functional properties of fish protein hydrolysates from fresh water carp (*C Atla Catla*) as influenced by the nature of enzyme. *Journal of Food Processing and Preservation*. 38(3): 1207-1214.
- Kurniawan, R. 2012. Pengaruh konsentrasi larutan garam dan waktu. *Jurnal Teknik Kimia*. 2(2): 127-135.
- Leni, G., Soetemans, L., Caligiani, A., Sforza, S., & Bastiaens, L. 2020. Degree of hydrolysis affects the techno-functional properties of lesser mealworm protein hydrolysates. *Foods*. 9(4): 381.

- Leo, E. E. M., Fernández, J. J. A., & Campos, M. R. S. 2016. Biopeptidas with antioxidant and anti-inflammatory potential in the prevention and treatment of diabetes disease. *Biomedicine & Pharmacotherapy*. 83: 816-826.
- Lestari, P. B., & Hartati, T. W. 2017. *Mikrobiologi berbasis inkuiry*. Penerbit Gunung Samudera.
- Mamboya, E. A. F., & Amri, E. 2012. Papain, a plant enzyme of biological importance: a review. *Am. J. Biochem. Biotechnol.* 8(2): 99-104.
- Mirzaei, M., Mirdamadi, S., Ehsani, M. R., & Aminlari, M. 2018. Production of antioxidant and ACE-inhibitory peptidas from *Kluyveromyces marxianus* protein hydrolysates: purification and molecular docking. *Journal of Food and Drug Analysis*. 26(2): 696-705.
- Mudgil, P., Kamal, H., Yuen, G. C., & Maqsood, S. 2018. Characterization and identification of novel antidiabetic and anti-obesity peptidas from camel milk protein hydrolysates. *Food Chemistry*. 259: 46-54.
- Mutamimah, D., Ibrahim, B., & Trilaksani, W. 2018. Antioxidant activity of protein hydrolysate produced from tuna eye (*Thunnus Sp.*) by enzymatic hydrolysis. *Jurnal Pengolahan Hasil Perikanan Indonesia*. 21(3): 522-531.
- Najafian, L., & Babji, A. S. 2014. Production of bioactive peptidas using enzymatic hydrolysis and identification antioxidative peptidas from patin (*Pangasius sutchi*) sarcoplasmic protein hydrolysate. *Journal of Functional Foods*. 9: 280-289.
- Najafian, L., & Babji, A. S. 2015. Isolation, purification and identification of three novel antioxidative peptidas from patin (*Pangasius sutchi*) myofibrillar protein hydrolysates. *LWT-Food Science and Technolog.* 60(1): 452-461.
- Naqash, S. Y., & Nazeer, R. A. 2011. Evaluation of bioactive properties of peptida isolated from exocoetus volitans backbone. *International Journal of Food Science & Technology*. 46(1): 37-43.
- Noman, A., Qixing, J., Xu, Y., Ali, A. H., Al-Bukhaiti, W. Q., Abed, S. M., & Xia, W. 2019. Influence of degree of hydrolysis on chemical composition, functional properties, and antioxidant activities of chinese sturgeon (*Acipenser sinensis*) hydrolysates obtained by using alcalase 2.4 l. *Journal of Aquatic Food Product Technology*. 28(6): 583-597.
- Normah, I., & Noorasma, M. 2015. Physicochemical properties of mud clam (*Polymesoda erosa*) hydrolysates obtained using different microbial enzymes. *International Food Research Journal*. 22(3).
- Ochiai, A., Ishii, G., & Neri, S. 2016. Phenotypic and functional heterogeneity of cancer-associated fibroblast within the tumor microenvironment. *Advanced Drug Delivery Reviews*. 99: 186-196.

- Ovando, C. A., Carvalho, J. C. D., Vinícius de Melo Pereira, G., Jacques, P., Soccol, V. T., & Soccol, C. R. 2018. Functional properties and health benefits of bioactive peptidas derived from spirulina: a review. *Food Reviews International*. 34(1): 34-51.
- Ovissipour, M., Abedian, A., Motamedzadegan, A., Rasco, B., Safari, R., & Shahiri, H. 2009. The effect of enzymatic hydrolysis time and temperature on the properties of protein hydrolysates from persian sturgeon (*Acipenser persicus*) viscera. *Food Chemistry*. 115(1): 238-242.
- Pramono, P. K. P. 2022. Rasio efisiensi protein ransum broiler yang disuplementasi mannan oligosakarida (MOS) hasil hidrolisis bungkil inti sawit dengan enzim mannanase. Universitas Jambi. Tesis.
- Prastika, H. H., Ratnayani, K., Puspawati, N. M., & Laksmiwati, A. A. I. A. M. 2019. Penggunaan enzim pepsin untuk produksi hidrolisat protein kacang gude (*Cajanus cajan* (L.) Millsp.) yang aktif antioksidan. *Cakra Kimia*. 7(2), 180-188.
- Priatni, S., Harimadi, Kezia Buana, Efendi Kosasih, W., & Rohmatussolihat, R. 2020. Production and characterization of spray-dried swamp eel (*Monopterus albus*) protein hydrolysate prepared by papain. *Sains Malaysiana*. 49(3): 545-552
- Prihanto, A. A., Nurdiani, R., & Sari, L. W. 2021. Production and characteristics of sailfin catfish (*Pterygoplichthys pardalis*) protein hydrolysate. *F1000 Research*. 10: 1089.
- Purbasari, D. 2008. Produksi dan karakterisasi hidrolisat protein dari kerang mas ngur (*Atactodea striata*). *Marine Niche*. 283-297.
- Puspawati, N.M., P.P. Dewi., N.W. Bogoriani., dan N.K. Ariati. 2020. Produksi hidrolisat protein antioksidan melalui hidrolisis enzimatis protein kulit ayam broiler dengan enzim papain. *Jurnal Kimia*. 14(2): 206-212.
- Qiao, M., Ying, G. G., Singer, A. C., & Zhu, Y. G. 2018. Review of antibiotic resistance in china and its environment. *Environment International*. 110: 160-172.
- Radhiyah., Windarti., Ridwan M. P. 2021. Pola pertumbuhan dan kebiasaan makan belut sawah (*Monopterus albus*) dari perairan umum stadion utama Riau. *Jurnal Ilmu Perairan*. 9(1), 68-76.
- Ranathunga, S., Rajapakse, N., & Kim, S. K. 2006. Purification and characterization of antioxidative peptida derived from muscle of conger eel (*Conger myriaster*). *European Food Research and Technology*. 222: 310-315.
- Roslan, J., Yunos, K. F. M., Abdullah, N., & Kamal, S. M. M. 2014. Characterization of fish protein hydrolysate from tilapia (*Oreochromis niloticus*) by-product. *Agriculture and Agricultural Science Procedia*. 2: 312-319.
- Saanin, H. 1984. Taksonomi dan kunci identifikasi ikan. Bina Cipta. Jakarta.

- Santi, Y. M. 2009. Analisis usaha agroindustri keripik belut sawah (*Monopterus albus zuieuw*) di Kabupaten Klaten. Universitas Sebelas Maret. Skripsi.
- Sari, R. K. 2019. Pengaruh konsentrasi asam klorida (HCl) dan waktu hidrolisis terhadap kandungan glukosa dan kadar bioetanol dari limbah kulit kakao (*Theobroma cacao L.*). Universitas Muhammadiyah Surakarta.
- Sarmadi & Ismail, 2010 Sarmadi, B. H., & Ismail, A. 2010. Antioxidative peptidas from food proteins: a review. *Peptidas*. 31(10): 1949-1956.
- Sarwono. 2003. Budidaya belut dan sidat. Penebar Swadaya. Jakarta.
- Setiawan, F., Yunita, O., & Kurniawan, A. 2018. Uji aktivitas antioksidan ekstrak etanol kayu secang (*Caesalpinia sappan*) menggunakan metode DPPH, ABTS, dan FRAP. *Media Pharmaceutica Indonesiana*. 2(2): 82-89.
- Shahidi, F., Han, X. Q., & Synowiecki, J. 1995. Production and characteristics of protein hydrolysates from capelin (*Mallotus villosus*). *Food chemistry*. 53(3): 285-293.
- Silaban, N. A. 2022. Pengaruh lama hidrolisis dengan papain terhadap aktivitas antibakteri hidrolisat protein belut pada bakteri pembentuk histamin. Universitas Gadjah Mada. Skripsi.
- Silvestre, M. P., da Silva, M. C., de Souza, M. W., Silva, V. D., de Aguiar, M. J., & Silva, M. R. 2013. Hydrolysis degree, peptida profile and phenylalanine removal from whey protein concentrate hydrolysates obtained by various proteases. *International Journal of Food Science & Technology*. 48(3): 588-595.
- Simanjorang, E., Kurniawati, N., & Hasan, Z. 2012. Pengaruh penggunaan enzim papain dengan konsentrasi yang berbeda terhadap karakteristik kimia kecap tutut. *Jurnal Perikanan Kelautan*. 3(4).
- Šližytė, R., Rustad, T., & Storrø, I. 2005. Enzymatic hydrolysis of cod (*Gadus morhua*) by-products: Optimization of Yield and Properties of Lipid and Protein Fractions. *Process Biochemistry*. 40(12): 3680-3692.
- Sunarlim, R. O. S. W. I. T. A., & Usmiati, S. 2009. Karakteristik daging kambing dengan perendaman enzim papain. *Teknologi Peternakan dan Veteriner*.
- Supratyami, M. 2016. Pemanfaatan belut (*Monopterus albus*) pada pembuatan crackers ber-omega-3. *Jurnal Agroindustri Halal*. 2(2): 081-089.
- Susanto, E., Rosyidi, D., Radiati, L. E., & Subandi, S. 2018. Optimasi aktivitas antioksidan peptida aktif dari cecek ayam melalui hidrolisis enzim papain. *Jurnal Ilmu dan Teknologi Hasil Ternak (JITEK)*. 13(1): 14-26.
- Susanto, E., Rosyidi, D., Radiati, L. E., & Subandi, S. 2018. Optimasi aktivitas antioksidan peptida aktif dari cecek ayam melalui hidrolisis enzim papain. *Jurnal Ilmu Dan Teknologi Hasil Ternak (JITEK)*. 13(1). 14-26.

- Susanty, A dan Indrati K. 2021. Pengaruh waktu hidrolisis terhadap karakteristik hidrolisat protein ikan toman (*Channa micropeltes*) asal das Kalimantan Timur. *JRTI*. 15(2): 463-475.
- Susi. 2012. Komposisi kimia dan asam amino pada tempe kacang nagara (*Vigna unguiculata* ssp. *Cylindrica*). *Jurnal Agroscientiae*. 19(1): 28-36.
- Taniyo, W., Salimi, Y. K., & Iyabu, H. 2021. Karakteristik dan aktivitas antioksidan hidrolisat protein ikan nike (*Awaous melanocephalus*). *Dalton: Jurnal Pendidikan Kimia dan Ilmu Kimia*. 4(2): 52-63.
- Theafelicia, Z., & Wulan, S. N. 2023. Perbandingan berbagai metode pengujian aktivitas antioksidan (DPPH, ABTS dan FRAP) pada teh hitam (*Camellia sinensis*). *Jurnal Teknologi Pertanian*. 24(1): 35-44.
- Utomo, B. S. B., Suryanigrum, T. D., & Harianto, H. R. 2014. Optimization of enzymatic hydrolysis of fish protein hydrolysate (FPH) processing from waste of catfish fillet production. *Journal of Squalen Bulletin of Marine and Fisheries Postharvest and Biotechnology*. 9(3): 115-126.
- Venugopal, V. 2016. Enzymes from seafood processing waste and their applications in seafood processing. *Advances in Food and Nutrition Research*. 78: 47-69.
- Vijaykrishnaraj, M., Roopa, B. S., & Prabhasankar, P. 2016. Preparation of gluten free bread enriched with green mussel (*Perna canaliculus*) protein hydrolysates and characterization of peptidas responsible for mussel flavour. *Food Chemistry*. 211: 715-725.
- Vilcacundo, R., Barrio, D. A., Piñuel, L., Boeri, P., Tombari, A., Pinto, A., & Carrillo, W. 2018. Inhibition of lipid peroxidation of kiwicha (*Amaranthus caudatus*) hydrolyzed protein using zebrafish larvae and embryos. *Plants*. 7(3): 69.
- Vilcacundo, R., Méndez, P., Reyes, W., Romero, H., Pinto, A., & Carrillo, W. 2018. Antibacterial activity of hen egg white lysozyme denatured by thermal and chemical treatments. *Scientia Pharmaceutica*. 86(4): 48.
- Wijayanti, I., & Setiyorini, E. 2018. Nutritional content of wild and cultured eel (*Anguilla bicolor*) from southern coast of central java. *Indonesian Journal of Marine Sciences*. 23(1): 37.
- Wijayanti, I., Romadhon dan Laras R. 2016. Karakteristik hidrolisat protein ikan bandeng (*Chanos chanos Forsk*) dengan konsentrasi enzim bromelin yang berbeda. *Indonesian Journal Fisheries Science and Technology*. 11(2): 129-133.
- Winarsi, H. 2007. *Antioksidan alami dan radikal*. Kanisius.
- Witono, Y., Iwan T., Wiwik S. W., Amelia R. 2014. Pengaruh waktu hidrolisis terhadap karakteristik hidrolisat protein ikan toman (*Channa micropeltes*) asal das Kalimantan Timur. *Jurnal Teknologi dan Industri Pangan*. 25(2): 140-145

- Witono, Y., Windrati, W. S., Taruna, I., Afriliana, A., & Assadam, A. 2014. characteristics and sensory analysis of ketchup and sauce products from “Bibian” fish hydrolysate. *American Journal of Food Science and Technology*. 2(6): 203-208.
- Wu, H. C., Chen, H. M., & Shiau, C. Y. 2003. Free amino acids and peptidas as related to antioxidant properties in protein hydrolysates of mackerel (*Scomber austriasicus*). *Food Research International*. 36(9-10): 949-957.
- Yoga, I. K. W. (2015). Penentuan konsentrasi optimum kurva standar antioksidan; asam galat, asam askorbat dan trolox® terhadap radikal bebas DPPH (2, 2-diphenyl-1-picrylhydrazyl) 0, 1 mM. *Prosiding Seminar Nasional MIPA*.
- Yuniarti T., Shanti D. L., Medal L. P Y. P.H., Heny B. P., Sonny K., Sri N. A., Nuberty T., Sherly R., Rufnia A. A., Adham P., Mirna Z. T. 2021. Pengetahuan bahan baku perikanan. *Yayasan Kita Menulis*.
- Yuniarti, T., Prayudi, A., Supenti, L., Suhrawardan, H., & Martosuyono, P. 2021. Produksi dan profil kimia hidrolisat protein dari hasil samping pengolahan udang segar. *Jurnal Perikanan Universitas Gadjah Mada*. 23(1): 63-69.
- Zahrotul Khoiriah, Z. E. S. I., Herpandi, H., & Dwita Lestari, S. 2019. Pemanfaatan limbah kepala belut (*Monopterus albus*) sebagai bahan campuran kopi (*Coffea arabica*). *Universitas Sriwijaya*. Skripsi.
- Zailanie, K. 2015. *Fish handling*. Universitas Brawijaya Press.