

DAFTAR PUSTAKA

- Agarwal, G., Wilson, J.R., Kurina, S.J., Anaya-Eugenio, G.D., Ninh, T.N., Burdette, J.E., Soejarto, D.D., Cheng, X., Carcache de Blanco, E.J., Rakotondraibe, L.H., and Kinghorn, A.D., 2019, Structurally Modified Cyclopenta[b]benzofuran Analogues Isolated from *Aglaia perviridis*, *J. Nat. Prod.*, 82, 2870-2877.
- Ahmadu, A., Tarimaledei, P., and Onanuga, A., 2013, Triterpenoids from *Gutenbergia Nigritana*(Benth). Oliv and Hiern, *Afr. J. Tradit. Complement. Altern. Med.*, 10, 405-409.
- Altemimi, A., Lakhssassi, N., Baharlouei, A., Watson, D.G., and Lightfoot, D.A., 2017, Phytochemicals: Extraction, Isolation, and Identification of Bioactive Compounds from Plant Extracts., *Plants*, 6, 42-65.
- Anthonsen, T., Bruun, T., Hemmer, E., Holme, D., Lamvik, A., Sunde, E., and Sørensen, N.A., 1970, Baccharis Oxide, a new Triterpenoid from *Baccharis halimifolia* L., *Acta Chem. Scand.*, 24, 2479-2488.
- Awang, K., Loong, X.M., Leong, K.H., Supratman, U., Litaudon, M., Mukhtar, M.R., and Mohamad, K., 2012, Triterpenes and steroids from the leaves of *Aglaia exima* (Meliaceae), *Fitoterapia*, 83, 1391-1395.
- Bondet, V., Brand-Williams, W., and Berset, C., 1997, Kinetics and mechanisms of antioxidant activity using the DPPH. free radical method, *Leb. + Technol.*, 30, 609-615.
- Cai, C., Ma, J., Han, C., Jin, Y., Zhao, G., and He, X., 2019, Extraction and antioxidant activity of total triterpenoids in the mycelium of a medicinal fungus, *Sanghuangporus sanghuang*, *Sci. Rep.*, 9, 7418-7427.
- Chen, B., Zhang, J., Han, J., Zhao, R., Bao, L., Huang, Y., and Liu, H., 2019, Lanostane Triterpenoids with Glucose-Uptake-Stimulatory Activity from Peels of the Cultivated Edible Mushroom *Wolfiporia cocos*, *J. Agric. Food Chem.*, 67, 7348-7364.
- Chudzik, M., Korzonek-Szlacheta, I., and Król, W., 2015, Triterpenes as potentially cytotoxic compounds, *Molecules*, 20, 1610-1625.
- Coskun, O., 2016, Separation Techniques: Chromatography, *North. Clin. Istanbul*, 3, 156-160.
- Cragg, G.M. and Newman, D.J., 2005, Plants as a source of anti-cancer agents., *J. Ethnopharmacol.*, 100, 72-79.
- Cragg, G.M. and Pezzuto, J.M., 2016, Natural Products as a Vital Source for the Discovery of Cancer Chemotherapeutic and Chemopreventive Agents, *Med. Princ. Pract.*, 25, 41-59.
- Dapat, E., Jacinto, S., and Efferth, T., 2013, A phenolic ester from *Aglaia loheri* leaves reveals cytotoxicity towards sensitive and multidrug-resistant cancer cells, *BMC Complement. Altern. Med.*, 13, 286-296.

- Davis, J.M., Navolanic, P.M., Weinstein-Opppenheimer, C.R., Steelman, L.S., Hu, W., Konopleva, M., Blagosklonny, M. V., and McCubrey, J.A., 2003, Raf-1 and Bcl-2 induce distinct and common pathways that contribute to breast cancer drug resistance, *Clin. cancer Res.*, 9, 1161-1170.
- Dias, D.A., Urban, S., and Roessner, U., 2012, A historical overview of natural products in drug discovery, *Metabolites*, 2, 303-336.
- Elyashberg, M. and Williams, A., 2015, M.E. Elyashberg, A.J. Williams, *Computer-based Structure Elucidation from Spectral Data. The Art of Solving Problems*, 1st Ed, Springer, Heidelberg.
- Fabio, G.D., Romanucci, V., De Marco, A., and Zarrelli, A., 2014, Triterpenoids from *Gymnema sylvestre* and Their Pharmacological Activities, *Molecules*, 19, 10956-10981.
- Farabi, K., Harneti, D., Nurlelasari, Maharani, R., Hidayat, A.T., Awang, K., Supratman, U., and Shiono, Y., 2017b, New cytotoxic protolimonoids from the stem bark of *Aglaia argentea* (Meliaceae), *Phytochem. Lett.*, 21, 211-215.
- Farabi, K., Harneti, D., Nurlelasari, Maharani, R., Hidayat, A.T., Supratman, U., Awang, K., and Shiono, Y., 2017, Cytotoxic steroids from the bark of *Aglaia argentea* (Meliaceae), *Chiang Mai Univ. J. Nat. Sci.*, 16, 293-306.
- Ferlay, J., Colombet, M., Soerjomataram, I., Parkin, D.M., Piñeros, M., Znaor, A., and Bray, F., 2021, Cancer statistics for the year 2020: An overview., *Int. J. cancer*, 149, 778-789.
- Fridlender, M., Kapulnik, Y., and Koltai, H., 2015, Plant derived substances with anti-cancer activity: from folklore to practice., *Front. Plant Sci.*, 6, 799.
- Geran, R.I., Greenberg, N.H., MacDonald, M.M., Schumacher, A.M., and Abbott, B., 1972, No Title, *J. Cancer Chemother. Rep.*, 13, 83-91.
- Gogavekar, S.S., Rokade, S.A., Ranveer, R.C., Ghosh, J.S., Kalyani, D.C., and Sahoo, A.K., 2014, Important nutritional constituents, flavour components, antioxidant and antibacterial properties of *Pleurotus sajor-caju*, *J. Food Sci. Technol.*, 51, 1483-1491.
- Greger, H., 2022, Comparative phytochemistry of flavaglines (= rocaglamides), a group of highly bioactive flavolignans from *Aglaia* species (Meliaceae), *Phytochem. Rev.*, 21, 725-764.
- Gurning, K., Haryadi, W., and Sastrohamidjojo, H., 2021, Isolation and characterization of antioxidant compounds of bangun-bangun (*Coleus amboinicus*, L.) leaves from North Sumatera, Indonesia, *Rasayan J. Chem.*, 14, 248-253.
- Hanson, J.R., 2003, *Natural Products: The Secondary Metabolites*, Royal Society of Chemistry, UK.
- Harneti, D. and Supratman, U., 2021, Phytochemistry and biological activities of *Aglaia* species, *Phytochemistry*, 181, 112540-112564.
- Harneti, D., Tjokronegoro, R., Safari, A., Supratman, U., Loong, X.M., Mukhtar,

- M.R., Mohamad, K., Awang, K., and Hayashi, H., 2012, Cytotoxic triterpenoids from the bark of *Aglaia smithii* (Meliaceae), *Phytochem. Lett.*, 5, 496-499.
- Huang, P.-Q., Deng, J.-W., Li, Y., Liao, Z.-B., Zhao, E., Tian, Y.-C., Tu, Y.-D., Li, D.-L., Jin, J.-W., Zhou, C.-X., Wu, R.-H., and Gan, L.-S., 2022, Terpenoids from the twigs and leaves of *Aglaia elaeagnoides* and their chemotaxonomic significance, *Biochem. Syst. Ecol.*, 103, 104427-104432.
- Hutagaol, R.P., Harneti, D., Hidayat, A.T., Nurlelarsari, Maharani, R., Katja, D.G., Supratman, U., Awang, K., and Shiono, Y., 2020, (22e,24s)-24-propylcholest-5en-3 α -acetate: A new steroid from the stem bark *aglaia angustifolia* (miq.) (meliaceae), *Molbank*, 2020, 3-8.
- Hwang, B.Y., Su, B., Chai, H., Mi, Q., Kardono, L.B.S., Afriastini, J.J., Riswan, S., Santarsiero, B.D., Mesecar, A.D., Wild, R., Fairchild, C.R., Vite, G.D., Rose, W.C., Farnsworth, N.R., Cordell, G.A., Pezzuto, J.M., Swanson, S.M., and Kinghorn, A.D., 2004, Derivatives from *Aglaia silvestris*, *J. Org. Chem.*, 69, 3350-3358.
- Joycharat, N., Plodpai, P., Panthong, K., Yingyongnarongkul, B.E., and Voravuthikunchai, S., 2010, Terpenoid constituents and antifungal activity of *Aglaia forbesii* seed against phytopathogens, *Can. J. Chem.*, 88, 937-944.
- Kampa, M., Nistikaki, A., Tsaousis, V., Maliataki, N., Notas, G., and Castanas, E., 2002, A new automated method for the determination of the total antioxidant capacity (TAC) of human plasma, based on the crocin bleaching assay. *BMC Clin. Pathol.*, 2, 19-25.
- Khazir, J., Riley, D.L., Pilcher, L.A., De-Maayer, P., and Mir, B.A., 2014, Anticancer Agents from Diverse Natural Sources, *Nat. Prod. Commun.*, 9, 1655-1669.
- Kumar, P., Mukherjee, M., Johnson, J.P.S., Patel, M., Huey, B., Albertson, D.G., and Simin, K., 2012, Cooperativity of Rb, Brca1, and p53 in malignant breast cancer evolution., *PLoS Genet.*, 8, 27-40.
- Leong, K.H., Looi, C.Y., Loong, X.M., Cheah, F.K., Supratman, U., Litaudon, M., Mustafa, M.R., and Awang, K., 2016, Cycloart-24-ene-26-ol-3-one, a new cycloartane isolated from leaves of *Aglaia exima* triggers tumour necrosis factor-receptor 1-mediated caspase-dependent apoptosis in colon cancer cell line, *PLoS One*, 11, 1-17.
- Li, J., Larregieu, C.A., and Benet, L.Z., 2016, Classification of natural products as sources of drugs according to the biopharmaceutics drug disposition classification system (BDDCS)., *Chin. J. Nat. Med.*, 14, 888-897.
- Ling, T., Boyd, L., and Rivas, F., 2022, Triterpenoids as Reactive Oxygen Species Modulators of Cell Fate, *Chem. Res. Toxicol.*, 35, 569-584.
- Liu, B. and Xu, Y.K., 2016, Cytotoxicity and Synergistic Effect of the Constituents from Roots of *Aglaia odorata* (Meliaceae), *Nat. Prod. Res.*, 30, 433-437.

- Liu, C.-H., Yen, M.-H., Tsang, S.-F., Gan, K.-H., Hsu, H.-Y., and Lin, C.-N., 2010, Antioxidant triterpenoids from the stems of *Momordica charantia*, *Food Chem.*, 118, 751-756.
- Loizzo, M.R., Menichini, F., and Tundis, R., 2013, Recent insights into the emerging role of triterpenoids in cancer therapy: Part I, *Elsevier*, 40, 1-31.
- Lourenço, S.C., Moldão-Martins, M., and Alves, V.D., 2019, Antioxidants of natural plant origins: From sources to food industry applications, *Molecules*, 24, 14-16.
- Mabberley, D.J. and Kubitzki, K., 2011, *Flowering plants. Eudicots: Sapindales, Cucurbitales, Myrtaceae*, Springer, Heidelberg.
- Martin, L., Birdsell, L., Macdonald, N., Reiman, T., Clandinin, M.T., McCargar, L.J., Murphy, R., Ghosh, S., Sawyer, M.B., and Baracos, V.E., 2013, Cancer cachexia in the age of obesity: skeletal muscle depletion is a powerful prognostic factor, independent of body mass index., *J. Clin. Oncol.*, 31, 1539-1547.
- van Meerloo, J., Kaspers, G.J.L., and Cloos, J., 2011, Cell sensitivity assays: the MTT assay., *Methods Mol. Biol.*, 731, 237-245.
- Miller, N.J., Diplock, A.T., Rice-Evans, C., Davies, M.J., Gopinathan, V., and Milner, A., 1993, A novel method for measuring antioxidant capacity and its application to monitoring the antioxidant status in premature neonates, *Clin. Sci.*, 84, 407-412.
- Mohamad, K., Martin, M.-T., Najdar, H., Gaspard, C., Sévenet, T., Awang, K., Hadi, H., and Païs, M., 1999, Cytotoxic 3,4-Secoapotirucallanes from *Aglaia argentea* Bark, *J. Nat. Prod.*, 62, 868-872.
- Müller, C., Schulte, F.W., Lange-Grünweller, K., Obermann, W., Madhugiri, R., Pleschka, S., Ziebuhr, J., Hartmann, R.K., and Grünweller, A., 2018, Broad-spectrum antiviral activity of the eIF4A inhibitor silvestrol against corona- and picornaviruses, *Antiviral Res.*, 150, 123-129.
- Newsholme, P., Cruzat, V.F., Keane, K.N., Carlessi, R., and De Bittencourt, P.I.H., 2016, Molecular mechanisms of ROS production and oxidative stress in diabetes, *Biochem. J.*, 473, 4527-4550.
- Nugroho, B.W., Güssregen, B., Wray, V., Witte, L., Bringmann, G., and Proksch, P., 1997, Insecticidal rocaglamide derivatives from *Aglaia elliptica* and *A. harmsiana*, *Phytochemistry*, 45, 1579-1585.
- Nurhayati, S. and Lusiyanti, Y., 2014, Apoptosis dan respon biologik sel sebagai, *Iptek Ilm. Pop.*, 7, 57-66.
- Oktaviani, D., Sukmawati, W., Farabi, K., Harneti, D., Nurlelasari, N., Darwati, D., Maharani, R., Mayanti, T., Safari, A., and Supratman, U., 2022, Terpenoids from the Stem bark of *Aglaia elaeagnoidea* and their cytotoxic activity against HeLa and DU145 Cancer Cell lines, *Molekul*, 17, 76-84.
- Othman, N., Pan, L., Mejin, M., Voong, J.C.L., Chai, H., Pannell, C.M.,

- Kinghorn, A.D., and Yeo, T.C., 2016, Cyclopenta[b]benzofuran and Secodammarane Derivatives from the Stems of *Aglaia stellatopilosa*, *J. Nat. Prod.*, 79, 784-791.
- Pan, L., Acuña, U.M., Li, J., Jena, N., Ninh, T.N., Pannell, C.M., Chai, H., Fuchs, J.R., Carcache de Blanco, E.J., Soejarto, D.D., and Kinghorn, A.D., 2013, Bioactive Flavaglines and Other Constituents Isolated from *Aglaia perviridis*, *J. Nat. Prod.*, 76, 394-404.
- Pan, L., Kardono, L.B.S., Riswan, S., Chai, H., Carcache De Blanco, E.J., Pannell, C.M., Soejarto, D.D., McCloud, T.G., Newman, D.J., and Kinghorn, A.D., 2010, Isolation and characterization of minor analogues of silvestrol and other constituents from a large-scale re-collection of *Aglaia foveolata*, *J. Nat. Prod.*, 73, 1873-1878.
- Pandey, A.K., Sonker, N., and Singh, P., 2016, Efficacy of Some Essential Oils Against *Aspergillus flavus* with Special Reference to *Lippia alba* Oil an Inhibitor of Fungal Proliferation and Aflatoxin B1 Production in Green Gram Seeds during Storage, *J. Food Sci.*, 81, 928-934.
- Prakash, O., Kumar, A., Kumar, P., and Ajeet, A., 2013, Anticancer Potential of Plants and Natural Products: A Review, *Am. J. Pharmacol. Sci.*, 1, 104-115.
- Prior, R.L., Wu, X., and Schaich, K., 2005, Standardized Methods for the Determination of Antioxidant Capacity and Phenolics in Foods and Dietary Supplements, *J. Agric. Food Chem.*, 53, 4290-4302.
- Ravanan, P., Singh, S.K., Rao, G.S., and Kondaiah, P., 2011, Growth inhibitory, apoptotic and anti-inflammatory activities displayed by a novel modified triterpenoid, cyano enone of methyl boswellates., *J. Biosci.*, 36, 297-307.
- Rebucci, M. and Michiels, C., 2013, Ac ce p te d us cr t, *Biochem. Pharmacol.*, 85, 1219-1226.
- Rice-Evans, C.A., Miller, N.J., and Paganga, G., 1996, Structure-antioxidant activity relationships of flavonoids and phenolic acids., *Free Radic. Biol. Med.*, 20, 933-956.
- Robinson, M.M. and Zhang, X., 2011, *The World Medicines Situation 2011 Traditional Medicines: Global Situation, Issues And Challenges*, 3rd Ed, WHO Press, Switzerland.
- Roux, D., Martin, M.-T., Adeline, M.-T., Sevenet, T., Hadi, A.H.A., and Païs, M., 1998, Foveolins A and B, dammarane triterpenes from *Aglaia foveolata*, *Phytochemistry*, 49, 1745-1748.
- Rudakov, O.B., Rudakova, L. V., and Selemenev, V.F., 2018, Acetonitrile as tops solvent for liquid chromatography and extraction, *J. Anal. Chromatogr. Spectrosc.*, 1, 864-883 .
- Salim, A.A., Chai, H.-B., Rachman, I., Riswan, S., Kardono, L.B.S., Farnsworth, N.R., Carcache-Blanco, E.J., and Kinghorn, A.D., 2007, Constituents of the leaves and stem bark of *Aglaia foveolata*, *Tetrahedron*, 63, 7926-7934.

- Schulz, G., Victoria, C., Kirschning, A., and Steinmann, E., 2021, Rocaglamide and silvestrol: a long story from anti-tumor to anti-coronavirus compounds, *Nat. Prod. Rep.*, 38, 12-18.
- Shahidi, F. and Zhong, Y., 2015, Measurement of antioxidant activity, *J. Funct. Foods*, 18, 757-781.
- Shang, Y. and Huang, S., 2019, Multi-omics data-driven investigations of metabolic diversity of plant triterpenoids, *Plant J.*, 97, 101-111.
- Sianturi, J., Purnamasari, M., Darwati, Harneti, D., Mayanti, T., Supratman, U., Awang, K., and Hayashi, H., 2015, New bisamide compounds from the bark of *Aglaia eximia* (Meliaceae), *Phytochem. Lett.*, 13, 297-301.
- Sirivibulkovit, K., Nouanthavong, S., and Sameenoi, Y., 2018, Paper-based DPPH Assay for Antioxidant Activity Analysis, *Anal. Sci.*, 34, 795-800.
- Sun, Yujin, Yin, Y., Sun, Yunpeng, Li, Q., Cui, L., Xu, W., Kong, L., and Luo, J., 2021b, Aglatestine A, a Rearranged Limonoid with a 3/6/6 Tricarbocyclic Framework from the Fruits of *Aglaia edulis*, *J. Org. Chem.*, 86, 11263-11268.
- Sung, H., Ferlay, J., Siegel, R.L., Laversanne, M., Soerjomataram, I., Jemal, A., and Bray, F., 2021, Global Cancer Statistics 2020: GLOBOCAN Estimates of Incidence and Mortality Worldwide for 36 Cancers in 185 Countries, *CA. Cancer J. Clin.*, 71, 209-249.
- Tan, Y.-S., Baskaran, A., Nallathamby, N., Chua, K.-H., Kuppusamy, U.R., and Sabaratnam, V., 2015, Influence of customized cooking methods on the phenolic contents and antioxidant activities of selected species of oyster mushrooms (*Pleurotus* spp.), *J. Food Sci. Technol.*, 52, 3058-3064.
- Tang, Y., Zhao, Z.-Z., Yao, J.-N., Feng, T., Li, Z.-H., Chen, H.-P., and Liu, J.-K., 2018, Irpeksins A–E, 1,10-seco-Eburicane-Type Triterpenoids from the Medicinal Fungus *Irpex lacteus* and Their Anti-NO Activity, *J. Nat. Prod.*, 81, 2163-2168.
- Tiwari, P.K., Kaur, M., and Kaur, H., 2011, Phytochemical screening and Extraction: A Review, *Inter.Pharm.Sci.*, 1, 98-106.
- Tripathy, V., Basak, B.B., Varghese, T.S., and Saha, A., 2015, Residues and contaminants in medicinal herbs-A review, *Phytochem. Lett.*, 14, 67-78.
- Wang, X., Qiu, Y., Wen, H., Weng, R., Chen, D., and Liu, H., 2023, Separation and structural characterization of unknown impurity in vancomycin by two-dimensional preparative liquid chromatography, LC-MS and NMR, *J. Pharm. Biomed. Anal.*, 232, 115403-115415.
- Wong, H.L., Bendayan, R., Rauth, A.M., and Wu, X.Y., 2006, Simultaneous delivery of doxorubicin and GG918 (Elacridar) by new polymer-lipid hybrid nanoparticles (PLN) for enhanced treatment of multidrug-resistant breast cancer, *J. Control. Release Soc.*, 116, 275-284.
- Xin, D.-Y., Jones, P.-J., and Gonnella, N.-C., 2018, DiCE: Diastereomeric in

Silico Chiral Elucidation, Expanded DP4 Probability Theory Method for Diastereomer and Structural Assignment, *J. Org. Chem.*, 83, 5035-5043.

Yodsaoue, O., Sonprasit, J., Karalai, C., Ponglimanont, C., Tewtrakul, S., and Chantrapromma, S., 2012, Diterpenoids and triterpenoids with potential anti-inflammatory activity from the leaves of *Aglaia odorata*, *Phytochemistry*, 76, 83-91.

Zhang, B., Xing, Y., Wen, C., Yu, X., Sun, W., Xiu, Z., and Dong, Y., 2017, Pentacyclic triterpenes as α -glucosidase and α -amylase inhibitors: Structure-activity relationships and the synergism with acarbose, *Bioorg. Med. Chem. Lett.*, 27, 5065-5070.

Zhang, Q.-W., Lin, L.-G., and Ye, W.-C., 2018, Techniques for extraction and isolation of natural products: a comprehensive review, *Chin. Med.*, 13, 20-46.

Zou, Y.H., Liu, W.T., Zhang, J.X., and Xiang, D.C., 2017, Triterpenoids from the bark of *Dysoxylum hainanense* and their anti-inflammatory and radical scavenging activity, *Fitoterapia*, 121, 159-163.