



DAFTAR PUSTAKA

- Abdillah, A. (2018). Kajian Kerentanan Airtanah terhadap Pencemar di Daerah Aliran Sungai Serang. *Skripsi*. Yogyakarta : Fakultas Geografi UGM.
- Adhesti, Y. M. P., & Widyastuti, M. (2021). Study of Groundwater Vulnerability to Contamination in Part of the Klaten District Area, Central Java-Indonesia. In *E3S Web of Conferences* (Vol. 325, p. 02006). EDP Sciences.
- Aller, L., Lehr, J. H., Petty, R. (1987). DRASTIC : A standardized system for evaluation groundwater pollution potential using hydrogeologic settings. Oklahoma : U.S. Environmental Protection Agency.
- Almasri, M. N. (2008). Assessment of intrinsic vulnerability to contamination for Gaza coastal aquifer, Palestine. *Journal of Environmental Management*, 88(4), 577-593.
- Anane, M., Abidi, B., Lachaal, F., Limam, A., & Jellali, S. (2013). GIS-based DRASTIC, Pesticide DRASTIC and the Susceptibility Index (SI): comparative study for evaluation of pollution potential in the Nabeul-Hammamet shallow aquifer, Tunisia. *Hydrogeology Journal*, 21(3), 715.
- Asadi, P., Ataie-Ashtiani, B., & Beheshti, A. (2017). Vulnerability assessment of urban groundwater resources to nitrate: the case study of Mashhad, Iran. *Environmental Earth Sciences*, 76(1), 41.
- Ayed, B., Jmal, I., Sahal, S., & Bouri, S. (2017). Assessment of groundwater vulnerability using a specific vulnerability method: case of Maritime Djefara shallow aquifer (Southeastern Tunisia). *Arabian Journal of Geosciences*, 10(12), 1-14.
- Baalousha, H. M. (2016). Groundwater Vulnerability Mapping of Qatar Aquifers. *Journal of African Earth Sciences*, vol.124, pp : 75-93, doi: <https://doi.org/10.1016/j.jafrearsci.2016.09.017>.
- Babiker, I. S., Mohamed, M. A., Hiyama, T., & Kato, K. (2005). A GIS-based DRASTIC model for assessing aquifer vulnerability in Kakamigahara Heights, Gifu Prefecture, central Japan. *Science of the Total Environment*, 345(1-3), 127-140.
- Badan Standardisasi Nasional (BSN).2002. Penyusunan Neraca Sumberdaya Bagian 1 : Sumberdaya Spasial. SNI 19-6728.1-2002.
- Bappeda Sragen. 2018. *Laporan Akhir Kegiatan Penyusunan dan Analisis Informasi*



- Bartzas, G., Tinivella, F., Medini, L., Zaharaki, D., & Komnitsas, K. (2015). Assessment of groundwater contamination risk in an agricultural area in north Italy. *Information Processing in Agriculture*, 2(2), 109-129.
- Cahyadi, A., Haryono, E., Adji, T. N., Widyastuti, M., Naufal, M., Ramadhan, F., ... & Riyanto, I. A. (2020). Analisis Konektivitas dan Karakteristik Lorong pada Sistem Hidrogeologi Mataair Beton, Kawasan Karst Gunungsewu, Kabupaten Gunungkidul dengan Uji Permutasi. *Jurnal Geografi*, 12(2), 105-114.
- DeVellis, R. F. (2003). *Scale development: Theory and applications* (2nd ed.). New York : SAGE Publication.
- DPMPTSP. 2020. Pembangunan Kawasan Industri Kecamatan Sumberlawang. Diakses melalui <https://sipelangi.sragenkab.go.id/investasi/detail/42> pada 1 januari 2022 pukul 08.30 WIB.
- Effendi, H. 2003. *Telaah Kualitas Air*. Yogyakarta : Kanisius.
- Febriarta, E., Larasti, A. 2020. Karakteristik Akuifer Air Tanah Dangkal di Enpdan Muda Merapi Yogyakarta. *Jurnal Sains dan Teknologi Lingkungan*. Vol. 12 No. 2. Hal 84.
- Febriarta, E., & Purnama, S. (2020). Identifikasi Keterdapatannya Airtanah Dengan Electromagnetic Very Low Frequency (EM-VLF) di Non Cekungan Airtanah Kecamatan Ungaran Timur. *Jurnal Geosains dan Teknologi*, 3(2), 52-62.
- Gaieb, S., & Hamza, M. H. (2013). Assessing vulnerability to agricultural pollution of groundwater Bou Arada Laroussa according to SI method applied by GIS. *Journal of Research in Environmental and Earth Sciences*, 1(2013), 01-10.
- Ghouili, N., Jarraya-Horriche, F., Hamzaoui-Azaza, F., Zaghrarni, M. F., Ribeiro, L., & Zammouri, M. (2021). Groundwater vulnerability mapping using the Susceptibility Index (SI) method: Case study of Takelsa aquifer, Northeastern Tunisia. *Journal of African Earth Sciences*, 173, 104035.
- Gogu, R. C., & Dassargues, A. (2000). Current trends and future challenges in groundwater vulnerability assessment using overlay and index methods. *Environmental geology*, 39, 549-559.
- Harter, T., dan Walker, L.G. 2001. *Booklet: Assessing Vulnerability of Groundwater*. California Department of Health Services.



Hendrayana, H. (2010). Pengendalian Daya Rusak Airtanah. *Journal UGM*, Vol 4 : 1 - 15.

Herlambang, A. (2006). Pencemaran air dan strategi penggulangannya. *Jurnal Air Indonesia*, 2(1).

Hsu, C. C., & Sandford, B. A. (2007). The Delphi technique: making sense of consensus. *Practical Assessment, Research, and Evaluation*, 12(1), 10.

Hutasoit, Lambok M., 2009. Kondisi Permukaan Air Tanah dengan dan tanpa peresapan buatan di daerah Bandung: Hasil Simulasi Numerik. *Jurnal Geologi Indonesia*, Vol. 4 No. 3 : 177-188.

Jang, W. S., Engel, B., Harbor, J., Theller, L. (2017). Aquifer vulnerability assessment for sustainable groundwater management using DRASTIC. *Journal Water*, 9 (10), 792.

Karan, S. K., Samadder, S. R., & Singh, V. (2018). Groundwater vulnerability assessment in degraded coal mining areas using the AHP–Modified DRASTIC model. *Land degradation & development*, 29(8), 2351-2365.

Kerlinger, F. N., Lee, H. B. (2000). *Foundations of behavioral research*. New York: Wadsworth Publishing.

Kumar, P., Bansod, B. K., Debnath, S. K., Thakur, P. K., & Ghanshyam, C. (2015). Index-based groundwater vulnerability mapping models using hydrogeological settings: a critical evaluation. *Environmental Impact Assessment Review*, 51, 38-49. Maria, R. (2018, February). Comparative studies of groundwater vulnerability assessment. In *IOP Conference Series: Earth and Environmental Science* (Vol. 118, No. 1, p. 012018). IOP Publishing.

Ncibi, K., Chaar, H., Hadji, R., Baccari, N., Sebei, A., Khelifi, F., & Hamed, Y. (2020). A GIS-based statistical model for assessing groundwater susceptibility index in shallow aquifer in Central Tunisia (Sidi Bouzid basin). *Arabian Journal of Geosciences*, 13(2), 1-21.

Neshat, A., Pradhan, B., Pirasteh, S., & Shafri, H. Z. M. (2014). Estimating groundwater vulnerability to pollution using a modified DRASTIC model in the Kerman agricultural area, Iran. *Environmental earth sciences*, 71, 3119-3131.

Noori, R., Ghahremanzadeh, H., Kløve, B., Adamowski, J. F., & Baghvand, A. (2019). Modified-DRASTIC, modified-SINTACS and SI methods for groundwater vulnerability assessment in the southern Tehran aquifer. *Journal of Environmental Science and Health, Part A*, 54(1), 89-100.

Peraturan Daerah Kabupaten Sragen No 1 Tahun 2020 tentang rencana Tata Ruang Wilayah Kabupaten Sragen Tahun 2010 - 2030.



Presiden Republik Indonesia. 2019. Undang – Undang No 19 Tentang Sumberdaya Air.

Putranto, T. T., Ali, R. K., & Putro, A. B. (2019). Studi Kerentanan Airtanah terhadap Pencemaran dengan Menggunakan Metode DRASTIC pada Cekungan Airtanah (CAT) Karanganyar-Boyolali, Provinsi Jawa Tengah. *Jurnal Ilmu Lingkungan*, 17(1), 159.

Putranto, T. T., & Marjuanto, A. A. (2021). Assessment of Contamination Vulnerability of Groundwater Using Susceptibility Index Method: Semarang City as Case Study. *Journal of Engineering Science and Technology*, 16(1), 438-454.

Purnama S. 2016. *Hidrologi Airtanah*. Yogyakarta : Penerbit PT Kanisius (Anggota IKAPI).

Purnama S. 2020. *Airtanah dan Intrusi Air Laut*. Yogyakarta : Penerbit PT Kanisius (Anggota IKAPI).

Putranto, T. T., & Kusuma, K. I. (2009). Permasalahan airtanah pada daerah urban. *Teknik*, 30(1), 48-56.

Putranto, T. T., Ali, R. K., & Putro, A. B. (2019). Studi Kerentanan Airtanah terhadap Pencemaran dengan Menggunakan Metode DRASTIC pada Cekungan Airtanah (CAT) Karanganyar-Boyolali, Provinsi Jawa Tengah. *Jurnal Ilmu Lingkungan*, 17(1), 159.

Putranto, T. T., Budihardjo, M. A., & Sarminingsih, A. (2022) Aplikasi Metode DRASTIC untuk Analisis Kerentanan Air Tanah terhadap Pencemaran di Kabupaten Tanah Laut, Kabupaten Banjar, dan Kota Banjarbaru, Provinsi Kalimantan Selatan. *TEKNIK*, 43(1), 45-56.

Rahman, M., Haque, M. M., & Tareq, S. M. (2021). Appraisal of groundwater vulnerability in south-central part of Bangladesh using DRASTIC model: An approach towards groundwater protection and health safety. *Environmental Challenges*, 5, 100391.

Ribeiro, L. (2000). SI: a new index of aquifer susceptibility to agricultural pollution. *ERSHA/CVRM, Instituto Superior Técnico, Lisboa, Portugal*.

Ribeiro, L., Pindo, J. C., & Dominguez-Granda, L. (2017)Assessment of Groundwater Vulnerability in The Daule Aquifer, Ecuador, Using The Susceptibility Index Method, *Science of The Total Environment*, 574, 1674-1683,

Riyadi, A. (2007). Karakteristik Airtanah di Kecamatan Tamansari Kota Tasikmalaya. *Jurnal Teknologi Lingkungan*, 8(3), 197 - 206.

Riyanto, I. A., & Widyastuti, M. (2016). Kerentanan intrinsik dan spesifik airtanah terhadap



pencemaran di kecamatan banjarnegara dan sekitarnya. *Jurnal Bumi Indonesia*, 5(4), 1 - 17.

Riyanto, I. A. (2016). Kerentanan intrinsik dan spesifik airtanah terhadap pencemaran di kecamatan banjarnegara dan sekitarnya. Skripsi. Yogyakarta : Fakultas Geografi UGM.

Rosen, L. (1994). A study of the DRASTIC methodology with emphasis on Swedish conditions. *Groundwater*, 32(2), 278-285.

Singha, S. S., Pasupuleti, S., Singha, S., Singh, R., & Venkatesh, A. S. (2019). A GIS-based modified DRASTIC approach for geospatial modeling of groundwater vulnerability and pollution risk mapping in Korba district, Central India. *Environmental Earth Sciences*, 78(21), 1-19.

Sartohadi J., Suratman., Jamulya., Dewi N.I.S., .2016. *Pengantar Geografi Tanah*. Yogyakarta : Pustaka Media.

Siregar, J. P. (2007). *Pendugaan geolistrik resistivitas sounding dalam penyelidikan lapisan akiferairtanah di Kabupaten Sragen*. Skripsi : Fakultas MIPA Universitas Sebelas Maret.

Stigter, T. Y., Ribeiro, L., & Dill, A. C. (2005). Evaluation of an intrinsic and a specific vulnerability assessment method in comparison with groundwater salinisation and nitrate contamination levels in two agricultural regions in the south of Portugal. *Hydrogeology Journal*, 14(1-2), 79-99.

Sudarmadji. 2013. *Mata Air*. Yogyakarta : Sekolah Pasca Sarjana UGM.

Sudarmaji, Hadi P, Wisyastuti M. 2016. *Pengelolaan Sumberdaya Air Terpadu*. Yogyakarta : Universitas Gadjah Mada Press.

Sugianti, K., Mulyadi, D., & Maria, R. (2017). Analisis kerentanan pencemaran airtanah dengan pendekatan metode drastic di Bandung Selatan. *Jurnal lingkungan dan bencana geologi*, 7(1), 19-33.

Suprayogi, S., Purnama, L.S., Darmanto, D. 2016. *Pengelolaan Daerah Aliran Sungai*. Yogyakarta:Universitas Gadjah Mada Press.

Todd, D.K, 1980. *Groundwater Hydrology*. New York.John Wiley and Sons Inc..

Voudouris, K., Kazakis, N., Polemio, M., Kareklas, K. (2010). Assessment of intrinsic vulnerability using the DRASTIC model and GIS in the kiti aquifer, cyprus. *European Water* 30. Hal : 13 – 28.



Widyastuti, M., Notosiswoyo, S., & Anggayana, K. (2006). Pengembangan Metode 'Drastic' untuk Prediksi Kerentanan Airtanah Bebas terhadap Pencemaran di Sleman. *Majalah Geografi Indonesia*, 20(2006).

Wijaya, Krishna A., and Ig L. S. Purnama. 2018. "Kajian Kerentanan Airtanah terhadap Potensi Pencemaran di Kecamatan Kasihan Kabupaten Bantul." *Jurnal Bumi Indonesia*, vol. 7, no. 1.