

DAFTAR PUSTAKA

- Aalim, H., Wang, D., & Luo, Z. (2021). Black rice (*Oryza sativa L.*) processing: Evaluation of physicochemical properties, in vitro starch digestibility, and phenolic functions linked to type 2 diabetes. *Food Research International*, 141(May 2020), 109898. <https://doi.org/10.1016/j.foodres.2020.109898>
- Abd Razak, D. L., Abd Rashid, N. Y., Jamaluddin, A., Sharifudin, S. A., Abd Kahar, A., & Long, K. (2017). Cosmeceutical potentials and bioactive compounds of rice bran fermented with single and mix culture of *Aspergillus oryzae* and *Rhizopus oryzae*. *Journal of the Saudi Society of Agricultural Sciences*, 16(2), 127–134. <https://doi.org/10.1016/j.jssas.2015.04.001>
- Abdulrachman, S., Karsono, S., Samaullah, M. Y., & Sembiring, H. (2011). *Prosedur Operasional Standar (POS) Budi Daya Padi Sawah*. Balai Besar Penelitian Tanaman Padi Kementerian Pertanian.
- Agu, C. M., Orakwue, C. C., Ani, K. A., Agulanna, A. C., & Dzarma, G. W. (2022). Process thermodynamic, kinetics and statistical analysis of Gmelina arborea seed oil extraction as clean environmental friendly industrial base-oil. *Case Studies in Chemical and Environmental Engineering*, 6(April), 100216. <https://doi.org/10.1016/j.csee.2022.100216>
- Ahmad, I., Prabowo, W. C., Nur, Y., Ardana, M., Rahayu, B. P., & Herman. (2020). OPTIMASI METODE EKSTRAKSI BERBANTU MIKROWAVE DENGAN PELARUT HIJAU (ASAM SITRAT-GLUKOSA) TERHADAP KADAR POLIFENOL TOTAL DARI DAUN KADAMBA (*Mitragyna speciosa* Korth. Havil) MENGGUNAKAN RESPONSE SURFACE METHODOLOGY. *Majalah Farmasi Dan Farmakologi*, 24(1), 11–16.
- Airouyuwa, J. O., Mostafa, H., Riaz, A., Stathopoulos, C., & Maqsood, S. (2023). Natural Deep Eutectic Solvents and Microwave-Assisted Green Extraction for Efficient Recovery of Bioactive Compounds from By-Products of Date Fruit (*Phoenix dactylifera L.*) Processing: Modeling, Optimization, and Phenolic Characterization. *Food and Bioprocess Technology*, 16(4), 824–843. <https://doi.org/10.1007/s11947-022-02960-8>
- Alamsyah, D., & Pratama, D. (2019). Segmentasi Warna Citra Bunga Daisy dengan Algoritma K-Means pada Ruang Warna Lab. *Jurnal Buana Informatika*, 10(2), 153. <https://doi.org/10.24002/jbi.v10i2.2458>
- Alara, O. R., & Abdurahman, N. H. (2019). Microwave-assisted extraction of phenolics from *Hibiscus sabdariffa* calyces: Kinetic modelling and process intensification. *Industrial Crops and Products*, 137(May), 528–535. <https://doi.org/10.1016/j.indcrop.2019.05.053>
- Alara, O. R., Abdurahman, N. H., & Abdul Mudalip, S. K. (2019). Optimizing Microwave-Assisted Extraction Conditions to Obtain Phenolic-Rich Extract from *Chromolaena odorata* Leaves. *Chemical Engineering and Technology*, 42(9), 1733–1740. <https://doi.org/10.1002/ceat.201800462>
- Ariyati, E. (2015). PENGARUH KONSENTRASI ASAM SITRAT TERHADAP KARAKTERISTIK EKSTRAK ANTOSIANIN DAUN JATI SERTA UJI STABILITASNYA DALAM ES KRIM. *Prosiding Seminar Nasional Pendidikan Biologi 2015, Yang Diselenggarakan Oleh Prodi Pendidikan*

- Biologi FKIP Universitas Muhammadiyah Malang, Tema: “Peran Biologi Dan Pendidikan Biologi Dalam Menyiapkan Generasi Unggul Dan Berdaya Saing Global”, Malang, 21, 4(2007), 339–345.*
- Aydar, A. Y., Aydın, T., Yılmaz, T., Kothakota, A., Claudia Terezia, S., Florin Leontin, C., & Pandiselvam, R. (2022). Investigation on the influence of ultrasonic pretreatment on color, quality and antioxidant attributes of microwave dried *Inula viscosa* (L.). *Ultrasonics Sonochemistry*, 90(August), 106184. <https://doi.org/10.1016/j.ultsonch.2022.106184>
- Bansod, S. P., Parikh, J. K., & Sarangi, P. K. (2023). Pineapple peel waste valorization for extraction of bio-active compounds and protein: Microwave assisted method and Box Behnken design optimization. *Environmental Research*, 221(December 2022), 115237. <https://doi.org/10.1016/j.envres.2023.115237>
- Bener, M., Şen, F. B., Önem, A. N., Bekdeşer, B., Çelik, S. E., Lalikoglu, M., Aşçı, Y. S., Capanoglu, E., & Apak, R. (2022). Microwave-assisted extraction of antioxidant compounds from by-products of Turkish hazelnut (*Corylus avellana* L.) using natural deep eutectic solvents: Modeling, optimization and phenolic characterization. *Food Chemistry*, 385(February). <https://doi.org/10.1016/j.foodchem.2022.132633>
- Bertoia, M. L., Rimm, E. B., Mukamal, K. J., Hu, F. B., Willett, W. C., & Cassidy, A. (2016). Dietary flavonoid intake and weight maintenance: Three prospective cohorts of 124 086 US men and women followed for up to 24 years. *BMJ (Online)*, 352. <https://doi.org/10.1136/bmj.i17>
- Bhat, F. M., & Riar, C. S. (2017). Characterizing the pigmented traditional rice cultivars grown in temperate regions of Kashmir (India) for free and bound phenolics compounds and in vitro antioxidant properties. *Journal of Cereal Science*, 76, 253–262. <https://doi.org/10.1016/j.jcs.2017.06.018>
- Bouras, M., Chadni, M., Barba, F. J., Grimi, N., Bals, O., & Vorobiev, E. (2015). Optimization of microwave-assisted extraction of polyphenols from *Quercus* bark. *Industrial Crops and Products*, 77, 590–601. <https://doi.org/10.1016/j.indcrop.2015.09.018>
- Cassol, L., Rodrigues, E., & Zapata Noreña, C. P. (2019). Extracting phenolic compounds from *Hibiscus sabdariffa* L. calyx using microwave assisted extraction. *Industrial Crops and Products*, 133(August 2018), 168–177. <https://doi.org/10.1016/j.indcrop.2019.03.023>
- Castañeda-Ovando, A., Pacheco-Hernández, M. de L., Páez-Hernández, M. E., Rodríguez, J. A., & Galán-Vidal, C. A. (2009). Chemical studies of anthocyanins: A review. *Food Chemistry*, 113(4), 859–871. <https://doi.org/10.1016/j.foodchem.2008.09.001>
- Chemat, F., Abert Vian, M., & Zill-E-Huma. (2009). Microwave assisted Extraction for Bioactive Compounds. In *Green Chemistry Research Trends*. Springer.
- Chen, P. N., Kuo, W. H., Chiang, C. L., Chiou, H. L., Hsieh, Y. S., & Chu, S. C. (2006). Black rice anthocyanins inhibit cancer cells invasion via repressions of MMPs and u-PA expression. *Chemico-Biological Interactions*, 163(3), 218–229. <https://doi.org/10.1016/j.cbi.2006.08.003>
- Cheong, K.-L., Wang, L.-Y., Wu, D.-T., Hu, D., Zhao, J., & Li, S.-P. (2016).

- Microwave-Assisted Extraction, Chemical Structures, and Chain Conformation of Polysaccharides from a Novel *Cordyceps Sinensis* Fungus UM01. *Journal of Food Science*, 81(9), 2167–2174.
- Colombo, F., Cappa, C., Bani, C., Magni, M., Biella, S., Restani, P., & Di Lorenzo, C. (2023). Characterization of color, phenolic profile, and antioxidant activity of Italian pigmented rice varieties after different technological treatments. *Food Bioscience*, 53(April), 102674. <https://doi.org/10.1016/j.fbio.2023.102674>
- Das, K., Medhabati, K., Nongalleima, K., & Devi, H. (2014). The Potential of Dark Purple Scented Rice- From Staple Food to Nutraceutical. *Current World Environment*, 9(3), 867–876. <https://doi.org/10.12944/cwe.9.3.38>
- Enggiwanto, S., Istiqomah, F., Daniati, K., Roanisca, O., & Mahardika, R. G. (2018). EKSTRAKSI DAUN PELAWAN (*Tristaniaopsis merguensis*) SEBAGAI ANTIOKSIDAN MENGGUNAKAN MICROWAVE ASSISTED EXTRACTION (MAE). *Indonesian Journal of Pure and Applied Chemistry*, 1(2), 50. <https://doi.org/10.26418/indonesian.v1i2.30528>
- Fathordoobady, F., Mirhosseini, H., Selamat, J., & Manap, M. Y. A. (2016). Effect of solvent type and ratio on betacyanins and antioxidant activity of extracts from *Hylocereus polyrhizus* flesh and peel by supercritical fluid extraction and solvent extraction. *Food Chemistry*, 202, 70–80. <https://doi.org/10.1016/j.foodchem.2016.01.121>
- Fatmawati, S., Nugraheni, F., & Bariroh, T. (2018). Optimasi Waktu dan Konsentrasi Etanol Pada Ekstraksi Berbantu Ultrasonik Serta Penetapan Kadar Kafein Daun Kopi Arabika (*Coffea arabica* L.). *Jurnal Fakultas Farmasi Dan Sains* 1, 1–6.
- Ferreira, L. F., Minuzzi, N. M., Rodrigues, R. F., Pauletto, R., Rodrigues, E., Emanuelli, T., & Bochi, V. C. (2020). Citric acid water-based solution for blueberry bagasse anthocyanins recovery: Optimization and comparisons with microwave-assisted extraction (MAE). *Lwt*, 133(May), 110064. <https://doi.org/10.1016/j.lwt.2020.110064>
- Firdaus, M. J., Purwoko, B. S., Dewi, I. S., & Suwarno, W. B. (2022). Karakterisasi Fisikokimia Beras Galur-galur Padi Hitam Dihaploid. *Jurnal Agronomi Indonesia (Indonesian Journal of Agronomy)*, 50(1), 1–9. <https://doi.org/10.24831/jai.v50i1.39850>
- Garcia, M., Knuutila, H. K., Aronu, U. E., & Gu, S. (2018). Influence of substitution of water by organic solvents in amine solutions on absorption of CO₂. *International Journal of Greenhouse Gas Control*, 78(November 2017), 286–305. <https://doi.org/10.1016/j.ijggc.2018.07.029>
- Ghosh, S., Deng, H., & Luan, Y. (2007). Error sum of squares and power comparisons for model search, identification and discrimination. *Journal of Statistical Planning and Inference*, 137(11), 3475–3482. <https://doi.org/10.1016/j.jspi.2007.03.025>
- Gómez, A. V., Tadini, C. C., Biswas, A., Buttrum, M., Kim, S., Boddu, V. M., & Cheng, H. N. (2019). Microwave-assisted extraction of soluble sugars from banana puree with natural deep eutectic solvents (NADES). *Lwt*, 107(December 2018), 79–88. <https://doi.org/10.1016/j.lwt.2019.02.052>

- Halee, A., Supavititpatana, P., Ruttarattanamongkol, K., Jittrepotch, N., Rojsuntornkitti, K., & Kongbangkerd, T. (2018). Effects of solvent types and citric acid concentrations on the extraction of antioxidants from the black rice bran of *Oryza sativa L.* CV. Hom Nin. *Journal of Microbiology, Biotechnology and Food Sciences*, 8(2), 765–769. <https://doi.org/10.15414/jmbfs.2018.8.2.765-769>
- Hao, J., Zhu, H., Zhang, Z., Yang, S., & Li, H. (2015). Identification of anthocyanins in black rice (*Oryza sativa L.*) by UPLC/Q-TOF-MS and their in vitro and in vivo antioxidant activities. *Journal of Cereal Science*, 64, 92–99. <https://doi.org/10.1016/j.jcs.2015.05.003>
- Hasdar, M., & Wadli. (2021). Ekstraksi Beras Hitam Sirampog Berbantu Gelombang Mikro (Microwave Assisted Extraction (Mae)) Extraction of Black Rice From Sirampog Using Microwave Assisted Extraction (Mae) Method. *Jurnal Pengolahan Pangan*, 6(2), 49–53.
- Hernando, D., Widodo, A. W., & Dewi, C. (2020). Pemanfaatan Fitur Warna dan Fitur Tekstur untuk Klasifikasi Jenis Penggunaan Lahan pada Citra Drone. *Jurnal Pengembangan Teknologi Informasi Dan Ilmu Komputer*, 4(2), 614–621. <http://j-ptiik.ub.ac.id>
- Izza, H. F., Susanti, D. Y., Mariyam, S., & Saputro, A. D. (2023). Performance of microwave-assisted extraction of proanthocyanidins from red sorghum grain in various power and citric acid concentration. *Journal of the Saudi Society of Agricultural Sciences*, xxxx. <https://doi.org/10.1016/j.jssas.2023.05.002>
- Jenssen, R., & Eltoft, T. (2008). A new information theoretic analysis of sum-of-squared-error kernel clustering. *Neurocomputing*, 72(1–3), 23–31. <https://doi.org/10.1016/j.neucom.2008.03.017>
- Jisieike, C. F., & Betiku, E. (2020). Rubber seed oil extraction: Effects of solvent polarity, extraction time and solid-solvent ratio on its yield and quality. *Biocatalysis and Agricultural Biotechnology*, 24(January), 101522. <https://doi.org/10.1016/j.bcab.2020.101522>
- Kaderides, K., Papaoikonomou, L., Serafim, M., & Goula, A. M. (2019). Microwave-assisted extraction of phenolics from pomegranate peels: Optimization, kinetics, and comparison with ultrasounds extraction. *Chemical Engineering and Processing - Process Intensification*, 137(November 2018), 1–11. <https://doi.org/10.1016/j.cep.2019.01.006>
- Kalt, W., Cassidy, A., Howard, L. R., Krikorian, R., Stull, A. J., Tremblay, F., & Zamora-Ros, R. (2020). Recent Research on the Health Benefits of Blueberries and Their Anthocyanins. *Advances in Nutrition*, 11(2), 224–236. <https://doi.org/10.1093/advances/nmz065>
- Kurniasari, L., Hartati, I., & Ratnani, R. D. (2006). Kajian Ekstraksi Minyak Jahe Assisted Extraction (Mae). *Momentum*, 4(1952), 47–52.
- Kusumawati, R., Palupi, N. S., & Budijanto, S. (2022). Aktivitas Antioksidan Makaroni Beras Hitam dan Kacang-Kacangan Berpigmen Metode Cold Extrusion. *AgriTECH*, 42(3), 195. <https://doi.org/10.22146/agritech.56064>
- Lang, G. H., Lindemann, I. da S., Ferreira, C. D., Hoffmann, J. F., Vanier, N. L., & de Oliveira, M. (2019). Effects of drying temperature and long-term storage conditions on black rice phenolic compounds. *Food Chemistry*,

- 287(September 2018), 197–204.
<https://doi.org/10.1016/j.foodchem.2019.02.028>
- Lee, J., Durst, R. W., & Wrolstad, R. E. (2005). Determination of total monomeric anthocyanin pigment content of fruit juices, beverages, natural colorants, and wines by the pH differential method: Collaborative study. *Journal of AOAC International*, 88(5), 1269–1278. <https://doi.org/10.1093/jaoac/88.5.1269>
- Loypimai, P., Moongngarm, A., & Chottanom, P. (2016). Thermal and pH degradation kinetics of anthocyanins in natural food colorant prepared from black rice bran. *Journal of Food Science and Technology*, 53(1), 461–470. <https://doi.org/10.1007/s13197-015-2002-1>
- Maharani, B. C., Lindriati, T., & Diniyah, N. (2016). Pengaruh Variasi Waktu Blanching dan Konsentrasi Asam Sitrat Terhadap Karakteristik dan Aktivitas Ekstrak Pigmen Ubi Jalar Ungu (*Ipomoea batatas L.*). *Jurnal Penelitian Pangan (Indonesian Journal of Food Research)*, 1(1), 60–67. <https://doi.org/10.24198/jp2.2016.vol1.1.10>
- Mangiri, J., Mayulu, N., & Kawengian, S. E. S. (2016). GAMBARAN KANDUNGAN ZAT GIZI PADA BERAS HITAM (*Oryza sativa L.*) KULTIVAR PARE AMBO SULAWESI SELATAN. *Jurnal E-Biomedik*, 4(1), 2–6. <https://doi.org/10.35790/ebm.4.1.2016.11050>
- Markakis, P. (1982). *Anthocyanins as Food Colors*. Academic Press.
- Mazza, G., & Miniati, E. (2018). *Anthocyanins in Fruits, Vegetables, and Grains*. CRC Press.
- Moldovan, B., David, L., Chisbora, C., & Cimpoiu, C. (2012). Degradation kinetics of anthocyanins from european cranberrybush (*viburnum opulus l.*) fruit extracts. Effects of temperature, pH and storage solvent. *Molecules*, 17(10), 11655–11666. <https://doi.org/10.3390/molecules171011655>
- Mustapa, A. N., Martin, A., Gallego, J. R., Mato, R. B., & Cocero, M. J. (2015). Microwave-assisted extraction of polyphenols from *Clinacanthus nutans* Lindau medicinal plant: Energy perspective and kinetics modeling. *Chemical Engineering and Processing: Process Intensification*, 97, 66–74. <https://doi.org/10.1016/j.cep.2015.08.013>
- Nurhayati, N., Oktaviani, O., & Basmal, J. (2018). Pengaruh Waktu Ekstraksi terhadap Mutu Ekstrak Cair Rumpun Laut *Gracilaria sp.* sebagai Bahan Baku Pupuk Cair. *Jurnal Pascapanen Dan Bioteknologi Kelautan Dan Perikanan*, 13(1), 33. <https://doi.org/10.15578/jpbkp.v13i1.505>
- Nurhidajah, N., Rosidi, A., Sya'di, Y. K., & Yonata, D. (2022). Efektivitas Berbagai Pelarut Organik Pada Ekstraksi Senyawa Fungsional Beras Hitam. *Agrointek: Jurnal Teknologi Industri Pertanian*, 16(1), 76–83. <https://doi.org/10.21107/agrointek.v16i1.12194>
- Ofoe, R., Gunupuru, L. R., & Abbey, Lord. (2022). Metabolites, elemental profile and chemical activities of *Pinus strobus* high temperature-derived pyrolygneous acid. *Chemical and Biological Technologies in Agriculture*, 9(1), 1–14. <https://doi.org/10.1186/s40538-022-00357-5>
- Pang, Y., Ahmed, S., Xu, Y., Beta, T., Zhu, Z., Shao, Y., & Bao, J. (2018). Bound phenolic compounds and antioxidant properties of whole grain and bran of white, red and black rice. *Food Chemistry*, 240(January 2017), 212–221.

<https://doi.org/10.1016/j.foodchem.2017.07.095>

- Peanparkdee, M., Patrawart, J., & Iwamoto, S. (2019). Effect of extraction conditions on phenolic content, anthocyanin content and antioxidant activity of bran extracts from Thai rice cultivars. *Journal of Cereal Science*, 86(November 2018), 86–91. <https://doi.org/10.1016/j.jcs.2019.01.011>
- Pramitasari, R., & Angelica, N. (2020). Ekstraksi, Pengeringan Semprot, dan Analisis Sifat Fisikokimia Antosianin Beras Hitam (*Oryza sativa L.*). *Jurnal Aplikasi Teknologi Pangan*, 9(2), 83–94. <https://doi.org/10.17728/jatp.5889>
- Putri, P. D. (2015). *EKSTRAKSI ANTOSIANIN BERAS HITAM (Oryza sativa L.) DENGAN BERBAGAI PELARUT DAN STABILITASNYA PADA SUHU TINGGI*. UNIVERSITAS GADJAH MADA.
- Rahim, M. A., Umar, M., Habib, A., Imran, M., Khalid, W., Mariana, C., Lima, G., Shoukat, A., Itrat, N., Nazir, A., Ejaz, A., Zafar, A., Awuchi, C. G., Sharma, R., Santana, R. F., & Emran, T. Bin. (2022). Photochemistry, Functional Properties, Food Applications, and Health Prospective of Black Rice. *Journal of Chemistry*, 2022, 21.
- Rajha, H. N., Darra, N. El, Hobaike, Z., Boussetta, N., Vorobiev, E., Maroun, R. G., & Louka, N. (2014). Extraction of Total Phenolic Compounds, Flavonoids, Anthocyanins and Tannins from Grape Byproducts by Response Surface Methodology. Influence of Solid-Liquid Ratio, Particle Size, Time, Temperature and Solvent Mixtures on the Optimization Process. *Food and Nutrition Sciences*, 05(04), 397–409. <https://doi.org/10.4236/fns.2014.54048>
- Ratnaningsih, N., & Ekawatiningsih, D. P. (2010). Potensi beras hitam sebagai sumber antosianin dan aplikasinya pada makanan tradisional yogyakarta. In *Bidang MIPA dan Sains* (Issue 5).
- Rosidah, I., Bahua, H., Mufidah, R., & Pongtuluran, O. B. (2015). Pengaruh Kondisi Proses Ekstraksi Batang Brotowali (*Tinospora crispa* (L) Hook.f & Thomson) Terhadap Aktivitas Hambatan Enzim Alfa Glukosidase. *Media Penelitian Dan Pengembangan Kesehatan*, 25(4). <https://doi.org/10.22435/mpk.v25i4.4586.203-210>
- Sanchez-Reinoso, Z., Mora-Adames, W. I., Fuenmayor, C. A., Darghan-Contreras, A. E., Gardana, C., & Gutiérrez, L. F. (2020). Microwave-assisted extraction of phenolic compounds from Sacha Inchi shell: Optimization, physicochemical properties and evaluation of their antioxidant activity. *Chemical Engineering and Processing - Process Intensification*, 153(May), 107922. <https://doi.org/10.1016/j.cep.2020.107922>
- Sani, M. S. A., Bakar, J., Azid, A., & Iqbal, M. J. (2022). Chemometrics-based evaluation on the effect of sonication, contact time and solid-to-solvent ratio on total phenolics and flavonoids, free fatty acids and antibacterial potency of Carica papaya seed against *S. enteritidis*, *B. cereus*, *V. vulnificus* and *P. . Food Chemistry Advances*, 1(February), 100033. <https://doi.org/10.1016/j.focha.2022.100033>
- Sant'Anna, V., Brandelli, A., Marczak, L. D. F., & Tessaro, I. C. (2012). Kinetic modeling of total polyphenol extraction from grape marc and characterization of the extracts. *Separation and Purification Technology*, 100, 82–87. <https://doi.org/10.1016/j.seppur.2012.09.004>

- Serea, D., Constantin, O. E., Horincar, G., Stănciuc, N., Aprodu, I., Bahrim, G. E., & Răpeanu, G. (2023). Optimization of Extraction Parameters of Anthocyanin Compounds and Antioxidant Properties from Red Grape (*Băbească neagră*) Peels. *Inventions*, 8(2). <https://doi.org/10.3390/inventions8020059>
- Setyaningsih, W., Putro, A. W., Fathimah, R. N., Kurnia, K. A., Darmawan, N., Yulianto, B., Jiwanti, P. K., Carrera, C. A., & Palma, M. (2022). A microwave-based extraction method for the determination of sugar and polyols: Application to the characterization of regular and peaberry coffees. *Arabian Journal of Chemistry*, 15(3), 103660. <https://doi.org/10.1016/j.arabjc.2021.103660>
- Sharma, M., & Dash, K. K. (2022). Microwave and ultrasound assisted extraction of phytochemicals from black jamun pulp: Kinetic and thermodynamics characteristics. *Innovative Food Science and Emerging Technologies*, 75(December 2021), 102913. <https://doi.org/10.1016/j.ifset.2021.102913>
- Shen, Y., Jin, L., Xiao, P., Lu, Y., & Bao, J. (2009). Total phenolics, flavonoids, antioxidant capacity in rice grain and their relations to grain color, size and weight. *Journal of Cereal Science*, 49(1), 106–111. <https://doi.org/10.1016/j.jcs.2008.07.010>
- Sholihah, A., Aini, N., & Dwiyanti, H. (2021). Optimasi Ekstraksi Antosianin Pada Beras Hitam Sirampog Menggunakan Metode Ultrasound Extract Assist (UAE). *Jurnal Aplikasi Teknologi Pangan*, 10(3), 71–76.
- Sinaga, A. S. (2019). SEGMENTASI RUANG WARNA L*a*b. *Jurnal Mantik Penusa*, 3(1), 43–46.
- Škerget, M., Kotnik, P., Hadolin, M., Hraš, A. R., Simonič, M., & Knez, Ž. (2005). Phenols, proanthocyanidins, flavones and flavonols in some plant materials and their antioxidant activities. *Food Chemistry*, 89(2), 191–198. <https://doi.org/10.1016/j.foodchem.2004.02.025>
- Sompong, R., Siebenhandl-Ehn, S., Linsberger-Martin, G., & Berghofer, E. (2011). Physicochemical and antioxidative properties of red and black rice varieties from Thailand, China and Sri Lanka. *Food Chemistry*, 124(1), 132–140. <https://doi.org/10.1016/j.foodchem.2010.05.115>
- Sonar, M. P., & Rathod, V. K. (2020). Microwave assisted extraction (MAE) used as a tool for rapid extraction of Marmelosin from *Aegle marmelos* and evaluations of total phenolic and flavonoids content, antioxidant and anti-inflammatory activity. *Chemical Data Collections*, 30, 100545. <https://doi.org/10.1016/j.cdc.2020.100545>
- Stănciuc, N., Ureșan, C. M., & Bahrim, G. E. (2020). RECOVERY OF BIOACTIVE COMPOUNDS FROM RED ONION SKINS USING CONVENTIONAL SOLVENT EXTRACTION AND MICROWAVE ASSISTED EXTRACTION. *The Annals of the University Dunarea de Jos of Galati Fascicle VI – Food Technology*, 44, 104–126.
- Supardan, M. D., Asnawi, T. M., Putri, Y., & Wahyuni, S. (2011). Application of Ultrasound-assisted Solvent Extraction for Recovery of Oil from Palm Oil Mill Effluent. *Agritech*, 31(4), 368–373.
- Susanti, D. Y., Sediawan, W. B., Fahrurrozi, M., & Hidayat, M. (2020). Optimization of Agitation and Kinetic Studies on Proanthocyanidin

- Compound Extraction from Red Sorghum Grains in Agitated Vessel. *IOP Conference Series: Materials Science and Engineering*, 778(1). <https://doi.org/10.1088/1757-899X/778/1/012085>
- Tensiska, Sukarminah, E., & Natalia, D. (2007). EKSTRAKSI PEWARNA ALAMI DARI BUAH ARBEN (*Rubus idaeus* Linn.) DAN APLIKASINYA PADA SISTEM PANGAN. *Jurnal. Teknol. Dan Industri Pangan*, XVIII(1).
- Tomasi, I. T., Santos, S. C. R., Boaventura, R. A. R., & Botelho, C. M. S. (2023). Optimization of microwave-assisted extraction of phenolic compounds from chestnut processing waste using response surface methodology. *Journal of Cleaner Production*, 395(July 2022). <https://doi.org/10.1016/j.jclepro.2023.136452>
- Upadhyay, R., Ramalakshmi, K., & Jagan Mohan Rao, L. (2012). Microwave-assisted extraction of chlorogenic acids from green coffee beans. *Food Chemistry*, 130(1), 184–188. <https://doi.org/10.1016/j.foodchem.2011.06.057>
- Utasee, S., Jamjod, S., Lordkaew, S., & Prom-U-Thai, C. (2022). Improve Anthocyanin and Zinc Concentration in Purple Rice by Nitrogen and Zinc Fertilizer Application. *Rice Science*, 29(5), 435–450. <https://doi.org/10.1016/j.rsci.2022.07.004>
- Vermerris, W., & Nicholson, R. (2006). PHENOLIC COMPOUND BIOCHEMISTRY. In *Estudios Atacamenos* (Issue 63). Springer. <https://doi.org/10.22199/issn.0718-1043-2019-0023>
- Wang, G., Lu, M., Zhang, S., Ji, J., Li, B., Li, J., Zhang, L., Yang, D., Wang, W., & Guan, C. (2022). Anthocyanin release and absorption properties of boiling pigmented rice using an in vitro digestion model. *Journal of Food Measurement and Characterization*, 16(4), 2649–2663. <https://doi.org/10.1007/s11694-022-01378-2>
- Wang, X., Xu, J., Wang, L., Gao, X., Fu, X., & Zhao, Y. (2019). Optimization of Microwave-Ultrasound-assisted Enzymatic Hydrolysis extraction of iodine amino acids in laminaria by high performance liquid chromatography with a photodiode array detector. *Algal Research*, 39(July 2018), 101452. <https://doi.org/10.1016/j.algal.2019.101452>
- Wei, C., Wan, C., Huang, F., & Guo, T. (2023). Extraction of *Cinnamomum longepaniculatum* deciduous leaves essential oil using solvent-free microwave extraction: Process optimization and quality evaluation. *Oil Crop Science*, 8(1), 7–15. <https://doi.org/10.1016/j.ocsci.2023.02.004>
- Wiyono, T., & Kartikawati, D. (2017). Pengaruh Metode Ekstraksi Sari Nanas secara Langsung dan Osmosis Dengan Variasi Perebusan Terhadap Kualitas Sirup Nanas (*Ananas comosus* L.). *Jurnal Ilmiah UNTAG Semarang. [Diakses 2020 Des 19];*, 6(2), 108–118.
- Wu, X., Beecher, G. R., Holden, J. M., Haytowitz, D. B., Gebhardt, S. E., & Prior, R. L. (2006). Concentrations of anthocyanins in common foods in the United States and estimation of normal consumption. *Journal of Agricultural and Food Chemistry*, 54(11), 4069–4075. <https://doi.org/10.1021/jf0603001>
- Xiaokang, W., Lyng, J. G., Brunton, N. P., Cody, L., Jacquier, J. C., Harrison, S. M., & Papoutsis, K. (2020). Monitoring the effect of different microwave extraction parameters on the recovery of polyphenols from shiitake

- mushrooms: Comparison with hot-water and organic-solvent extractions. *Biotechnology Reports*, 27, e00504. <https://doi.org/10.1016/j.btre.2020.e00504>
- Xiong, G., Jia, L., Luo, L., Ding, Y., Lin, Q., & Liu, C. (2023). Improvement in texture and cooking quality of black rice (*Oryza sativa L.*) using different pretreatments. *Journal of Cereal Science*, 109(December 2022), 103611. <https://doi.org/10.1016/j.jcs.2022.103611>
- Yi, J., Qiu, M., Zhu, Z., Dong, X., Andrew Decker, E., & McClements, D. J. (2021). Robust and recyclable magnetic nanobiocatalysts for extraction of anthocyanin from black rice. *Food Chemistry*, 364(March), 130447. <https://doi.org/10.1016/j.foodchem.2021.130447>
- Zhou, Z., Chen, X., Zhang, M., & Blanchard, C. (2014). Phenolics, flavonoids, proanthocyanidin and antioxidant activity of brown rice with different pericarp colors following storage. *Journal of Stored Products Research*, 59, 120–125. <https://doi.org/10.1016/j.jspr.2014.06.009>
- Ziegler, V., Ferreira, C. D., Hoffmann, J. F., Chaves, F. C., Vanier, N. L., de Oliveira, M., & Elias, M. C. (2018). Cooking quality properties and free and bound phenolics content of brown, black, and red rice grains stored at different temperatures for six months. *Food Chemistry*, 242(March 2017), 427–434. <https://doi.org/10.1016/j.foodchem.2017.09.077>