



DAFTAR PUSTAKA

- Ahima, R. S. (2011). Digging deeper into obesity. *The Journal of Clinical Investigation*, 121(6), 2078.
- Ahima, R. S., & Lazar, M. A. (2013). The health risk of obesity - Better metrics imperative. *Science*, 341(6148), 856–858.
<https://doi.org/10.1126/science.1241244>
- Apridamayanti, P., Pratiwi, R., Purwestri, Y. A., Tunjung, W. A. S., & Rumiyati, R. (2017). Anthocyanin, nutrient contents, and antioxidant activity of black rice bran of *Oryza sativa* L.'Cempo Ireng' from Sleman, Yogyakarta, Indonesia. *Indonesian Journal of Biotechnology*, 22(1), 49-54.
- Aulbach, A. D., & Amuzie, C. J. (2017). Biomarkers in nonclinical drug development. In *A Comprehensive guide to toxicology in nonclinical drug development* (pp. 447-471). Academic Press.
- Auliya, P., Oenzil, F., & Dia Rofinda, Z. D. (2016). Gambaran kadar gula darah pada mahasiswa fakultas kedokteran Universitas Andalas yang memiliki berat badan berlebih dan obesitas. *Jurnal Kesehatan Andalas*, 5(3), 528–533.
<https://doi.org/10.25077/jka.v5i3.571>
- Banerjee, P. & Ray, D. P. (2019). Functional food: A brief overview. *International Journal of Bioresource Science*, 6(2), 57–60. <https://doi.org/10.30954/2347-9655.02.2019.2>
- Banjerdpongchai, R., Wudtiwai, B., & Sringsarm, K. (2013). Cytotoxic and apoptotic-inducing effects of purple rice extracts and chemotherapeutic drugs on human cancer cell lines. *Asian Pacific Journal of Cancer Prevention*, 14(11), 6541-6548.
- Barham D, Trinder P. An improved color reagent for the determination of blood glucose by the oxidase system. Analyst. 1972;97:142-5.

- Barritt, A. S., Watkins, S., Gitlin, N., Klein, S., Lok, A. S., Loomba, R., ... & Sanyal, A. J. (2021). Patient determinants for histologic diagnosis of NAFLD in the real world: a TARGET-NASH study. *Hepatology Communications*, 5(6), 938-946.
- Belwal, T., Nabavi, S. F., Nabavi, S. M., & Habtemariam, S. (2017). Dietary anthocyanins and insulin resistance: When food becomes a medicine. *Nutrients*, 9(10), 1111.
- BPOM. 2011. Peraturan Kepala Badan Pengawas Obat dan Makanan Republik Indonesia tentang Pengawasan Klaim Dalam Label dan Iklan Pangan Olahan No. HK.03.1.23.11.11.09909. Jakarta.
- Carpentier, A. C. (2021). 100th anniversary of the discovery of insulin perspective: insulin and adipose tissue fatty acid metabolism. *American Journal of Physiology-Endocrinology and Metabolism*, 320(4), E653-E670.
- Chasanah, S. N. & Pratiwi, R. (2019). Kadar Serum Glutamic Pyruvate Transaminase Darah Tikus (*Rattus norvegicus*, Berkenhout, 1769) Hiperlipidemia dengan asupan pelet nasi dan bekatul beras hitam padi (*Oryza sativa L.*) "Cempo Ireng". *Jurnal Ilmiah Cendekia Eksakta*, 4(1), 33-37.
- Daeli, E., & Ardiaria, M. (2018). Pengaruh Pemberian Nasi Beras Merah (*Oryza nivara*) dan Nasi Beras Hitam (*Oryza sativa L. indica*) terhadap Perubahan Kadar Gula Darah dan Trigliserida Tikus Wistar (*Rattus norvegicus*) Diabetes Melitus Tipe 2. *JNH (Journal of Nutrition and Health)*, 6(2), 42-56.
- Divella, R., Mazzocca, A., Daniele, A., Sabbà, C., & Paradiso, A. (2019). Obesity, nonalcoholic fatty liver disease and adipocytokines network in promotion of cancer. *International journal of biological sciences*, 15(3), 610.
- DPKP DIY. (2022, Oktober 27). *Sembada Hitam Sembodo Rupo Sembodo Roso*. <https://dpkp.jogjaprov.go.id/baca/Sembada+Hitam+Sembodo+Rupo+Sembo do+Roso/271022/8fa8883bd5df388296038be7fefb04178139b4466d0940f0f0 4d42dba489fed3563>



Fajrian, F. M. 2020. Enzim transferase dengan bilirubin total penderita ikterus obstruktif. *Jurnal Ilmiah Kesehatan Sandi Husada*, 11(1), 178.

Geisler, C. E., & Renquist, B. J. (2017). Hepatic lipid accumulation: cause and consequence of dysregulated glucoregulatory hormones. *Journal of Endocrinology*, 234(1), R1-R21.

Granato, D., Barba, F. J., Lorenzo, J. M., Cruz, A. G., & Putnik, P. (2019). *Functional Foods: Product Development, Technological Trends, Efficacy Testing, and Safety*. 1–26.

Granato, D., Nunes, D. S., & Barba, F. J. (2017). An integrated strategy between food chemistry, biology, nutrition, pharmacology, and statistics in the development of functional foods: A proposal. *Trends in Food Science and Technology*, 62(December 2016), 13–22.
<https://doi.org/10.1016/j.tifs.2016.12.010>

Guo, H., Ling, W., Wang, Q., Liu, C., Hu, Y., Xia, M., ... Xia, X. (2007). Effect of anthocyanin-rich extract from black rice (*Oryza sativa L. indica*) on hyperlipidemia and insulin resistance in fructose-fed rats. *Plant Foods for Human Nutrition*, 62(1), 1–6. <https://doi.org/10.1007/s11130-006-0031-7>

Gusdon, A. M., Song, K. X., & Qu, S. (2014). Nonalcoholic fatty liver disease: pathogenesis and therapeutics from a mitochondria-centric perspective. *Oxidative medicine and cellular longevity*, 2014. <http://dx.doi.org/10.1155/2014/637027>

Hadizadeh, F., Faghihimani, E., & Adibi, P. (2017). Nonalcoholic fatty liver disease: Diagnostic biomarkers. *World journal of gastrointestinal pathophysiology*, 8(2), 11.

Hardy, O. T., Czech, M. P., & Corvera, S. (2012). What causes the insulin resistance underlying obesity?. *Current opinion in endocrinology, diabetes, and obesity*, 19(2), 81.

Hioki, C., Yoshida, T., Kogure, A., Yoshimoto, K., & Shimatsu, A. (2010). Growth

hormone administration controls body composition associated with changes of thermogenesis in obese KK-Ay Mice. *The Open Endocrinology Journal*, 4(1).

Ilmi, W., Pratiwi, R., & Purwestri, Y. A. (2018). Total anthocyanin content and antioxidant activity of Brown rice, endosperm, and rice bran of three Indonesian Black rice (*Oryza sativa L.*) cultivars. In *Proceeding of the 2nd International Conference on Tropical Agriculture* (pp. 205-216). Springer International Publishing.

Indrisari, M., Nurkhairi, N., & Herdayanti, R. (2015). Uji Aktivitas Ekstrak Beras Hitam (*Oryza sativa L.*) Terhadap Penurunan Berat Badan Tikus (*Rattus norvegicus*). *Jurnal Farmasi dan Bahan Alam: FARBAL*, 3(2), 38-40.

Jendrassik, L. (1938). Vereinfachte photometrische methoden zur bestimmung des blutbilirubins. *Biochem z*, 297, 81-89.

Kahar, H. (2017). Pengaruh hemolisis terdapat kadar serum *glutamate pyruvate transaminase* (SGPT) sebagai salah satu parameter fungsi hati. *The Journal of Muhammadiyah Medical Laboratory Technologist*, 1(1), 38. <https://doi.org/10.30651/jmlt.v1i1.981>

Kementerian Kesehatan Republik Indonesia. 2014. Laporan Riset Kesehatan Dasar 2013. http://labdata.litbang.kemkes.go.id/images/download/laporan/RKD/2013/Laporan_riskesdas_2013_final.pdf. Diakses tanggal 12 Maret 2022, jam 09.25.

Kementerian Kesehatan Republik Indonesia. 2019. Laporan Riset Kesehatan Dasar 2018. http://labdata.litbang.kemkes.go.id/images/download/laporan/RKD/2018/Laporan_Nasional_RKD2018_FINAL.pdf. Diakses tanggal 12 Maret 2022, jam 10.05.

Koek, G. H., Liedorp, P. R., & Bast, A. (2011). The role of oxidative stress in non-alcoholic steatohepatitis. *Clinica chimica acta*, 412(15-16), 1297-1305.

Kurdanti, W., Suryani, I., Syamsiatun, N. H., Siwi, L. P., Adityanti, M. M., Mustikaningsih, D., & Sholihah, K. I. (2015). Faktor-faktor yang

mempengaruhi kejadian obesitas pada remaja. *Jurnal Gizi Klinik Indonesia*, 11(4), 179. <https://doi.org/10.22146/ijcn.22900>

Kwak, M. S., Kim, D., Chung, G. E., Kang, S. J., Park, M. J., Kim, Y. J., Yoon, J. H., & Lee, H. S. (2012). Serum bilirubin levels are inversely associated with nonalcoholic fatty liver disease. *Clinical and molecular hepatology*, 18(4), 383.

Lee, S. I., Kim, J. W., Lee, Y. K., Yang, S. H., Lee, I., Suh, J. W., & Kim, S. D. (2011). Anti-obesity effect of monascus pilosus mycelial extract in high fat diet-induced obese rats. *Journal of Applied Biological Chemistry*, 54(3), 197-205.

Ma, X., Liu, S., Zhang, J., Dong, M., Wang, Y., Wang, M., & Xin, Y. (2020). Proportion of NAFLD patients with normal ALT value in overall NAFLD patients: a systematic review and meta-analysis. *BMC gastroenterology*, 20, 1-8.

Marques, C., Meireles, M., Norberto, S., Leite, J., Freitas, J., Pestana, D., ... & Calhau, C. (2016). High-fat diet-induced obesity Rat model: a comparison between Wistar and Sprague-Dawley Rat. *Adipocyte*, 5(1), 11-21.

Mastuti, T., S., T. S. M., Matita, I. C., Pokatong, W. D. R., & Neysha, A. (2020). Penyuluhan peran pangan fungsional dan pola hidup sehat untuk menghadapi pandemi corona. *Prosiding Konferensi Nasional Pengabdian Kepada Masyarakat Dan Corporate Social Responsibility (PKM-CSR)*, 3, 428–432. <https://doi.org/10.37695/pkmcsr.v3i0.972>

McMurray, F., Patten, D. A., & Harper, M. E. (2016). Reactive oxygen species and oxidative stress in obesity—recent findings and empirical approaches. *Obesity*, 24(11), 2301-2310.

Mohan, S., Thiagarajan, K., Sundaramoorthy, B., Gurung, V., Barpande, M., Agarwal, S., & Chandrasekaran, R. (2016). Alleviation of 4-nitroquinoline 1-oxide induced oxidative stress by Oroxyllum indicum (L.) leaf extract in albino Wistar rats. *BMC complementary and alternative medicine*, 16, 1-11.



Nashrurrokhman, M., Sayekti, P. R., Safitri, A., Purwestri, Y. A., & Pratiwi, R.

(2019). Macronutrient and mineral contents of five local black rice (*Oryza sativa*) cultivars in Indonesia. *Biodiversitas Journal of Biological Diversity*, 20(12).

Nastiti, R. D. W., Nurhidajah, N., & Yusuf, M. (2020). Berat Badan, Feed Conversion Ratio (Fcr), Dan Berat Jaringan Adiposa Pada Tikus Hiperkolesterolemia Dengan Diet Beras Hitam. *Jurnal Pangan dan Gizi*, 10(2), 147-158.

Novelli, E. L. B., Diniz, Y. S., Galhardi, C. M., Ebaid, G. M. X., Rodrigues, H. G., Mani, F., ... Novelli Filho, J. L. V. B. (2007). Anthropometrical parameters and markers of obesity in rats. *Laboratory Animals*, 41(1), 111–119. <https://doi.org/10.1258/002367707779399518>

Nowicka, P., Wojdyło, A., & Samoticha, J. (2016). Evaluation of phytochemicals, antioxidant capacity, and antidiabetic activity of novel smoothies from selected *Prunus* fruits. *Journal of Functional Foods*, 25, 397-407.

Nurpalah, R., Kusmiati, M., & Intan Saproh, J. (2018). Gambaran kadar sgpt pada obesitas sentral. *Prosiding Seminar Nasional Dan Diseminasi Penelitian Kesehatan STIKes Bakti Tunas Husada Tasikmalaya, 21 April 2018 ISBN:978-602-72636-3-*, (April), 168–171.

Octaviany, V. D., Yusmaini, H., & Simanjuntak, K. (2017). Uji efektivitas ekstrak kulit apel (*Malus sylvestris*-mill) var. rome beauty terhadap kadar enzim SGPT tikus (*Rattus norvegicus*) galur Wistar yang diinduksi CCl4 (karbon tetraklorida). *Jurnal Profesi Medika*, 11(2), 56-62.

Oteiza, P. I., Cremonini, E., & Fraga, C. G. (2022). Anthocyanin actions at the gastrointestinal tract: Relevance to their health benefits. *Molecular Aspects of Medicine*, 101156.

Pawar, S. V., Zanwar, V. G., Choksey, A. S., Mohite, A. R., Jain, S. S., Surude, R. G., ... & Varthakavi, P. K. (2017). Most overweight and obese Indian children have nonalcoholic fatty liver disease. *Annals of hepatology*, 15(6), 853-861.

Panjaitan. R. G. P., Handharyani, E., Chairul, Masriani, Zakiah, Z., & Manalu, W. (2007). Pengaruh pemberian karbon tetraklorida terhadap fungsi hati dan ginjal tikus. *Makara Kesehatan*, 11(1), 13.

Pengkumsri, N., Chaiyasut, C., Saenjum, C., Sirilun, S., Peerajan, S., Suwannaert, P., ... Sivamaruthi, B. S. (2015). Physicochemical and antioxidative properties of black, brown and red rice varieties of northern Thailand. *Food Science and Technology (Brazil)*, 35(2), 331–338. <https://doi.org/10.1590/1678-457X.6573>

Poonia, A., & Pandey, S. (2021). Bioactive compounds, nutritional benefits and food applications of black rice: a review. *Nutrition & Food Science*, 52(3), 466-482.

Pratiwi, R., & Purwestri, Y. A. (2017). Black rice as a functional food in Indonesia. *Functional Foods in Health and Disease*, 7(3), 182–194. <https://doi.org/10.31989/ffhd.v7i3.310>

Pratiwi, R., Tunjung, W. A. S., Rumiyati, R., & Amalia, A. R. (2016). Black rice bran extracts and fractions containing Cyanidin 3-glucoside and Peonidin 3-glucoside induce apoptosis in human cervical cancer Cells. *Indonesian Journal of Biotechnology*, 20(1), 69. <https://doi.org/10.22146/ijbiotech.15271>

Preguiça, I., Alves, A., Nunes, S., Fernandes, R., Gomes, P., Viana, S. D., & Reis, F. (2020). Diet-induced rodent models of obesity-related metabolic disorders—a guide to a translational perspective. *Obesity Reviews*, 21(12), e13081.

Prihapsara, F., Artanti, A. N., & Harini, M. (2017). Acute and subchronic toxicity of Self Nanoemulsifying Drug Delivery Systems (SNEDDS) from chloroform bay leaf extract (*Eugenia Polyantha* W.) with palm kernel oil as a carrier. In *IOP Conference Series: Materials Science and Engineering* (Vol. 333, No. 1, p. 012066). IOP Publishing.

Purwestri, Y. A., Pratiwi, R., Nuringtyas, T. Ri., Rumiyati, Fauzia, A. N., & Garusti. (2022). *Makanan Fungsional Berbahan Dasar Beras Hitam dan*

Proses Pembuatannya.

- Qi, S. S., He, J., Yuan, L. P., Le Wu, J., Zu, Y. X., & Zheng, H. X. (2020). Cyanidin-3-glucoside from black rice prevents renal dysfunction and renal fibrosis in streptozotocin-diabetic rats. *Journal of Functional Foods*, 72, 104062.
- Rabbani, F. (2018). *Perbedaan kadar SGPT berdasarkan frekuensi penggunaan kuvet*. Tesis. Fakultas Ilmu Keperawatan dan Kesehatan Unibersitas Muhammadiyah Semarang. Semarang.
- Rahayu, M. S., & Wahyuni, S. (2021). Effects of Oral Administration of Monosodium Glutamate (MSG) on Obesity in Male Wistar Rats (*Rattus Norvegicus*). *Bioscientia Medicina: Journal of Biomedicine and Translational Research*, 5(9), 879-882.
- Rosida, A. (2016). Pemerikasaan laboratorium penyakit hati. *Berkala Kedokteran*, 12(1), 123-131.
- Ruiz, A. R. G., Crespo, J., López Martínez, R. M., Iruzubieta, P., Casals Mercadal, G., Lalana Garcés, M., ... & Morales Ruiz, M. (2021). Measurement and clinical usefulness of bilirubin in liver disease. *Advances in Laboratory Medicine/Avances en Medicina de Laboratorio*, 2(3), 352-361.
- Samyor, D., Das, A. B., & Deka, S. C. (2017). Pigmented rice a potential source of bioactive compounds: a review. *International Journal of Food Science and Technology*, 52(5), 1073–1081. <https://doi.org/10.1111/ijfs.13378>
- Sancho, R. A. S., & Pastore, G. M. (2012). Evaluation of the effects of anthocyanins in type 2 diabetes. *Food Research International*, 46(1), 378-386.
- Saxena, T., Arya, A., Rathore, A. J., Rajak, N., Naz, S., & Shah, R. (2014). GGT and SGPT-A rising marker in diagnosis of non-alcoholic fatty liver disease. *Biomedical & Pharmacology Journal*, 7(1), 277.
- Serra, D., Mera, P., Malandrino, M. I., Mir, J. F., & Herrero, L. (2013). Mitochondrial fatty acid oxidation in obesity. *Antioxidants & redox signaling*, 19(3), 269-284.



Shiyan S, Herlina BM, Amriani A. Antiobesitas dan antihipercolesterolemia seduhan white tea (*Camellia sinensis*) pada tikus yang diberi diet lemak tinggi. *Pharmaciana* 2017; 7(2): 280.

Snel, M., Jonker, J. T., Schoones, J., Lamb, H., de Roos, A., Pijl, H., ... & Jazet, I. M. (2012). Ectopic fat and insulin resistance: pathophysiology and effect of diet and lifestyle interventions. *International journal of endocrinology*, 2012.

Song, H., Shen, X., Zhou, Y., & Zheng, X. (2021). Black rice anthocyanins alleviate hyperlipidemia, liver steatosis and insulin resistance by regulating lipid metabolism and gut microbiota in obese mice. *Food & Function*, 12(20), 10160-10170.

Stephens, D. N. (1980). Does the Lee obesity index measure general obesity?. *Physiology & Behavior*, 25(2), 313-315.

Suter, I. K. (2013). Pangan fungsional dan prospek pengembangannya. *Pangan Fungsional Dan Prospek Pengembangannya*, 1, 17.

Te Sligte, K., Bourass, I., Sels, J. P., Driessen, A., Stockbrügger, R. W., & Koek, G. H. (2004). Non-alcoholic steatohepatitis: review of a growing medical problem. *European journal of internal medicine*, 15(1), 10-21.

Thanuja, B., & Parimalavalli, R. (2018). Role of black rice in health and diseases. *Int. J. Health Sci. Res*, 8, 241-248.

Tsalissavrina, I., Wahono, D., & Handayani, D. (2006). Pengaruh pemberian diet tinggi karbohidrat dibandingkan diet tinggi lemak terhadap kadar trigliserida dan HDL darah pada *Rattus norvegicus* galur wistar. *Jurnal Kedokteran Brawijaya*, 22(2), 80-89.

Utzschneider, K. M., & Kahn, S. E. (2006). The role of insulin resistance in nonalcoholic fatty liver disease. *The Journal of Clinical Endocrinology & Metabolism*, 91(12), 4753-4761.

Wanti, S., Andriani, M. A. ., & Herriyadi, N. (2015). Pengaruh berbagai jenis beras



terhadap aktivitas antimikrobia pada angkak oleh *Monascus purpureus*. *Biofarmasi*, 13(1), 1–5. <https://doi.org/10.13057/biofar/f130101>

Wen, J. J., Gao, H., Hu, J. L., Nie, Q. X., Chen, H. H., Xiong, T., ... & Xie, M. Y. (2019). Polysaccharides from fermented *Momordica charantia* ameliorate obesity in high-fat induced obese rats. *Food & function*, 10(1), 448-457.

WHO. Regional Office for the Western Pacific. (2000). The Asia-Pacific perspective : redefining obesity and its treatment. Sydney : Health Communications Australia. <https://apps.who.int/iris/handle/10665/206936>

Widyaningsih, D., Wijayanti, N., & Nugrahini, N. I. P. (2017). *Pangan Fungsional: Aspek Kesehatan, Evaluasi, dan Regulasi*. Universitas Brawijaya Press. Malang. Hal: 6-7.

Widyastuti, S., & Suarsana, I. N. (2011). Ekstrak air tapak data menurunkan kadar gula dan meningkatkan jumlah sel beta pankreas kelinci hiperglykemia. *Jurnal Veteriner*, 12(1): 7-12.

Wright, S. M., & Aronne, L. J. (2012). Causes of obesity. *Abdominal Imaging*, 37(5), 730–732. <https://doi.org/10.1007/s00261-012-9862-x>

Wu, T., Guo, X., Zhang, M., Yang, L., Liu, R., & Yin, J. (2017). Anthocyanins in black rice, soybean and purple corn increase fecal butyric acid and prevent liver inflammation in high fat diet-induced obese mice. *Food and Function*, 8(9), 3178–3186. <https://doi.org/10.1039/c7fo00449d>

Xu, Y., Zhang, M., Wu, T., Dai, S., Xu, J., & Zhou, Z. (2015). The anti-obesity effect of green tea polysaccharides, polyphenols and caffeine in rats fed with a high-fat diet. *Food & Function*, 6(1), 296-303.

Yao, S. L., Xu, Y., Zhang, Y. Y., & Lu, Y. H. (2013). Black rice and anthocyanins induce inhibition of cholesterol absorption in vitro. *Food and Function*, 4(11), 1602–1608. <https://doi.org/10.1039/c3fo60196j>

Zaupa, M., Calani, L., Del Rio, D., Brighenti, F., & Pellegrini, N. (2015). Characterization of total antioxidant capacity and (poly)phenolic compounds



UNIVERSITAS
GADJAH MADA

Pengaruh Pemberian Produk Olahan Pangan Fungsional Black Rice Crunch terhadap Indeks Lee,
Kadar Glukosa, SGPT, dan Bilirubin Darah Tikus Putih (*Rattus norvegicus* Berkenhout, 1769) Obesitas Baik Aisyah Adania, Dra. Rarastoeti Pratiwi, M.Sc., Ph.D.

Universitas Gadjah Mada, 2023 | Diunduh dari <http://etd.repository.ugm.ac.id/>

of differently pigmented rice varieties and their changes during domestic cooking. *Food Chemistry*, 187, 338–347.
<https://doi.org/10.1016/j.foodchem.2015.04.055>

Žiberna, L., Jenko-Pražnikar, Z., & Petelin, A. (2021). Serum bilirubin levels in overweight and obese individuals: The importance of anti-inflammatory and antioxidant responses. *Antioxidants*, 10(9).
<https://doi.org/10.3390/antiox10091352>