

## DAFTAR PUSTAKA

- [AOAC] Association of Official Analytical. (1990). *Official Methods of Analysis of AOAC International* (15<sup>th</sup> ed). AOAC, Virginia.
- Abdelmeguid, N. E., Fakhoury, R., Kamal, S. M., & Wafai, R. J. A. L. (2010). Effects of *Nigella sativa* and thymoquinone on biochemical and subcellular changes in pancreatic b -cells of streptozotocin-induced diabetic rats. *Journal of Diabetes* 2, 2, 256–266. <https://doi.org/10.1111/j.1753-0407.2010.00091.x>
- Adilah, N., Idris, N., & Sulaiman, A. Z. (2017). Comparison between conventional extraction and ultrasound assisted extraction of *Labisia pumila* sp . in 25-L mobile extractor using water as solvent of extraction. *Chemical Engineering Transactions*, 56, 781–786. <https://doi.org/10.3303/CET1756131>
- Aguilar-Rosas, S. F., Ballinas-Casarrubias, M. L., Nevarez-Moorillon, G. V., Martin-Belloso, O., & Ortega-Rivas, E. (2007). Thermal and pulsed electric fields pasteurization of apple juice: Effects on physicochemical properties and flavour compounds. *Journal of Food Engineering*, 83(1), 41–46. <https://doi.org/10.1016/j.jfoodeng.2006.12.011>
- Ahmad, A., Husain, A., Mujeeb, M., Khan, S. A., Najmi, A. K., Siddique, N. A., Damanhour, Z. A., & Anwar, F. (2013). A review on therapeutic potential of *Nigella sativa*: A miracle herb. *Asian Pacific Journal of Tropical Biomedicine*, 3(5), 337–352. [https://doi.org/10.1016/S2221-1691\(13\)60075-1](https://doi.org/10.1016/S2221-1691(13)60075-1)
- Alfonso, A. A., & Mongan, A. E. (2016). Gambaran kadar kreatinin serum pada pasien penyakit ginjal kronik stadium 5 non dialisis. *Jurnal E-Biomedik*, 4(1), 178–183.
- Ali, H., Yesmin, R., Satter, M. A., Habib, R., & Yeasmin, T. (2018). Antioxidant and antineoplastic activities of methanolic extract of *Kaempferia galanga* Linn. Rhizome against Ehrlich ascites carcinoma cells. *Journal of King Saud University - Science*, 30(3), 386–392. <https://doi.org/10.1016/j.jksus.2017.05.009>
- Almas, I., Innocent, E., Machumi, F., & Kisinza, W. (2021). Chemical composition of essential oils from *Eucalyptus globulus* and *Eucalyptus maculata* grown in Tanzania. *Scientific African*, 12, e00758. <https://doi.org/10.1016/j.sciaf.2021.e00758>
- Aloud, A. A., Veeramani, C., Govindasamy, C., Alsaif, M. A., El Newehy, A. S., & Al-Numair, K. S. (2017). Galangin, a dietary flavonoid, improves antioxidant status and reduces hyperglycemia-mediated oxidative stress in streptozotocin-induced diabetic rats. *Redox Report*, 22(6), 290–300.
- Alsahli, M. S., Elumalai, K., Devanesan, S., Govindarajan, M., Krishnappa, K., & Maggi, F. (2020). The aromatic ginger *Kaempferia galanga* L. (Zingiberaceae) essential oil and its main compounds are effective larvicidal agents against *Aedes vittatus* and *Anopheles maculatus* without toxicity on the non-target aquatic fauna. *Industrial Crops & Products*, 158(113012), 1–8. <https://doi.org/10.1016/j.indcrop.2020.113012>
- Álvarez, C. (2016). Identification of the volatile compounds in the roasting Venezuela Criollo cocoa beans by Gas Chromatography-Spectrometry Mass.

- Journal of Nutritional Health & Food Engineering*, 5(4), 659–666.  
<https://doi.org/10.15406/jnhfe.2016.05.00178>
- Angkasa, Dudung & Sulaeman, A. (2012). *Pengembangan minuman fungsional sumber serat dan antioksidan dari daun hantap (Sterculia oblongata R. Brown.)*. Institut Pertanian Bogor.
- Ani, T. A., Calinescu, I., & Lavric, V. (2009). *Ultrasonic vs microwave extraction intensification of active principles from medicinal plants*.  
<https://doi.org/10.3303/cet0917171>
- Aprotosoiaie, A. C., Luca, S. V., & Miron, A. (2016). Flavor chemistry of cocoa and cocoa products — An overview. *Comprehensive Reviews in Food Science and Food Safety*, 15, 73–91. <https://doi.org/10.1111/1541-4337.12180>
- Ariva, F. S. (2018). Penentuan kandungan antioksidan dalam rimpang Kencur (*Kaempferia galanga* L.) yang diekstraksi dengan bantuan gelombang ultrasonik. *Chempublish Journal*, 3(2), 64–74.  
<https://doi.org/10.22437/chp.v3i2.5769>
- Arivazhagan, P., Thilakavathy, T., & Panneerselvam, C. (2000). Antioxidant lipoate and tissue antioxidants in aged rats. *Journal of Nutritional Biochemistry*, 11(3), 122–127. [https://doi.org/10.1016/S0955-2863\(99\)00079-0](https://doi.org/10.1016/S0955-2863(99)00079-0)
- Ashari, H., Sadeghi, N., Oveisi, M. R., & Hajimahmoodi, M. (2020). Total phenol, total flavonoid and ascorbic acid content of Iranian commercial orange juice. *Journal of Food Safety and Hygiene*, 6(1), 22–27.
- Astuti, Y., Sundari, D., Winarno, W. (1996). *Tanaman kencur (Kaempferia galanga. Linn) informasi tentang fitokimia dan efek farmakologis*. Pusat Penelitian dan Pengembangan Farmasi.
- Ayala, A., Muñoz, M. F., & Argüelles, S. (2014). Lipid peroxidation: Production, metabolism, and signaling mechanisms of Malondialdehyde and 4-Hydroxy-2-Nonenal. *Oxidative Medicine and Cellular Longevity*, 1–31.  
<https://doi.org/10.1155/2014/360438>
- Az-Zahro, S. A. J., Umami, S. H., Hasanah, U., & Wijayanti, E. D. (2019). Aktivitas antihiperurisemia teh asam daun tin (*Ficus carica*) terhadap tikus putih (*Rattus norvegicus*). *Kartika : Jurnal Ilmiah Farmasi*, 7(1), 22.  
<https://doi.org/10.26874/kjif.v7i1.176>
- Balasundram, N., Sundram, K., & Samman, S. (2006). Phenolic compounds in plants and agri-industrial by-products : Antioxidant activity , occurrence , and potential uses. *Food Chemistry*, 99, 191–203.  
<https://doi.org/10.1016/j.foodchem.2005.07.042>
- Baynes, J. W. (1991). Perspectives in diabetes role of oxidative stress in development of complications in diabetes. *Diabetes*, 40, 405–412.
- Beauchamp, P. (2016). *Advanced organic spectroscopy tools for beginning organic spectroscopists: using simulated spectra to learn how to solve complicated organic structures*. Createspace Independent Publishing Platform.
- Belgis, M., Nafi, A., & Wulandari, A. D. (2021). Antibacterial activity of *Kaempferia galanga* L. hard candy against *Streptococcus pyogenes* and *Staphylococcus aureus* bacteria growth. *International Journal on Food, Agriculture, Adn Natural Resources*, 02(01), 1–8.

- Bellincampi, D., Cervone, F., & Lionetti, V. (2014). Plant cell wall dynamics and wall-related susceptibility in plant – pathogen interactions. *Frontiers in Plant Science*, 5(228), 1–8. <https://doi.org/10.3389/fpls.2014.00228>
- Benchaar, C., & Greathead, H. (2011). Essential oils and opportunities to mitigate enteric methane emissions from ruminants. *Animal Feed Science and Technology*, 166–167, 338–355. <https://doi.org/10.1016/j.anifeedsci.2011.04.024>
- Benzie, I. F. F., Szeto, Y. T., & Africa, N. (2000). Total antioxidant capacity of Teas by the ferric reducing / antioxidant power assay. *Journal Agriculture Food Chemistry*, 633–636.
- Berlett, B. S., Levine, R. L., Chock, P. B., Chevion, M., & Stadtman, E. R. (2001). Antioxidant activity of Ferrozine – iron – amino acid complexes. *Article in Proceedings of the National Academy of Science*, 98(2), 451–456. <https://doi.org/10.1073/pnas.021552298>
- Bou, M., Matar, J., Farhat, A., Abdelkhalek, S., Bou-maroun, E., & El, L. (2022). Antioxidant and anti-diabetic activities in commercial and homemade pomegranate molasses in Lebanon. *Food Bioscience*, 46, 101540. <https://doi.org/10.1016/j.fbio.2021.101540>
- Brodowska, K. M. (2017). Natural flavonoids : classification , potential role , and application of flavonoid analogues. *European Journal of Biological Research*, 7(2), 108–123.
- Bryant, S., Huerta, V., Mihalik, K., Crixell, S. H., & Vatter, D. A. (2008). Medicinal plants from Indian Subcontinent decrease quorum sensing dependent virulence in *Pseudomonas aeruginosa*. *Journal of Natural Remedies*, 8(2), 138–150.
- Cabré, M., Camps, J., Paternáin, J., Ferré, N., & Joven, J. (2000). Time-course of changes in hepatic lipid peroxidation and glutathione metabolism in rats with carbon tetrachloride-induced cirrhosis. *Clinical and Experimental Pharmacology and Physiology*, 27(9), 694–699. <https://doi.org/10.1046/j.1440-1681.2000.03322.x>
- Caillet, S., Salmiéri, S., & Lacroix, M. (2006). Evaluation of free radical-scavenging properties of commercial grape phenol extracts by a fast colorimetric method. *Food Chemistry*, 95(1), 1–8. <https://doi.org/10.1016/j.foodchem.2004.12.011>
- Calderón-Montaña, J. M., Burgos-Morón, E., Pérez-Guerrero, C., & López-Lázaro, M. (2011). A review on the dietary flavonoid Kaempferol. *Mini Reviews in Medicinal Chemistry*, 11(4), 298–344.
- Cameron, D. K., & Wang, Y. J. (2006). Application of protease and high-intensity ultrasound in corn starch isolation from degermed corn flour. *Cereal Chemistry*, 83(5), 505–509. <https://doi.org/10.1094/CC-83-0505>
- Caminiti, I. M., Noci, F., Muñoz, A., Whyte, P., Morgan, D. J., Cronin, D. A., & Lyng, J. G. (2011). Impact of selected combinations of non-thermal processing technologies on the quality of an apple and cranberry juice blend. *Food Chemistry*, 124(4), 1387–1392. <https://doi.org/10.1016/j.foodchem.2010.07.096>
- Canitez, N. (1998). *Pasteurization of Apple Cider with UV Irradiation*.

- Carabelly, A. N., Putri, D. K. T., Rezki, N., & Apriasari, M. L. (2020). SGOT and SGPT level of Wistar rat after the administration of *Channa micropeltes* extract micropeltes extract. *International Journal of Research in Pharmaceutical Sciences*, 11(4), 5415–5421. <https://doi.org/10.26452/ijrps.v11i4.3169>
- Chan, E. W. C., Lim, Y. Y., Wong, L. F., Lianto, F. S., Wong, S. K., Lim, K. K., Joe, C. E., & Lim, T. Y. (2008). Antioxidant and tyrosinase inhibition properties of leaves and rhizomes of ginger species. *Food Chemistry*, 109(3), 477–483. <https://doi.org/10.1016/j.foodchem.2008.02.016>
- Chang, H., Huang, G., Agrawal, D. C., Kuo, C., Wu, C., & Tsay, H. (2007). Antioxidant activities and polyphenol contents of six folk medicinal ferns used as “Gusuibu.” *Botanical Studies*, 48, 397–406.
- Chen, C. Z., & Cooper, S. L. (2002). Interactions between dendrimer biocides and bacterial membranes. *Biomaterials*, 23, 3359–3368.
- Chen, X., Wang, R., Hu, H., Zhao, X., Yin, Z., Zou, Y., Li, L., Jia, R., Zhang, Y., & Song, X. (2022). Antiviral effect of an extract from *Kaempferia galanga* L. rhizome in mice infected with pseudorabies virus. *Journal of Virological Methods*, 307(114573), 1–10. <https://doi.org/10.1016/j.jviromet.2022.114573>
- Chew, K. K., Ng, S. Y., Thoo, Y. Y., Khoo, M. Z., Wan Aida, W. M., & Ho, C. W. (2011). Effect of ethanol concentration, extraction time and extraction temperature on the recovery of phenolic compounds and antioxidant capacity of *Centella asiatica* extracts. *International Food Research Journal*, 18(2), 571–578.
- Chew, Y., Goh, J., & Lim, Y. (2009). Assessment of in vitro antioxidant capacity and polyphenolic composition of selected medicinal herbs from Leguminosae family in Peninsular Malaysia. *Food Chemistry*, 116(1), 13–18. <https://doi.org/10.1016/j.foodchem.2009.01.091>
- Cos, P., Ying, L., Calomme, M., Hu, J. P., Cimanga, K., Poel, B. Van, Pieters, L., Vlietinck, A. J., & Berghe, D. Vanden. (1998). Structure - Activity relationship and classification of flavonoids as inhibitors of xanthine oxidase and superoxide scavengers. *Journal of Natural Products*, 3864(32), 71–76. <https://doi.org/10.1021/np970237h>
- Coskun, O., Kanter, M., Korkmaz, A., & Oter, S. (2005). Quercetin, a flavonoid antioxidant, prevents and protects streptozotocin-induced oxidative stress and  $\beta$ -cell damage in rat pancreas. *Pharmacological Research*, 51(2), 117–123. <https://doi.org/10.1016/j.phrs.2004.06.002>
- Costa, M., Sezgin-bayindir, Z., Losada-barreiro, S., Paiva-martins, F., Saso, L., & Bravo-d, C. (2021). Polyphenols as Antioxidants for Extending Food Shelf-Life and in the Prevention of Health Diseases : Encapsulation and Interfacial Phenomena. *Biomedicines*, 9(1909), 1–38.
- Czaja, A. J. (2014). Hepatic inflammation and progressive liver fibrosis in chronic liver disease. *World Journal of Gastroenterology*, 20(10), 2515–2532. <https://doi.org/10.3748/wjg.v20.i10.2515>
- Dalle-donne, I., Rossi, R., Colombo, R., Giustarini, D., & Milzani, A. (2006). Biomarkers of oxidative damage in human disease. *Clinical Chemistry*, 52(4), 601–623. <https://doi.org/10.1373/clinchem.2005.061408>

- Damien Dorman, H. J., Deans, S. G., Noble, R. C., & Surai, P. (1995). Evaluation in vitro of plant essential oils as natural antioxidants. *Journal of Essential Oil Research*, 7(6), 645–651. <https://doi.org/10.1080/10412905.1995.9700520>
- Davis, W. W., & Stout, T. R. (1971). Disc plate method of microbiological antibiotic assay. II. Novel procedure offering improved accuracy. *Applied Microbiology*, 22(4), 666–670. <https://doi.org/10.1128/aem.22.4.666-670.1971>
- De Alencar, D. B., Da Silva, S. R., Pires-Cavalcante, K. M. S., De Lima, R. L., Pereira, F. N., De Sousa, M. B., Viana, F. A., Nagano, C. S., Do Nascimento, K. S., Cavada, B. S., Sampaio, A. H., & Saker-Sampaio, S. (2014). Antioxidant potential and cytotoxic activity of two red seaweed species, *amansia multifida* and *meristiella echinocarpa*, from the coast of Northeastern Brazil. *Anais Da Academia Brasileira de Ciencias*, 86(1), 251–263. <https://doi.org/10.1590/0001-37652014116312>
- De, P., Baltas, M., & Bedos-Belval, F. (2011). Cinnamic acid derivatives as anticancer agents - A review. *Current Medicinal Chemistry*, 18, 1672–1703.
- De, Prithwiraj, Veau, D., & Bedos-belval, F. (2005). Cinnamic derivatives in tuberculosis. In *Understanding tuberculosis - new approaches to fighting against drug resistance* (pp. 337–362).
- Decker, E. A. (1998). Strategies for manipulating the prooxidative / antioxidative balance of foods to maximize oxidative stability. *Trends in Food Science and Technology*, 9, 241–248.
- Dewi, I. G. A. M., Ganda Putra, G. P., & Wrsiati, L. P. (2021). Karakteristik ekstrak kulit biji kakao (*Theobroma cacao* L.) sebagai sumber antioksidan pada perlakuan suhu dan waktu maserasi. *Jurnal Rekayasa Dan Manajemen Agroindustri*, 8(1), 139–149. <https://doi.org/10.24843/jrma.2021.v09.i01.p01>
- Duarte, A., Luís, Â., Oleastro, M., & Domingues, F. C. (2015). Antioxidant properties of Coriander essential oil and Linalool and their potential to control *Campylobacter* spp. *Food Control*. <https://doi.org/10.1016/j.foodcont.2015.09.033>
- Duque, L., Restrepo, C., Sáez, J., Gil, J., Schneider, B., & Otálvaro, F. (2010). Synthesis of musafluorone: a naphthoxanthone isolated from *Musa acuminata*. *Tetrahedron Letters*, 51, 4640–4643. <https://doi.org/10.1016/j.tetlet.2010.06.128>
- Dwita, L. P., Hikmawanti, N. P. E., Yeni, & Supandi. (2021). Extract, fractions, and ethyl-p-methoxycinnamate isolate from *Kaempferia galanga* Elicit anti-inflammatory activity by limiting leukotriene B4 (LTB4) production. *Journal of Traditional and Complementary Medicine*, 11(6), 563–569. <https://doi.org/10.1016/j.jtcme.2021.06.004>
- El-haskoury, R., Al-waili, N., Kamoun, Z., Makni, M., & Al-waili, H. (2018). Antioxidant activity and protective effect of Carob Honey in CCl4 - induced kidney and liver injury. *Archives of Medical Research*, 49(5), 306–313. <https://doi.org/10.1016/j.arcmed.2018.09.011>
- El Jemli, M., Kamal, R., Marmouzi, I., Zerrouki, A., Cherrah, Y., & Alaoui, K. (2016). Radical-Scavenging activity and ferric reducing ability of *juniperus thurifera* (L.), *J. oxycedrus* (L.), *J. phoenicea* (L.) and *Tetraclinis articulata*



- (L.). *Advances in Pharmacological Sciences*, 1–6. <https://doi.org/10.1155/2016/6392656>
- Elmastaş, M., Gülçin, I., Işildak, Ö., Küfrevioğlu, Ö. I., Ibaoglu, K., & Aboul-Enein, H. Y. (2006). Radical scavenging activity and antioxidant capacity of bay leaf extracts. *Journal of the Iranian Chemical Society*, 3(3), 258–266. <https://doi.org/10.1007/BF03247217>
- Enogieru, A. B., Haylett, W., Hiss, D. C., Bardien, S., & Ekpo, O. E. (2018). Rutin as a potent antioxidant: implications for neurodegenerative disorders. *Oxidative Medicine and Cellular Longevity*, 1–17. <https://doi.org/10.1155/2018/6241017>
- Fajeriati, N. dan A. (2017). Uji aktivitas antibakteri ekstrak etanol rimpang kencur (*Kaempferia galanga* L.) pada bakteri *Bacillus subtilis* dan *Escherichia coli*. *Journal of Current Pharmaceutical Science*, 1(1), 36–41.
- Farida. (2018). *Pengaruh jenis kemasan dan kondisi penyimpanan terhadap kadar fenol, sifat fisikokimia, mikrobiologis, dan organoleptik minuman beras kencur dari beras merah (Oryza nivara)*.
- Gawlik-dziki, U. (2008). Effect of hydrothermal treatment on the antioxidant properties of broccoli (*Brassica oleracea* var. botrytis italica) florets. *Food Chemistry*, 109, 393–401. <https://doi.org/10.1016/j.foodchem.2007.12.058>
- Gilman, A. G., Goodman, L. S., & Gilman, A. (1981). The pharmacological basis of therapeutics. In *Macmillan Publishing Co.* (p. 1609).
- Gordon, M. H. (1990). *The mechanism of antioxidants action in vitro*. Elsevier Applied Science.
- Gowda, S., Desai, P. B., Hull, V. V., Math, A. A. K., & Sonal, N. (2009). A review on laboratory liver function tests. *PanAfrican Medical Journal*, 1–11.
- Grotto, D., Federal, U., Maria, D. S., & Garcia, S. (2009). Importance of the lipid peroxidation biomarkers and methodological aspects. *Quimica Nova*, 32(1), 169–174. <https://doi.org/10.1590/S0100-40422009000100032>
- Guinda, Á., Castellano, J. M., Santos-Lozano, J. M., Delgado-Hervás, T., Gutiérrez-Adán, P., & Rada, M. (2015). Determination of major bioactive compounds from olive leaf. *LWT*, 64(1), 431–438. <https://doi.org/10.1016/j.lwt.2015.05.001>
- Gül, A., & Pehlivan, T. (2018). Antioxidant activities of some monofloral honey types produced across Turkey. *Saudi Journal of Biological Sciences*, 25(6), 1056–1065. <https://doi.org/10.1016/j.sjbs.2018.02.011>
- Gulcin, I. (2011). Antioxidant activity of Eugenol: A Structure - Activity relationship study. *Journal of Medicinal Food*, 14(9), 975–985. <https://doi.org/10.1089/jmf.2010.0197>
- Hadriyono, K.R.P., & Kurniawati, A. (2011). *Karakter kulit manggis, kadar polifenol dan potensi antioksidan manggis pada berbagai umur buah dan setelah buah dipanen*. Institut Pertanian Bogor.
- Hari, N., & Nair, V. P. (2018). FTIR spectroscopic analysis of leaf extract in hexane in *Jasminum azoricum* L. *International Journal of Scientific Research and Technology*, 4(8), 170–172.
- Hasanah, A. N., Nazaruddin, F., Febrina, E., & Zuhrotun, A. (2011). Analisis kandungan minyak atsiri dan uji aktivitas antiinflamasi ekstrak rimpang

- Kencur (*Kaempferia galanga* L.). *Jurnal Matematika & Sains*, 16(3), 147–152.
- Hayati, E. K., Ningsih, R., & Latifah, L. (2016). Antioxidant activity of flavonoid from rhizome *Kaempferia galanga* L. extract. *Alchemy*, 4(2), 127. <https://doi.org/10.18860/al.v4i2.3203>
- Heim, K. E., Tagliaferro, A. R., & Bobilya, D. J. (2002). Flavonoid antioxidants: chemistry, metabolism and structure-activity relationships. *Journal of Nutritional Biochemistry*, 13, 572–584.
- Henderson, N. C., & Iredale, J. P. (2007). Liver fibrosis : cellular mechanisms of progression and resolution. *Clinical Sciences*, 280, 265–280. <https://doi.org/10.1042/CS20060242>
- Herdiana, D. D., Utami, R., & Anandito, R. B. katri. (2014). Kinetika degradasi termal aktivitas antioksidan pada minuman tradisional wedang uwuh siap minum. *Jurnal Teknosains Pangan*, 3(3), 44–53.
- Hihat, S., Remini, H., & Madani, K. (2017). Effect of oven and microwave drying on phenolic compounds and antioxidant capacity of coriander leaves. *International Food Research Journal*, 24(2), 503–509.
- Hong, T. K., Kim, S. Il, Heo, J. W., Lee, J. K., Choi, D. R., & Ahn, Y. J. (2011). Toxicity of *Kaempferia galanga* rhizome constituents to *Meloidogyne incognita* juveniles and eggs. *Nematology*, 13(2), 235–244. <https://doi.org/10.1163/138855410X517048>
- Horváthová, J., Suhaj, M., & Šimko, P. (2007). Effect of thermal treatment and storage on antioxidant activity of some spices. *Journal of Food and Nutrition Research*, 46(1), 20–27.
- Hubmann, A. M., Jammer, A., & Monschein, S. (2022). Activities of H<sub>2</sub>O<sub>2</sub> - Converting enzymes in apple leaf buds during dormancy release in consideration of the ratio change between bud scales and physiologically active tissues. *Horticulturae*, 8(982), 2–17.
- Hur, J., Pak, S. C., Koo, B., Jeon, S., Hur, J., Pak, S. C., Koo, B., & Jeon, S. (2013). Borneol alleviates oxidative stress via upregulation of Nrf2 and Bcl-2 in SH-SY5Y cells orneol alleviates oxidative stress via upregulation of. *Pharmaceutical Biology*, 51(1), 30–35. <https://doi.org/10.3109/13880209.2012.700718>
- Ibrahim, A. M., Sriherfyna, F. H., & Yuniarta. (2015). Pengaruh suhu dan lama waktu ekstraksi terhadap sifat kimia dan fisik pada pembuatan minuman sari jahe merah (*Zingiber officinale* var. *Rubrum*) dengan kombinasi penambahan madu sebagai pemanis. *Jurnal Pangan Dan Agroindustri*, 3(2), 530–541.
- Ichwan, S. J. A., Husin, A., Hidayati, W., Lestari, W., Nor, M., & Razak, A. (2019). Anti-neoplastic potential of ethyl-p-methoxycinnamate of *Kaempferia galanga* on oral cancer cell lines. *Materials Today: Proceedings*, 16, 2115–2121. <https://doi.org/10.1016/j.matpr.2019.06.100>
- Iida, Y., Tuziuti, T., Yasui, K., Towata, A., & Kozuka, T. (2008). Control of viscosity in starch and polysaccharide solutions with ultrasound after gelatinization. *Innovative Food Science and Emerging Technologies*, 9(2), 140–146. <https://doi.org/10.1016/j.ifset.2007.03.029>
- Indiarto, R., Pranoto, Y., Santoso, U., & Supriyanto. (2019). In vitro antioxidant activity and profile of polyphenol compounds extracts and their fractions on

- cacao beans. *Pakistan Journal of Biological Sciences*, 22(1), 34–44. <https://doi.org/10.3923/pjbs.2019.34.44>
- Innamorato, V., Longobardi, F., Cervellieri, S., Cefola, M., Pace, B., Capotorto, I., Gallo, V., Rizzuti, A., Logrieco, A. F., Chimica, D., Aldo, B., & Orabona, V. (2020). Quality evaluation of table grapes during storage by using <sup>1</sup>H NMR, LC- HRMS, MS-eNose and multivariate statistical analysis. *Food Chemistry*, 315(126247), 1–12. <https://doi.org/10.1016/j.foodchem.2020.126247>
- Irnawati, I., Riyanto, S., Martono, S., Windarsih, A., & Rohman, A. (2022). Physicochemical properties and antioxidant activities of pumpkin seed oil as affected by different origins and extraction methods. *Journal of Applied Pharmaceutical Science*, 12(03), 115–122. <https://doi.org/10.7324/JAPS.2022.120312>
- Islamadina, R., Chan, A., & Rohman, A. (2020). Chemometrics application for grouping and determining volatile compound which related to antioxidant activity of Turmeric essential oil (*Curcuma longa*). *Journal of Food and Pharmaceutical Sciences*, 8(2), 225–239.
- Jagadish PC, Latha KP, Mudgal J, N. G. (2016). Extraction, characterization and evaluation of *Kaempferia galanga* L. (Zingiberaceae) rhizome extracts against acute and chronic inflammation in rats. *Journal of Ethnopharmacology*, 194, 434–439. <https://doi.org/10.1016/j.jep.2016.10.010>
- Jahangiri, Y., Ghahremanib, H., Torghabeh, J. A., & Salehi, E. A. (2011). Effect of temperature and solvent on the total phenolic compounds extraction from leaves of *Ficus carica*. *Journal of Chemical and Pharmaceutical Research*, 3(5), 253–259.
- Jeon, T. Il, Hwang, S. G., Park, N. G., Jung, Y. R., Shin, S. I., Choi, S. D., & Park, D. K. (2003). Antioxidative effect of chitosan on chronic carbon tetrachloride induced hepatic injury in rats. *Toxicology*, 187(1), 67–73. [https://doi.org/10.1016/S0300-483X\(03\)00003-9](https://doi.org/10.1016/S0300-483X(03)00003-9)
- Jiang, H., Ling, B., Zhou, X., & Wang, S. (2020). Effects of combined radio frequency with hot water blanching on enzyme inactivation, color and texture of sweet potato. *Innovative Food Science and Emerging Technologies*, 66(June), 102513. <https://doi.org/10.1016/j.ifset.2020.102513>
- Jirovetz, L., Buchbauer, G., Shafi, M. P., & Leela, N. K. (2003). Analysis of the essential oils of the leaves, stems, rhizomes and roots of the medicinal plant *Alpinia galanga* from southern India. *Acta Pharm*, 53, 73–81.
- Kairupan, C. P., & Lolo, W. A. (2014). Uji daya hambat ekstrak etanol daun kembang sepatu (*Hibiscus rosa-sinensis* L) terhadap pertumbuhan bakteri *escherichia coli*. *Pharmacon*, 3(2), 93–98. <https://doi.org/10.35799/pha.3.2014.4779>
- Kandyliari, A., Potsaki, P., Bousdouni, P., Kaloteraki, C., Christofilea, M., Almpounioti, K., Moutsou, A., Fasoulis, C. K., Polychronis, L. V., Gkalpinos, V. K., Tzakos, A. G., & Koutelidakis, A. E. (2023). Development of dairy products fortified with plant extracts: antioxidant and phenolic content characterization. *Antioxidants*, 12(500), 1–21.
- Karaciğer, K., Serum, H., & Arasındaki, S. (2003). The relationship between serum malondialdehyde and ceruloplasmin in chronic liver disease. *Turkish Journal*



- of Biochemistry*, 28(May), 32–34.
- Khadhraoui, B., Ummat, V., Tiwari, B. K., Fabiano-tixier, A. S., & Chemat, F. (2021). Review of ultrasound combinations with hybrid and innovative techniques for extraction and processing of food and natural products. *Ultrasonics Sonochemistry*, 76, 105625. <https://doi.org/10.1016/j.ultsonch.2021.105625>
- Khairy, H. L., Saadoon, A. F., Zzaman, W., Yang, T. A., & Mat, A. (2018). Identification of flavor compounds in rambutan seed fat and its mixture with cocoa butter determined by SPME-GCMS. *Journal of King Saud University - Science*, 30(3), 316–323. <https://doi.org/10.1016/j.jksus.2017.03.001>
- Khlif, I., Jellali, K., Michel, T., Halabalaki, M., Skaltsounis, A. L., & Allouche, N. (2015). Characteristics, phytochemical analysis and biological activities of extracts from Tunisian Chetoui Olea europaea variety. *Journal of Chemistry*.
- Khokhar, S., & Owusu Apenten, R. K. (2003). Iron binding characteristics of phenolic compounds: Some tentative structure-activity relations. *Food Chemistry*, 81(1), 133–140. [https://doi.org/10.1016/S0308-8146\(02\)00394-1](https://doi.org/10.1016/S0308-8146(02)00394-1)
- Kilic-akyilmaz, M., Ozer, B., Bulat, T., & Topcu, A. (2022). Effect of heat treatment on micronutrients , fatty acids and some bioactive components of milk. *International Dairy Journal*, 126, 105231. <https://doi.org/10.1016/j.idairyj.2021.105231>
- Kiptiyah, S. Y., Harmayani, E., & Santoso, U. (2017). Study of microbiological quality and antioxidant activity beras kencur drink with heating process. *Indonesian Food and Nutrition Progress*, 14(2), 91–100. <https://doi.org/10.22146/ifnp.29725>
- Konaté, K., Souza, A., Thérèse, K. Y., Dibala, I. C., & Barro, N. (2011). Phytochemical composition , Antioxidant and Anti-inflammatory potential of bioactive fractions from extracts of three medicinal plants traditionally used to treat liver diseases in Burkina Faso . *International Journal of Phytomedicine*, 3, 406–415.
- Kumar, A. (2020). Phytochemistry, pharmacological activities and uses of traditional medicinal plant *Kaempferia galanga* L. – An overview. *Journal of Ethnopharmacology*, 253(October 2019), 112667. <https://doi.org/10.1016/j.jep.2020.112667>
- Kumar, S., & Pandey, A. K. (2013). Chemistry and biological activities of flavonoids : An overview. *The Scientific World Journal*, 1–16.
- Kumar, T., Zareen, K., Dasharath, K., & Bijendra, O. (2020). Profiling of polyphenols in phalsa (*Grewia asiatica* L) fruits based on liquid chromatography high resolution mass spectrometry. *Journal of Food Science and Technology*, 57(2), 606–616. <https://doi.org/10.1007/s13197-019-04092-y>
- Kusumowidyo, R. L., & Nurrokhman. (2013). *Kemampuan ekstrak air rim pang kencur (Kaempferia galanga L.) dalam menghambat pertumbuhan candida albicans secara in vitro*. Universitas Gadjah Mada.
- Lai, L., Chou, S., & Chao, W. (2001). Studies on the antioxidative activities of Hsian-tsao (*Mesona procumbens* Hemsl) leaf gum. *Journal Agriculture Food Chemistry*, 49(2), 963–968.

- Lahey-beitia, J., Burillo, A. M., Penna, G. La, Hegde, M. L., Rao, K. S., & Discovery, D. (2021). Polyphenols as potential metal chelation compounds against Alzheimer's disease. *Journal Alzheimers Disease*, 82(1), 8335–8357. <https://doi.org/10.3233/JAD-200185>. Polyphenols
- Lamia, S. S., Emran, T., Rikta, J. K., Chowdhury, N. I., Sarker, M., Jain, P., Islam, T., Gias, Z. T., Shill, M. C., & Reza, H. M. (2021). Coenzyme Q10 and silymarin reduce ccl4-induced oxidative stress and liver and kidney injury in ovariectomized rats-implications for protective therapy in chronic liver and kidney diseases. *Pathophysiology*, 28(1), 50–63. <https://doi.org/10.3390/pathophysiology28010005>
- Lapeyre, C., Bedos-belval, F., Duran, H., & Ne, A. (2005). Derivatives as Antiatherogenic Agents. *Journal of Medicinal Chemistry*, 4826, 8115–8124.
- Lee, H. Y., Kim, S. W., Lee, G. H., Choi, M. K., Jung, H. W., Kim, Y. J., Kwon, H. J., & Chae, H. J. (2016). Turmeric extract and its active compound, curcumin, protect against chronic CCl4-induced liver damage by enhancing antioxidation. *BMC Complementary and Alternative Medicine*, 16(1), 1–10. <https://doi.org/10.1186/s12906-016-1307-6>
- Li, F., Chen, G., Zhang, B., & Fu, X. (2017). Current applications and new opportunities for the thermal and non-thermal processing technologies to generate berry product or extracts with high nutraceutical contents. *Food Research International*, 100(August), 19–30. <https://doi.org/10.1016/j.foodres.2017.08.035>
- Lim, Y. H., Kim, I. H., & Seo, J. J. (2007). In vitro activity of kaempferol isolated from the *Impatiens balsamina* alone and in combination with erythromycin or clindamycin against *Propionibacterium acnes*. *Journal of Microbiology*, 45(5), 473–477.
- Lin, C. C., & Huang, P. C. (2000). Antioxidant and hepatoprotective effects of *Acatopanax senticosus*. *Phytotherapy Research*, 14(7), 489–494. [https://doi.org/10.1002/1099-1573\(200011\)](https://doi.org/10.1002/1099-1573(200011))
- Lin, Y., Shi, R., Wang, X., & Shen, H.-M. (2009). Luteolin, a flavonoid with potentials for cancer prevention and therapy. *Curr Cancer Drug Targets*, 8(7), 634–646.
- Liu, H., Cao, J., & Jiang, W. (2015). Evaluation and comparison of vitamin C, phenolic compounds, antioxidant properties and metal chelating activity of pulp and peel from selected peach cultivars. *LWT - Food Science and Technology*, 63, 1042–1048.
- Loizzo, M. R., Tundis, R., Bonesi, M., Menichini, F., Mastellone, V., Avallone, L., & Menichini, F. (2012). Radical scavenging, antioxidant and metal chelating activities of *Annona cherimola* Mill. (cherimoya) peel and pulp in relation to their total phenolic and total flavonoid contents. *Journal of Food Composition and Analysis*, 25, 179–184. <https://doi.org/10.1016/j.jfca.2011.09.002>
- Lourenço, S. C., Mold, M., & Alves, V. D. (2019). Antioxidants of natural plant origins: From sources to food industry applications. *Molecules*, 24(4132), 1–25.
- Lukito, L. P. (2017). *Application of steam blanching and citric acid pretreatments on the quality of kencur (*Kaempferia galanga* L.) dried by solar tunnel dryer.*

Universitas Sogiejapranata.

- Mahae, N., & Chaiseri, S. (2009). Antioxidant activities and antioxidative components in extracts of *Alpinia galanga* (L.) Sw. *Kasetsart Journal - Natural Science*, 43(2), 358–369.
- Malanovic, N., & Lohner, K. (2016). Antimicrobial peptides targeting gram-positive bacteria. *Pharmaceuticals*, 9(59), 2–33. <https://doi.org/10.3390/ph9030059>
- Mandha, J., Shumoy, H., Matemu, A. O., & Raes, K. (2023). Characterization of fruit juices and effect of pasteurization and storage conditions on their microbial, physicochemical, and nutritional quality. *Food Bioscience*, 51(102335), 1–15. <https://doi.org/10.1016/j.fbio.2022.102335>
- Markham, K. R. (1988). *Cara mengidentifikasi flavonoid*. Institut Teknologi Bandung.
- Martemucci, G., Costagliola, C., Mariano, M., Luca, D., Napolitano, P., & Alessandro, A. G. D. (2022). Free radical properties, source and targets, antioxidant consumption and health. *Oxygen*, 2, 48–78.
- Masubuchi, Y., & Horie, T. (2007). Toxicological significance of mechanism-based inactivation of Cytochrome P450 enzymes by drugs. *Clinical Reviews in Toxicology*, 45(0), 389–412. <https://doi.org/10.1080/10408440701215233>
- Mathialagan, R., Mansor, N., Shamsuddin, M. R., Uemura, Y., & Majeed, Z. (2017). Optimisation of Ultrasonic-Assisted Extraction (UAE) of Allicin from Garlic (*Allium sativum* L.). *Chemical Engineering Transactions*, 56(December 2016), 1747–1752. <https://doi.org/10.3303/CET1756292>
- Mendes-oliveira, G., Deering, A. J., Martin-gonzalez, M. F. S., & Campanella, O. H. (2020). Microwave pasteurization of apple juice : Modeling the inactivation of *Escherichia coli* O157 : H7 and *Salmonella* Typhimurium at 80 – 90 ° C. *Journal Of Food Microbiology*, 87(September 2019), 103382. <https://doi.org/10.1016/j.fm.2019.103382>
- Minarti, A., Budiana, T. E. (2014). Bioaktivitas turunan metil sinamat terhadap pertumbuhan bakteri *Escherichia coli*, *Bacillus aureus*, *Bacillus subtilis*, *Pseudomonas aureognosa* dan jamur *Candida albicans*. *Jurnal Kimia Valensi*, 1(1), 60–64.
- Moghimi, R., Aliahmadi, A., & Rafati, H. (2017). Ultrasonic nanoemulsification of food grade trans -cinnamaldehyde : 1 , 8-Cineol and investigation of the mechanism of antibacterial activity. *Ultrasonics - Sonochemistry*, 35, 415–421. <https://doi.org/10.1016/j.ultsonch.2016.10.020>
- Mouradov, A., & Spangenberg, G. (2014). Flavonoids : a metabolic network mediating plants adaptation to their real estate. *Frontiers in Plant Science*, 5(620), 1–16. <https://doi.org/10.3389/fpls.2014.00620>
- Movasaghi, Z., Rehman, S., & Rehman, I. (2018). Fourier Transform Infrared ( FTIR ) spectroscopy of biological tissues. *Applied Spectroscopy Reviews*, 43(2), 134–179. <https://doi.org/10.1080/05704920701829043>
- Moyer, R. A., Hummer, K. E., Finn, C. E., Frei, B., & Wrolstad, R. E. (2002). Anthocyanins, phenolics, and antioxidant capacity in diverse small fruits: *Vaccinium*, *Rubus*, and *Ribes*. *Journal of Agricultural and Food Chemistry*, 50, 519–525.

- Muhafidzah, Z., Seniwati, & Syarif, R. A. (2010). Aktivitas antioksidan fraksi rimpang kencur (*Kaempferia rhizoma*) dengan menggunakan metode peredaman 1,1 Diphenyl-2-picrylhydrazil (DPPH). *As-Syifaa*, 10(01), 44–50.
- Munde, T., Brand, S., Hidalgo, W., Maddula, R. K., Svatoš, A., & Schneider, B. (2013). Biosynthesis of tetraoxygenated phenylphenalenones in *Wachendorfia thyrsoiflora*. *Phytochemistry*, 91, 165–176. <https://doi.org/10.1016/j.phytochem.2012.02.020>
- Mustafa, R. A., Hamid, A. A., Mohamed, S., & Bakar, F. A. (2010). Total phenolic compounds, flavonoids, and radical scavenging activity of 21 selected tropical plants. *Journal of Food Science*, 75(1), 28–35. <https://doi.org/10.1111/j.1750-3841.2009.01401.x>
- Mutha, R. E., Tatiya, A. U., & Surana, S. J. (2021). Flavonoids as natural phenolic compounds and their role in therapeutics : an overview. *Future Journal of Pharmaceutical Sciences*, 7(25), 1–13.
- Nabella, H. (2011). *Hubungan asupan protein dengan kadar ureum dan kreatinin, dan asam urat*. Universitas Diponegoro.
- Nandi, A., Yan, L., Jana, C. K., & Das, N. (2019). Role of catalase in oxidative stress- and age-associated degenerative diseases. *Oxidative Medicine and Cellular Longevity*, 1–19.
- Nazzaro, F., Fratianni, F., De Martino, L., Coppola, R., & De Feo, V. (2013). Effect of essential oils on pathogenic bacteria. *Pharmaceuticals*, 6(12), 1451–1474. <https://doi.org/10.3390/ph6121451>
- Ndhlala, A. R., Moyo, M., & Staden, J. Van. (2010). Natural antioxidants : fascinating or mythical biomolecules. *Molecules*, 15, 6905–6930. <https://doi.org/10.3390/molecules15106905>
- Neoh, Y. Y., Matanjun, P., & Lee, J. S. (2016). Comparative study of drying methods on chemical constituents of Malaysian red seaweed. *Drying Technology*, 34(14), 1745–1751. <https://doi.org/10.1080/07373937.2016.1212207>
- Nie, J., Chen, D., Lu, Y., & Dai, Z. (2021). Effects of various blanching methods on fucoxanthin degradation kinetics, antioxidant activity, pigment composition, and sensory quality of *Sargassum fusiforme*. *LWT*, 143, 111179. <https://doi.org/10.1016/j.lwt.2021.111179>
- Nurlita, F., & Suja, I. W. (2004). *Buku ajar praktikum kimia organik*. IKIP NEgeri Singaraja.
- Oliveira, T. M. De, Bruno, R., Carvalho, F. De, Fernandes, H., Antônio, G., Oliveira, L. De, Souza, A. A. De, Lima, S. G. De, De, R. M., Oliveira, T. M. De, Bruno, R., Carvalho, F. De, & Henrique, I. (2015). Evaluation of p - cymene , a natural antioxidant. *Pharmaceutical Biology*, 53(3), 423–428. <https://doi.org/10.3109/13880209.2014.923003>
- Oobrattee, M. O. A. S., & Ruoma, O. K. I. A. (2002). Antioxidant activities of phenolic, proanthocyanidin, and flavonoid components in extracts of *Cassia fistula*. *Journal Agriculture Food Chemistry*, 50, 5042–5047. <https://doi.org/10.1021/jf0201172>
- Oonmetta-aree, J., Suzuki, T., Gasaluck, P., & Eumkeb, G. (2006). Antimicrobial properties and action of galangal (*Alpinia galanga* Linn.) on *Staphylococcus*

- aureus. *LWT*, 39, 1214–1220. <https://doi.org/10.1016/j.lwt.2005.06.015>
- Ortelli, D., Rudaz, S., Cognard, E., & Veuthey, J. L. (2000). Analysis of dihydroartemisinin in plasma by liquid chromatography - Mass spectrometry. *Chromatographia*, 52(7–8), 445–450. <https://doi.org/10.1007/BF02535717>
- Panjaitan, R. G. P., Handharyani, E., Chairul, Masriani, Zakiah, Z., & Manalu, W. (2007). Pengaruh pemberian karbon tetraklorida terhadap fungsi hati dan ginjal tikus. *Makara Kesehatan*, 11(1), 11–16.
- Park, K., Lee, H., Cho, M., & Yun, H. (2020). A phytochemical constituent, (E)-Methyl-Cinnamate isolated from *Alpinia katsumadai* Hayata suppresses cell survival, migration, and differentiation in pre-osteoblasts. *International Journal of Molecular Science*, 21(3700), 1–14.
- Parwata, I. M. O. (2016). Kimia Organik Bahan Alam Flavanoid. *Diklat / Bahan Ajar*, 1–51.
- Parwata, I. M. O., & Dewi, P. F. S. (2008). Isolasi dan uji aktivitas antibakteri minyak atsiri dari rimpang lengkuas (*Alpinia galanga* L.). *Jurnal Kimia*, 2(2), 100–104.
- Pathak, D. K., & Behl, S. (2019). Extraction of essential oil and identification of clove Oil by Clevenger Apparatus and chromatographic analysis. *Journal of Emerging Technologies and Innovative Research*, 6(6), 363–367.
- Patras, A., Tiwari, B. K., & Brunton, N. P. (2011). Influence of blanching and low temperature preservation strategies on antioxidant activity and phytochemical content of carrots, green beans and broccoli. *LWT - Food Science and Technology*, 44(1), 299–306. <https://doi.org/10.1016/j.lwt.2010.06.019>
- Payne, A. C., Mazzer, A., Clarkson, G. J. J., & Taylor, G. (2013). Antioxidant assays - consistent findings from FRAP and ORAC reveal a negative impact of organic cultivation on antioxidant potential in spinach but not watercress or rocket leaves. *Food Science & Nutrition*, 1(6), 439–444. <https://doi.org/10.1002/fsn3.71>
- Phanthong, P., Lomarat, P., Chomnawang, M. T., & Bunyapraphatsara, N. (2013). Antibacterial activity of essential oils and their active components from Thai spices against foodborne pathogens. *ScienceAsia*, 39(5), 472–476. <https://doi.org/10.2306/scienceasia1513-1874.2013.39.472>
- Pietta, P. G. (2000). Flavonoids as antioxidants. *Journal of Natural Products*, 63(7), 1035–1042. <https://doi.org/10.1021/np9904509>
- Pinent, M., Castell, A., Baiges, I., Montagut, G., Arola, L., & Ardévol, A. (2008). Bioactivity of flavonoids on insulin-secreting cells. *Comprehensive Reviews in Food Science and Food Safety*, 7(4), 299–308. <https://doi.org/10.1111/j.1541-4337.2008.00048.x>
- Pizzino, G., Irrera, N., Cucinotta, M., Pallio, G., Mannino, F., Arcoraci, V., Squadrito, F., Altavilla, D., & Bitto, A. (2017). Oxidative stress: Harms and benefits for human health. *Oxidative Medicine and Cellular Longevity*, 1–13. <https://doi.org/10.1155/2017/8416763>
- Plaper, A., Golob, M., Hafner, I., Oblak, M., Šolmajer, T., & Jerala, R. (2003). Characterization of quercetin binding site on DNA gyrase. *Biochemical and Biophysical Research Communications*, 306(2), 530–536. [https://doi.org/10.1016/S0006-291X\(03\)01006-4](https://doi.org/10.1016/S0006-291X(03)01006-4)



- Prakash, A., Rigelhof, F., & Miller, E. (2001). Antioxidant activity. *Journal of Medical Sciences*, 3(1), 87–98.
- Pratama, I. B. (2012). *Metode pengambilan minyak atsiri dalam rimpang kencur (Kaempferia galanga. L) menggunakan ekstraksi gelombang mikro*. Universitas Diponegoro.
- Primawati, S.N., Nofisulastri dan Nufida, B. A. (2013). Efektivitas senyawa bioaktif ekstrak Kencur (*Kaempferia galanga* L.) menggunakan pelarut aquades dengan pelarut metanol. *Jurnal Ilmiah Biologi "Bioscientist,"* 1(2), 102–105.
- Procházková, D., Boušová, I., & Wilhelmová, N. (2011). Antioxidant and prooxidant properties of flavonoids. *Fitoterapia*, 82, 513–523. <https://doi.org/10.1016/j.fitote.2011.01.018>
- Pujimulyani, D; Santoso, U; Raharjo, S; Marsono, Y. (2012). The effect of blanching on antioxidant activity and glycosides of white saffron (*Curcuma mangga* Val.). *International Food Research Journal*, 19(2), 617–621.
- Pujimulyani, D., Santoso, U., D, S. L., & Maruf, A. (2020). Orally administered pressure-blanching white saffron (*Curcuma mangga* Val.) improves antioxidative properties and lipid profiles in vivo. *Heliyon*, 6(e04219). <https://doi.org/10.1016/j.heliyon.2020.e04219>
- Qiang, Y., Si, R., Tan, S., Wei, H., Huang, B., Wu, M., Shi, M., Fang, L., Fu, J., & Zeng, S. (2021). Spatial variation of volatile organic compounds and antioxidant activity of turmeric (*Curcuma longa* L.) essential oils harvested from four provinces of China. *Current Research in Food Science*, 4, 882–890. <https://doi.org/10.1016/j.crfs.2021.11.002>
- R, F. J., M, D. A. C., Nirwani, B., Nurmasitoh, T., & Bowo, E. T. (n.d.). Manfaat sirih merah (*Piper crocatum*) sebagai agen anti bakterial terhadap bakteri gram positif dan gram negatif. *Jurnal Kedokteran Dan Kesehatan Indonesia*.
- Rababah, T. M., Al-Mahasneh, M. A., Kilani, I., Yang, W., Alhamad, M. N., Ereifej, K., & Al-u'datt, M. (2011). Effect of jam processing and storage on total phenolics, antioxidant activity, and anthocyanins of different fruits. *Journal of the Science of Food and Agriculture*, 91(6), 1096–1102. <https://doi.org/10.1002/jsfa.4289>
- Rabib, H., Elagdi, C., Hsaine, M., Fougrach, H., Koussa, T., & Badri, W. (2020). Antioxidant and antibacterial activities of the essential oil of Moroccan *Tetraclinis articulata* (Vahl) masters. *Biochemistry Research International*, 1–5.
- Randhir, R., Kwon, Y. I., & Shetty, K. (2008). Effect of thermal processing on phenolics, antioxidant activity and health-relevant functionality of select grain sprouts and seedlings. *Innovative Food Science and Emerging Technologies*, 9(3), 355–364. <https://doi.org/10.1016/j.ifset.2007.10.004>
- Rao, N., & Kaladhar, D. (2014). Antioxidant and antimicrobial activities of rhizome extracts of *Kaempferia galanga*. *World Journal of Pharmacy and Pharmaceutical Sciences*, 3(5), 1180–1189.
- Regianto, H. (2009). *Minyak atsiri rimpang kencur (Kaempferia galanga L) karakterisasi simplisia, isolasi, dan analisis komponen minyak atsiri secara GC-MS*.

- Reineccius. (2016). *Flavor chemistry and technology*. CRC Press.
- Rice-Evans, C. A., Miller, N. J., & Paganga, G. (1997). Antioxidant properties of phenolic compounds. *Trends in Plant Science*, 2(4), 152–159. [https://doi.org/10.1016/S1360-1385\(97\)01018-2](https://doi.org/10.1016/S1360-1385(97)01018-2)
- Riitta Puupponen-Pimiä, Suvi T Häkkinen, Marjukka Aarni, Tapani Suortti, Anna-Maija Lampi, Merja Eurola, Vieno Piironen, A. M. N. and, & Oksman-Caldentey, K.-M. (2003). Blanching and long-term freezing affect various bioactive compounds of vegetables. *Journal of the Science of Food and Agriculture*, 83, 1389–1402. <https://doi.org/10.1002/jsfa.1589>
- Riyani, D. W. W., Rohadi, & Pratiwi, E. (2018). *Variasi suhu maserasi terhadap rendemen dan karakteristik minyak atsiri jahe emprit*. Universitas Semarang.
- Rohman, A., Riyanto, S., & Utari, D. (2006). Antioxidant activities, total phenolic and flavonoid contents of ethyl acetate extract of Mengkudu (*Morinda citrifolia*, L) fruit and its fractions. *Majalah Farmasi Indonesia*, 17(3), 136–142.
- Rosa, A., Deiana, M., Casu, V., Paccagnini, S., Appendino, G., Ballero, M., & Dessi, M. A. (2002). Antioxidant activity of Capsinoids. *Journal Agricultural and Food Chemistry*, 50, 7396–7401. <https://doi.org/10.1021/jf020431w>
- Sari, D. A. P., Ningrum, A., & Anggrahini, S. (2021). Effect of temperature and drying time on Angiotensin Converting Enzyme (Ace) inhibitor and antioxidant activity of Soybean (*Glycine max* L.) - Jackbean (*Canavalia ensiformis* L.) mixed grains tempeh flour. *Indonesian Food and Nutrition Progress*, 18(1), 1–8. <https://doi.org/10.22146/ifnp.53535>
- Schymanski, E. L., Singer, H. P., Slobodnik, J., Ipolyi, I. M., Oswald, P., Krauss, M., Schulze, T., Haglund, P., Letzel, T., Grosse, S., Thomaidis, N. S., Bletsou, A., Zwiener, C., Ibáñez, M., Portolés, T., Boer, R. De, Reid, M. J., & Onghena, M. (2015). Non-target screening with high-resolution mass spectrometry: critical review using a collaborative trial on water analysis. *Anal Bioanal Chem*, 407, 6237–6255. <https://doi.org/10.1007/s00216-015-8681-7>
- Sekarsari, S., Rai Widarta, I. W., & Anom Jambe, A. A. G. N. (2019). Pengaruh suhu dan waktu ekstraksi dengan gelombang ultrasonik terhadap aktivitas antioksidan ekstrak daun jambu biji (*Psidium guajava* L.). *Jurnal Ilmu Dan Teknologi Pangan*, 8(3), 267–277.
- Serly, S. (2022). *Pengaruh suhu dan lama waktu termosonikasi terhadap senyawa antioksidan dan kualitas minuman beras kencur*.
- Shahidi, F., & Ambigaipalan, P. (2015). Phenolics and polyphenolics in foods, beverages and spices: Antioxidant activity and health effects - A review. *Journal of Functional Foods*, 18, 820–897. <https://doi.org/10.1016/j.jff.2015.06.018>
- Sharma, N., Biswas, S., Al-dayan, N., & Alhegaili, A. S. (2021). Antioxidant role of Kaempferol in prevention of hepatocellular carcinoma. *Antioxidants*, 10(1419), 1–17.
- Shebis, Y., Iluz, D., Kinel-Tahan, Y., Dubinsky, Z., & Yehoshua, Y. (2013). Natural antioxidants: Function and sources. *Food and Nutrition Sciences*, 4, 643–649. <https://doi.org/10.4236/fns.2013.46083>
- Shirazi, M. T., Gholami, H., Kavosi, G., Rowshan, V., & Tafsiry, A. (2014).

- Chemical composition, antioxidant, antimicrobial and cytotoxic activities of *Tagetes minuta* and *Ocimum basilicum* essential oils. *Food Science & Nutrition*, 2(2), 146–155. <https://doi.org/10.1002/fsn3.85>
- Sikorska, M., & Matlawska, I. (2008). Polyphenolic compounds from *Abutilon grandiflorum* leaves. *Acta Poloniae Pharmaceutica - Drug Research*, 65(4), 467–471.
- Silalahi, M. (2017). Essensial oil pada *Alpinia galanga* (L.) willd dan pemanfaatannya. *Jurnal Pro-Life*, 4(1), 287–296.
- Singal, A. K., Jampana, S. C., & Weinman, S. A. (2011). Antioxidants as therapeutic agents for liver disease. *Liver International*, 31(10), 1432–1448. <https://doi.org/10.1111/j.1478-3231.2011.02604.x>
- Singh, G., Kapoor, I. P. S., Singh, P., Heluani, C. S. De, Lampasona, M. P. De, & Catalan, C. A. N. (2010). Comparative study of chemical composition and antioxidant activity of fresh and dry rhizomes of turmeric (*Curcuma longa* Linn.). *Food and Chemical Toxicology*, 48(4), 1026–1031. <https://doi.org/10.1016/j.fct.2010.01.015>
- Singh, S., Sahoo, S., & Sahoo, B. C. (2022). Derivatives of cinnamic acid esters and terpenic diversity in volatiles of thirty-six sand ginger (*Kaempferia galanga* L.). *Molecules*, 27(1116), 1–15.
- Solecki, O., Mosbah, A., Floc, B., & Felden, B. (2015). Converting a *Staphylococcus aureus* toxin into effective cyclic Pseudopeptide antibiotics. *Chemistry & Biology*, 22, 329–335. <https://doi.org/10.1016/j.chembiol.2014.12.016>
- Song, L., Wu, X., Xie, J., Zhang, H., Yang, H., Zeng, Q., Yang, X., & Xie, W. (2021). *Kaempferia galanga* Linn. Extract – A potential antibacterial agent for preservation of poultry products. *LWT - Food Science and Technology*, 147(111553), 1–11. <https://doi.org/10.1016/j.lwt.2021.111553>
- Song, Y., & March, J. (2022). Hyperuricemia and the small intestine : Transport mechanisms and. *Biotechnology Notes*, 3(November 2021), 32–37. <https://doi.org/10.1016/j.biotno.2022.05.001>
- Soong, Y., & Barlow, P. J. (2004). Antioxidant activity and phenolic content of selected fruit seeds. *Food Chemistry*, 88, 411–417. <https://doi.org/10.1016/j.foodchem.2004.02.003>
- Souza, T.P., de Oliveira, P.R. Pereira, B. (2005). Physical exercise and oxidative stress. Effect of intense physical exercise on the urinary chemiluminescence and plasmatic malondialdehyde. *Revista Brasileira de Medicina Do Esporte*, 11(1).
- Sova, M. (2012). Antioxidant and antimicrobial activities of Cinnamic Acid derivatives. *Mini Reviews in Medical Chemistry*, 12, 749–767.
- Sripour, W., & Jinda, N. (2014). Effect of *Alpinia galanga* Essential Oil on Bacteria Spoilage. *The 26th Annual Meeting of the Thai Society for Biotechnology and International Conference*, 15(3), 382–387.
- Srivastava, N., Mishra, S., Iqbal, H., Chanda, D., & Shanker, K. (2021). Standardization of *Kaempferia galanga* L. rhizome and vasorelaxation effect of its key metabolite ethyl p -methoxycinnamate. *Journal of Ethnopharmacology*, 271(113911), 1–10.

- <https://doi.org/10.1016/j.jep.2021.113911>
- Srivastava, N., Ranjana, Singh, S., Gupta, A. C., Shanker, K., Bawankule, D. U., & Luqman, S. (2019). Aromatic ginger (*Kaempferia galanga* L.) extracts with ameliorative and protective potential as a functional food, beyond its flavor and nutritional benefits. *Toxicology Reports*, 6(June), 521–528. <https://doi.org/10.1016/j.toxrep.2019.05.014>
- Stephane Caillet, Stephane Salmieri, M. L. (2006). Evaluation of free radical-scavenging properties of commercial grape phenol extracts by a fast colorimetric method. *Journal of Food Chemistry*, 95, 1–8. <https://doi.org/10.1016/j.foodchem.2004.12.011>
- Subandrate, Faisal, M. E., & Anggraini, N. W. (2017). Peranan stres oksidatif pada preeklampsia. *Cermin Dunia Kedokteran*, 44(5), 353–355. <http://www.who>
- Suganthi, N., Nisha, S. A., Pandian, S. K., & Devi, K. P. (2013). Evaluation of *Gelidiella acerosa*, the red algae inhabiting South Indian coastal area for antioxidant and metal chelating potential. *Biomedicine & Preventive Nutrition*, 3, 399–406.
- Sultan, M. T., Butt, M. S., Anjum, F. M., Jamil, A., Akhtar, S., & Nasir, M. (2009). Nutritional profile of indigenous cultivar of black cumin seeds and antioxidant potential of its fixed and essential oil. *Pakistan Journal of Botany*, 41(3), 1321–1330.
- Suryanti, V., Marliyana, S. D., & Musmualim, M. (2018). Identifikasi senyawa kimia dalam buah Kundur (*Benincasa hispida* (Thunb) Cogn.) dengan Kromatografi Gas-Spektrometer Massa (KG-SM). *Alchemy Jurnal Penelitian Kimia*, 14(1), 100. <https://doi.org/10.20961/alchemy.14.1.13496.100-110>
- Sutthanont, N., Choochote, W., Tuetun, B., Junkum, A., Jitpakdi, A., Chaithong, U., Riyong, D., & Pitasawat, B. (2010). Chemical composition and larvicidal activity of edible plant-derived essential oils against the pyrethroid-susceptible and -resistant strains of *Aedes aegypti* (Diptera: Culicidae). *Journal of Vector Ecology*, 35(1), 106–115. <https://doi.org/10.1111/j.1948-7134.2010.00066.x>
- Sy, E., Kadri, H., & Yerizel, E. (2015). Efek pemberian vitamin C terhadap aktifitas katalase hati tikus galur wistar yang Terpapar ion Pb. *Jurnal Kesehatan Andalas*, 4(1), 279–285.
- Tabaraki, R., & Sadeghinejad, N. (2013). Comparison of the antioxidant activity of volatile compounds of traditional herbal waters per serving cup. *Iranian Journal of Pharmaceutical Sciences*, 9(1), 47–54.
- Taha, A., Mehany, T., Pandiselvam, R., Siddiqui, S. A., Mir, N. A., Malik, M. A., Sujayasree, O. J., Chaitanya, K., Khanashyam, A. C., Casanova, F., Xu, X., Khanashyam, A. C., Casanova, F., Xu, X., Pan, S., & Hu, H. (2023). Sonoprocessing : mechanisms and recent applications of power ultrasound in food. *Critical Reviews in Food Science and Nutrition*, 0(0), 1–39. <https://doi.org/10.1080/10408398.2022.2161464>
- Tappi, E. S., Lintong, P., & Loho, L. L. (2013). Gambaran histopatologi hati tikus wistar yang diberikan jus tomat (*Solanum Lycopersium*) pasca kerusakan hati wistar yang diinduksi karbon tetraklorida (CCl<sub>4</sub>). *Jurnal E-Biomedik*, 1(3), 1126–1129.
- Tawata, S., Taira, S., Kobamoto, N., Zhu, J., Toyama, S., & Ata, S. T. A. W. (1996).

- Synthesis and antifungal activity of Cinnamic Acid esters. *Bioscience, Biotechnology, and Biochemistry*, 60(5), 909–910. <https://doi.org/10.1271/bbb.60.909>
- Tayman, C., Cekmez, F., Kafa, I. M., Canpolat, F. E., Cetinkaya, M., Uysal, S., & Tunc, T. (2012). *Beneficial Effects of Nigella Sativa Oil on Intestinal Damage in Necrotizing Enterocolitis*. 286–294. <https://doi.org/10.3109/08941939.2011.639849>
- Teow, C. C., Truong, V. Den, McFeeters, R. F., Thompson, R. L., Pecota, K. V., & Yencho, G. C. (2007). Antioxidant activities, phenolic and  $\beta$ -carotene contents of sweet potato genotypes with varying flesh colours. *Food Chemistry*, 103(3), 829–838. <https://doi.org/10.1016/j.foodchem.2006.09.033>
- Teselkin, Y. O., Babenkova, I. V., Kolhir, V. K., Baginskaya, A. I., Tjukavkina, N. A., Kolesnik, Y. A., Selivanova, I. A., & Eichholz, A. A. (2000). Dihydroquercetin as a means of antioxidative defence in rats with tetrachloromethane hepatitis. *Phytotherapy Research*, 14(3), 160–162. [https://doi.org/10.1002/\(SICI\)1099-1573\(200005\)14\(3\)<160::AID-PTER160>3.0.CO;2-1](https://doi.org/10.1002/(SICI)1099-1573(200005)14(3)<160::AID-PTER160>3.0.CO;2-1)
- Tewtrakul, S., Yuenyongsawad, S., Kummee, S., & Atsawajaruwan, L. (2005). Chemical components and biological activities of volatile oil of *Kaempferia galanga* Linn. *Songklanarin Journal of Science and Technology*, 27(2), 503–507.
- Tian, C., Liu, X., Chang, Y., Wang, R., Lv, T., Cui, C., & Liu, M. (2021). Investigation of the antiinflammatory and antioxidant activities of luteolin, kaempferol, apigenin and quercetin. *South African Journal of Botany*, 137, 257–264. <https://doi.org/10.1016/j.sajb.2020.10.022>
- Tohge, T., Watanabe, M., Hoefgen, R., & Fernie, A. R. (2013). Shikimate and phenylalanine biosynthesis in the green lineage. *Frontiers in Plant Science*, 4(62), 1–13. <https://doi.org/10.3389/fpls.2013.00062>
- Tsimogiannis, D., & Oreopoulou, V. (2019). Classification of phenolic compounds in plants. *Polyphenols in Plants*, 263–284. <https://doi.org/10.1016/b978-0-12-813768-0.00026-8>
- Turkmen, N., Sari, F., & Velioglu, Y. S. (2005). The effect of cooking methods on total phenolics and antioxidant activity of selected green vegetables. *Food Chemistry*, 93(4), 713–718. <https://doi.org/10.1016/j.foodchem.2004.12.038>
- Ukieyanna, E., Suryani, & Roswien, A. P. (2012). *Aktivitas antioksidan kadar fenolik dan flavonoid total tumbuhan suruhan*. Institut Pertanian Bogor.
- Ullah, A., Munir, S., Badshah, S. L., Khan, N., Ghani, L., Poulson, B. G., Emwas, A., & Jaremko, M. (2020). Important flavonoids and their role as a therapeutic agent. *Molecules*, 25(2543), 1–39. <https://doi.org/10.3390/molecules25225243>
- Ultee, A., Bennik, M. H. J., & Moezelaar, R. (2002). The phenolic hydroxyl group of carvacrol is essential for action against the food-borne pathogen *Bacillus cereus*. *Applied and Environmental Microbiology*, 68(4), 1561–1568. <https://doi.org/10.1128/AEM.68.4.1561-1568.2002>
- Umar, M. I., Asmawi, M. Z. Bin, Sadikun, A., Altaf, R., & Iqbal, M. A. (2011). Phytochemistry and medicinal properties of *kaempferia galanga* L. (Zingiberaceae) extracts. *African Journal of Pharmacy and Pharmacology*,



- 5(14), 1638–1647. <https://doi.org/10.5897/AJPP11.388>
- Umar, M. I., Asmawi, M. Z., Sadikun, A., Atangwho, I. J., Yam, M. F., Altaf, R., & Ahmed, A. (2012). Bioactivity-guided isolation of ethyl-p-methoxycinnamate, an anti-inflammatory constituent, from *Kaempferia galanga* L. extracts. *Molecules*, 17(7), 8720–8734. <https://doi.org/10.3390/molecules17078720>
- Umemura, T., Kodama, Y., Hioki, K., Inoue, T., Nomura, T., & Kurokawa, Y. (2001). Butylhydroxytoluene (BHT) increases susceptibility of transgenic rasH2 mice to lung carcinogenesis. *Journal of Cancer Research and Clinical Oncology*, 127(10), 583–590. <https://doi.org/10.1007/s004320100268>
- Vegara, S., Mena, P., Martí, N., Saura, D., & Valero, M. (2013). Approaches to understanding the contribution of anthocyanins to the antioxidant capacity of pasteurized pomegranate juices. *Food Chemistry*, 141(3), 1630–1636. <https://doi.org/10.1016/j.foodchem.2013.05.015>
- Verdecchia, P., Schillaci, G., Reboldi, G., Santeusano, F., Porcellati, C., & Brunetti, P. (2000). Relation between serum uric acid and risk of cardiovascular disease in essential hypertension. *Hypertension*, 1072–1079.
- Walkowiak-Tomczak, D. (2007). Changes in antioxidant activity of Black Chokeberry juice concentrate solutions during storage. *ACTA Acta Sci. Pol., Technol. Aliment*, 6(2), 49–55. [https://www.food.actapol.net/pub/5\\_2\\_2007.pdf](https://www.food.actapol.net/pub/5_2_2007.pdf)
- Wannissorn, B., & Jarikasem, S. (2005). Antibacterial properties of essential oils from Thai medicinal plants. *Fitoterapia*, 76, 233–236. <https://doi.org/10.1016/j.fitote.2004.12.009>
- Wen, C., Zhang, J., Zhang, H., Dzah, C. S., & Zandile, M. (2018). Advances in ultrasound assisted extraction of bioactive compounds from cash crops – A review. *Ultrasonics - Sonochemistry*, 48(May), 538–549. <https://doi.org/10.1016/j.ultsonch.2018.07.018>
- Weston-Green, K., Clunas, H., & Jimenez Naranjo, C. (2021). A review of the potential use of Pinene and Linalool as Terpene-based medicines for brain health: Discovering novel therapeutics in the flavours and fragrances of Cannabis. *Frontiers in Psychiatry*, 12(August), 1–19. <https://doi.org/10.3389/fpsy.2021.583211>
- Widodo, H., Sisindari, S., Asmara, W., & Rohman, A. (2019). Antioxidant activity, total phenolic and flavonoid contents of selected medicinal plants used for liver diseases and its classification with chemometrics. *Journal of Applied Pharmaceutical Science*, 9(06), 99–105. <https://doi.org/10.7324/JAPS.2019.90614>
- Wiesner, J., Mitsch, A., Wiûner, P., & Schlitzer, M. (2001). Structure ± activity Relationships of novel Antimalarial agents. *Bioorganic & Medicinal Chemistry Letters*, 11, 423–424.
- X. Yue and Z. Xu. (2008). Changes of anthocyanins, anthocyanidins, and antioxidant activity in Bilberry extract during dry heating. *Journal of Food Chemistry*, 73(6), 494–499. <https://doi.org/10.1111/j.1750-3841.2008.00845.x>
- Xiao, H., Deng, L., El-mashad, H. M., Yang, X., Mujumdar, A. S., & Gao, Z.

- (2017). Recent developments and trends in thermal blanching-a comprehensive review. *Information Processing in Agriculture*. <https://doi.org/10.1016/j.inpa.2017.02.001>
- Xu, B. J., & Chang, S. K. C. (2007). A comparative study on phenolic profiles and antioxidant activities of legumes as affected by extraction solvents. *Journal of Food Science*, 72(2). <https://doi.org/10.1111/j.1750-3841.2006.00260.x>
- Xu, Dong-ping, Li, Y., Meng, X., Zhou, T., Zhou, Y., Zheng, J., & Zhang, J. (2017). Natural Antioxidants in foods and medicinal plants: Extraction, assessment and resources. *International Journal of Molecular Sciences*, 18(96), 1–32. <https://doi.org/10.3390/ijms18010096>
- Xu, Dong, Hu, M., Wang, Y., & Cui, Y. (2019). Antioxidant activities of Quercetin and its complexes for medicinal application. *Molecules*, 24(1123), 1–15. <https://doi.org/10.3390/molecules24061123>
- Xu, J., Hu, Q., & Liu, Y. (2012). Antioxidant and DNA-protective activities of Chlorogenic Acid isomers. *Agricultural and Food Chemistry*, 60, 11625–11630. <https://doi.org/10.1021/jf303771s>
- Yang, L., Liu, H., Xia, D., & Wang, S. (2020). Antioxidant properties of camphene-based thiosemicarbazones: Experimental and theoretical evaluation. *Molecules*, 25(5). <https://doi.org/10.3390/molecules25051192>
- Yang, X., Ji, H., Feng, Y., Yu, J., & Liu, A. (2018). Structural characterization and antitumor activity of polysaccharides from kaempferia galanga L. *Oxidative Medicine and Cellular Longevity*, 2018. <https://doi.org/10.1155/2018/9579262>
- Yao, F., Huang, Y., Wang, Y., & He, X. (2018). Anti-inflammatory diarylheptanoids and phenolics from the rhizomes of kencur (*Kaempferia galanga* L.). *Industrial Crops and Products*, 125, 454–461. <https://doi.org/10.1016/j.indcrop.2018.09.026>
- Yuenyongsawad, S. (2005). Chemical components and biological activities of volatile oil of *Kaempferia galanga* Linn. *Songklanarin Journal of Science and Technology*, 27(2), 503–507.
- Yuliantari, N. W. A., Widarta, I. W. R., & Permana, I. D. G. M. (2017). Pengaruh suhu dan waktu ekstraksi terhadap kandungan flavonoid dan aktivitas antioksidan daun sirih (Annona muricata L.) menggunakan ultrasonik. *Media Ilmiah Teknologi Pangan*, 4(1), 35–42.
- Yusof, N. B., Hasan, M. H., Armayni, U. A., Bin, M. S., Mohsin, H. F., & Wahab, I. A. (2013). The ferrous ion chelating assay of Pandanus extracts. *The Open Conference Proceedings Journal*, 4, 155.
- Zainol, M., Bakar, A., & Dek, P. (2009). Effect of different drying methods on the degradation of selected flavonoids in *Centella asiatica*. *International Food Research Journal*, 16, 531–537.
- Zeng, X., Tang, W., Ye, G., Ouyang, T., Tian, L., Ni, Y., & Li, P. (2010). Studies on disinfection mechanism of electrolyzed oxidizing water on *E. coli* and *S. aureus*. *Journal of Food Sc*, 75(5), 253–260. <https://doi.org/10.1111/j.1750-3841.2010.01649.x>
- Zhang, Q. W., Lin, L. G., & Ye, W. C. (2018). Techniques for extraction and isolation of natural products : a comprehensive review. *Chinese Medicine*, 1–

26. <https://doi.org/10.1186/s13020-018-0177-x>
- Zhang, Y., Liu, X., Wang, Y., Jiang, P., & Quek, S. (2015). Antibacterial activity and mechanism of cinnamon essential oil against *Eschericia coli* and *Staphlococcus aureus*. *Food Control*. <https://doi.org/10.1016/j.foodcont.2015.05.032>
- Zhao, L., Zhang, H., Hao, T., Li, S., & Page, S. D. S. (2015). In vitro antibacterial activities and mechanism of sugar fatty acid esters against five food-related bacteria. *Food Chemistry*, 187, 370–377. <https://doi.org/10.1016/j.foodchem.2015.04.108>
- Zi-min, C., Jian-qing, P., Yi, C., Ling, T., Yan-yan, Z., Ling-yun, F., Qing-de, L., & Xiang-chun, S. (2020). 1,8-Cineole: a review of source, biological activities, and application. *Journal of Asian Natural Products Research*, 0(0), 1–17. <https://doi.org/10.1080/10286020.2020.1839432>
- Zulak, K. G., & Bohlmann, J. (2010). Terpenoid biosynthesis and specialized vascular cells of conifer defense. *Journal of Integrative Plant Biology*, 52(1), 86–97. <https://doi.org/10.1111/j.1744-7909.2010.00910.x>