

DAFTAR PUSTAKA

- Alothman, M., Bhat, R. and Karim, A.A. (2009) Antioxidant Capacity and Phenolic Content of Selected Tropical Fruits from Malaysia, Extracted with Different Solvents. *Food Chemistry*, 115, 785-788. <http://dx.doi.org/10.1016/j.foodchem.2008.12.005>.
- Hairmansis, A., Supartopo, Kustianto, B., Suwarno, dan Pane, H. (2012). Perakitan dan Pengembangan Varietas Unggul Baru Padi Toleran Rendaman Air Inpara 4 dan Inpara 5 untuk Daerah Rawan Banjir.
- AOAC. (2005). *Official Methods of Analysis of the Association of Official Analytical Chemist*. Association of Official Analytical Chemist. Washington DC.
- Astawan, M., dan Wresdiyati, T. (2004). *Diet Sehat dengan Makanan Berserat*. Tiga Serangkai Pustaka Mandiri. Solo.
- Badan Pengawas Obat dan Makanan (BPOM). (2011). *Peraturan Kepala Badan Pengawas Obat dan Makanan tentang Pengawasan Klaim dalam dan Iklan Pangan Olahan*. BPOM. Jakarta.
- Badan Pusat Statistik (BPS). (2022, 1 Maret). Luas Panen dan Produksi Padi di Indonesia 2021. Diakses pada 10 Januari 2023. <https://www.bps.go.id/website/images/Luas-Panen-dan-Produksi-Padi-2021--Angka-Tetap--ind.jpg>.

- Barros, M., Fleuri, L. F., & Macedo, G. A.. (2010). Seed lipases: sources, applications and properties - a review. *Brazilian Journal of Chemical Engineering*, 27(1), 15-29.
- Butsat, S. dan Siriamornpun, S. (2010). Antioxidant capacities and phenolic compounds of the husk, bran and endosperm of Thai rice. *Food Chemistry* 119: 606-613.
- Capanzana, M. V., & Buckle, K. A. (1997). Optimisation of germination conditions by response surface methodology of a high amylose rice (*Oryza sativa*) cultivar. *LWT – Food Science and Technology*, 30, 155–163.
- Chinma, C. E., Anuonye, J. C., Simon, O. C., Ohiare, R. O., & Danbaba, N. (2015). Effect of germination on the physicochemical and antioxidant characteristics of rice flour from three rice varieties from Nigeria. *Food Chemistry*, Vol. 185, 454–458.
<https://doi.org/10.1016/j.foodchem.2015.04.010>.
- Chin, W. W. (1998). The Partial Least Squares Approach to Structural Equation Modeling. *Modern Methods for Business Research*, 295, 336
- Cho, D. H., dan Lim, S. T. (2015). Germinated brown rice and its bio-functional compounds. *Food Chemistry*, 196(8), 259–271.
<https://doi:10.1016/j.foodchem.2015.09.025>.
- Cevallos-Casals, B. A., & Cisneros-Zevallos, L. (2010). Impact of germination on phenolic content and antioxidant activity of 13 edible seed species. *Food Chemistry*, 119(4), 1485–1490.
<https://doi.org/10.1016/j.foodchem.2009.09.030>.

- Czernicka, L., Kukula-Koch, W., dan Marzec, Z. (2017). Quantitative Analysis of Phenolic Compounds in Different Types o Pickleg Ginger. *Journal Problemy Higieny I Epidemiologi*, 98(3): 240-244.
- David, J. M., dan Gurdev, S. K. (2018). IR-64: a high quality and high yielding mega variety. Rice (N Y). <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5890005/>. Diakses pada 9 Juni 2023.
- Fernandez-Orozco, R., Frias, J., Zielinski, H., Piskula, M. K., Kozłowska, H., & Vidal-Valverde, C. (2008). Kinetic study of the antioxidant compounds and antioxidant capacity during germination of *Vigna radiata* cv. Emerald, *Glycine max* cv. jutro and *Glycine max* cv. merit. *Food Chemistry*, 111(3), 622–630.
- Gepstin S, Han I, (1980). Evidence for the involvement of cytokinin in the regulation of proteolytic activity in cotyledons of germinating beans. *Plant and Cell Physiology.*, 21(1):57-63.
- Gordon, M. H. (2001). *Measuring Antioxidant Activity*. Dalam Jan Pokorny, Nedyalka Yaishlieva-Maslarova, and Micael Gordon (ed). *Antioxidants in Food Practical Application*. Woodhead Publishing LTD. London.
- Ghozali, I. (2016) Aplikasi Analisis Multivariete Dengan Program IBM SPSS 23. Edisi 8. Semarang: Badan Penerbit Universitas Diponegoro.
- Guo, W. dan Beta, T. (2013). Phenolic acid composition and antioxidant potential of insoluble and soluble dietary fibre extracts derived from select whole-grain cereals. *Food Research International* 51: 518-525.

- Haryadi. (2008). *Teknologi Pengolahan Beras*. Gadjah Mada University Press. Yogyakarta.
- Hartati, S., Marsono, Y., Suparmo, Santoso, U. (2015). Komposisi Kimia serta Aktivitas Antioksidan Ekstrak Hidrofilik Bekatul Beberapa Varietas Padi. *AGRITECH*, Vol. 35, No. 1.
- He, D., Han, C., Yao, J., Shen, S., & Yang, P. (2011). Constructing the metabolic and regulatory pathways in germinating rice seeds through proteomic approach. *Proteomics*, 11, 2693.
- Huang, Y. S., & Ho, S. C. (2010). Polymethoxy flavones are responsible for the anti-inflammatory activity of citrus fruit peel. *Food Chemistry*, 119(3), 868–873. <https://doi.org/10.1016/J.FOODCHEM.2009.09.092>
- Jiamyangyuen, S., dan Oraikul, B. (2009). Effect of germination on antioxidants, textural, and sensorial property of germinated brown rice. *FDA journal*, 16(1): 25-33.
- Juliano, B.O. (1972). A Simplified Assay for Milled Rice Amylose Measurement. *J. of Cereal Sci*, 161. 334-336.
- Kamil. (1982). *Teknologi Benih I*. Angkasa. Bandung.
- Kahkonen, M. P., Hopta, A. I. Vourela, H. J., Rauha, J. P., Pihlaja, K., Kujala, T. S., dan Heinonen, M. (1999). Antioxidant Activity of Plant Extracts Containing Phenolic Compound. *Journal of Agricultural and Food Chemistry*, 47 (10): 3954-3962.

Kayahara, H., Tsukahara, K., & Tatai, T. (2000). Flavor, health, and nutritional quality of pre-germinated brown rice. *In 10th international flavor conference*, p. 546–551. Paros, Greece.

Kementerian Pertanian (Kementan). (2019, 27 November). Varietas Unggul Padi.

Diakses pada 10 Januari 2023.

<http://cybex.pertanian.go.id/mobile/artikel/83962/VARIETAS-UNGGUL-PADI/>.

Liu, Q., dan Yao, H. (2007). Antioxidant activities of barley seeds extract. *Food Chemistry*, 102: 732-737.

Lu, Z. H., Zhang, Y., Li, L. T., Curtis, R. B., Kong, X. L., Fulcher, R. G., ... Cao, W. (2010). Inhibition of microbial growth and enrichment of gamma-aminobutyric acid during germination of brown rice by electrolyzed oxidizing water. *Journal of Food Protection*, 73, 483–487.

Manna, K. M., Naing, K. M., Pe, H. (1995). Amylase activity of some roots and sprouted cereals an beans. *Food and Nutrition Bulletin*, 16(2):1-5. doi:10.1177/156482659501600211.

Miyoshi, K., Sato, T., & Takahashi, N. (1996). Differences in the effects of dehusking during formation of seeds on the germination of seeds of indica and japonica rice (*Oryza sativa L.*). *Annals of Botany*, 77, 599–604.

Molyneux, P. (2004). The Use of Stable Free Radical Diphenilpicrylhydrazil (DPPH) for Estimating Antioxidant Activity. *J. Sci. Technol.* 26(2): 211-219.

- Muhammad, S. I., Ismail, M., Mahmud, R. B., Salisu, A. M., & Zakaria, Z. A. (2013). Germinated brown rice and its bioactives modulate the activity of uterine cells in oophorectomised rats as evidenced by gross cytohistological and immunohistochemical changes. *BMC complementary and alternative medicine*, 13, 198. <https://doi.org/10.1186/1472-6882-13-198>.
- Mamta, Misra, K., Dhillon, G, Brar, S., & Verma, M. (2014). Biotransformation of waste biomass into high value biochemicals. In *Biotransformation of Waste Biomass into High Value Biochemicals*, Chapter 6 Antioxidant. Springer Science + Business Media. <https://doi.org/10.1007/978-1-4614-8005-1>
- Oh, S.-K., Lee, J.-H., Won, Y.-J., Lee, D.-H., & Kim, C.-K. (2014). Changes of Physicochemical Properties According to the Shoot Length in Germinated Brown Rice. *Korean Journal of Cro Science*, 59(3), 223–229. <https://doi.org/10.7740/KJCS.2014.59.3.223>.
- Ohtsubo, K., Suzuki, K., Yasui, Y., Kasumi, T., (2005). Bio-functional components in the processed pre-germinated brown rice by a twin-screw extruder. *J. Food Compos. Anal.*, The International Year of Rice 2004 18, 303–316. <https://doi.org/10.1016/j.jfca.2004.10.003>.
- Panesar, P. S., dan Kaur, S. (2016). Rice: Types and Composition, Editor(s): Benjamin Caballero, Paul M. Finglas, Fidel Toldrá, *Encyclopedia of Food and Health*, Academic Press, pp. 646-652. <https://doi.org/10.1016/B978-0-12-384947-2.00596-1>.

- Paques FW, Macedo GA., (2006). Lipases de Látex Vegetais: Propriedades e Aplicações Industriais: A Review. *Química Nova*. 29(1):93.
- Praengam, K., Sahasakul, Y., Kupradinun, P., Sakarin, S., Sanitchua, W., Rungsipipat, A., Rattanapinyopituk, K., Angkasekwinai, P., Changsri, K., Mhuantong, W., Tangphatsornruang, S., & Tuntipopipat, S. (2017). Brown rice and retrograded brown rice alleviate inflammatory response in dextran sulfate sodium (DSS)-induced colitis mice. *Food & function*, 8(12), 4630–4643. <https://doi.org/10.1039/c7fo00305f>.
- Pranoto, H. S., W. Q. Mugnisjah., E. Muniarti. (1990). *Biologi Benih*. IPB. Bogor
- Pradeep, P.M., dan Sreerama, Y.N., (2015). Impact of processing on the phenolic profiles of small millets: evaluation of their antioxidant and enzyme inhibitory properties associated with hyperglycemia. *Food Chem*. 169, 455–463.
- P. J. Caceres, Martinez V. M., Amigo, L., dan Frias, J. (2014). Maximizing the phytochemical content and antioxidant activity of ecuadorian brown rice sprout through optimal germination conditions. *Food Chemistry*, 152: 407-414.
- Quettier AL, Eastmond PJ, (2009). Storage oil hydrolysis during early seedling growth. *Plant Physiology and Biochemistry*.47:485.
- Reische, D. W., Lillard, D. A., dan Eitenmiller, D. (2002). Antioxidant. Dalam: Casmir C. Akoh dan David B. Min. *Food Lipids: Chemistry, Nutrition, and Biotechnology*. Marcel Dekker, Inc. New York.
- Silalahi, J. (2006). Makanan Fungsional. *Kanisius*. Yogyakarta.

- Shutov AD, Vaintraub IA., (1987). Degradation of storage proteins in germinating seeds. *Phytochemistry*.26:1557-1566.
- Srisang, N., S. Prachayawarakorn, S. Soponronnarit, and W. Varanyanond. (2009). Comparative study of germinated brown rice drying using hot air and superheated steam fluidized bed. *In Proc. 6th Asia-Pacific Drying Conf.* 336-342. Bangkok, Thailand: King Mongkut's University of Technology Thonburi and Thammasat University.
- Sutopo, L. (2002). *Teknologi Benih Ed. Revisi*. PT Raja Grafindo Persada. Jakarta.
- Sugiyono. (2018). *Metode Penelitian Kuantitatif*. Bandung: Alfabeta.
- Vermerris, W., dan Nicholson, R. (2008). *Phenolic Compound Biochemistry*. Springer. Berlin.
- Winarno, F. G. (1997). *Kimia Pangan dan Gizi*. Gramedia Pustaka Utama. Jakarta.
- Winarsi, H. (2007). *Antioksidan Alami dan Radikal Bebas*. Penerbit Kanisius. Yogyakarta.
- Wu, F. F., Yang, N., Tour´e, A., Jin, Z. Y., Xu, X. M. (2013). Germinated brown rice and its role in human health. *Critical Reviews in Food Science and Nutrition*, 53(5), 451- 463. <https://doi:10.1080/10408398.2010.542259>.
- Wu, N. N., Li, R., Li, Z. J., Tan, B. (2022). Effect of germination in the form of paddy rice and brown rice on their phytic acid, GABA, γ -oryzanol, phenolics, flavonoids and antioxidant capacity. *Food Research International*, vol. 159. <https://doi.org/10.1016/j.foodres.2022.111603>.

- Wu, X., Guo, T., Luo, F., Lin, Q. (2023). Brown rice: a missing nutrient-rich health food. *Food Science and Human Wellness*, Vol. 12(5), pp. 1458-1470. <https://doi.org/10.1016/j.fshw.2023.02.010>.
- Yodpitaka, S., Mahatheeranont, S., Boonyawan, D., Sookwong, P., Roytrakul, S., Norkaew, O. (2019). Cold plasma treatment to improve germination and enhance the bioactive phytochemical content of germinated brown rice. *Food Chemistry*, 289, 328–339. <https://doi.org/10.1016/j.foodchem.2019.03.061>.
- Yue, X. dan Xu, Z. (2008). Changes of Anthocyanins, Anthocyanidins, and antioxidant activity compositions in black rice. *Agricultural Sciences in China*, 5(6): 431-440.
- Yulianthi, N. N., Suhendra, L., & Wrasiasi, L. (2017). Pengaruh perbandingan jenis pelarut terhadap kandungan senyawa total fenolik, α -Tokoferol, dan total karotenoid ekstrak *Sargassum polycystum*. *Jurnal Rekayasa Dan Manajemen Agroindustri*, 5(4), 1–10.
- Zielinski, H., Kozłowska, H., Lewczuk, B. (2001). Bioactive compounds in the cereal grains before and after hydrothermal processing. *Innovative Food Science & Emerging Technologies*, Vol. 2(3), pp. 159-169, [https://doi.org/10.1016/S1466-8564\(01\)00040-6](https://doi.org/10.1016/S1466-8564(01)00040-6).