

## DAFTAR PUSTAKA

- Almeida, H., Amaral, M. H., Lobão, P., & Lobo, J. M. (2014). In situ gelling systems: a strategy to improve the bioavailability of ophthalmic pharmaceutical formulations. *Drug discovery today*, 19(4), 400–412. <https://doi.org/10.1016/j.drudis.2013.10.001>
- AOAC (1993) *Official Methods of Analysis, Journal of AOAC International*.
- AOAC (2019). Apendix F: Guidelines for Standar Method Performance Requierements, in *AOAC Official Methods of Analysis*. AOAC International, pp. 1–18.
- Aranaz, I., Alcántara, A. R., Civera, M. C., Arias, C., Elorza, B., Heras Caballero, A., & Acosta, N. (2021). Chitosan: An Overview of Its Properties and Applications. *Polymers*, 13(19), 3256. <https://doi.org/10.3390/polym13193256>.
- Austin, A., Lietman, T. and Rose-Nussbaumer, J. (2017) “Update on the Management of Infectious Keratitis,” *Ophthalmology*, 124(11), pp. 1678–1689. doi: 10.1016/j.ophtha.2017.05.012.
- Byrd, L. B. and Martin, N. (2021) *Corneal Ulcer*. Treasure Island (FL): StatPearls Publishing.
- Cao, Y., Zhang, W., Wu, J., Zhang, H., & Zhou, H. (2017). Peripheral Ulcerative Keratitis Associated with Autoimmune Disease: Pathogenesis and Treatment. *Journal of ophthalmology*, 2017, 7298026. <https://doi.org/10.1155/2017/7298026>.
- Chaudhari, P., Shetty, D. and Lewis, S. A. (2022). Recent progress in colloidal nanocarriers loaded in situ gel in ocular therapeutics, *Journal of Drug Delivery Science and Technology*, 71(April). doi: 10.1016/j.jddst.2022.103327.
- Chaudhari, S. P. and Dugar, R. P. (2017). Application of surfactants in solid dispersion technology for improving solubility of poorly water soluble drugs, *Journal of Drug Delivery Science and Technology*, 41, pp. 68–77. doi: 10.1016/j.jddst.2017.06.010.
- Cheng, M., Cui, Y., Guo, Y., Zhao, P., Wang, J., Zhang, R., & Wang, X. (2023). Design of carboxymethyl chitosan-reinforced pH-responsive hydrogels for on-demand release of carvacrol and simulation of release kinetics. *Food chemistry*, 405(Pt A), 134856. <https://doi.org/10.1016/j.foodchem.2022.134856>.
- Cho, H. J., Balakrishnan, P., Park, E. K., Song, K. W., Hong, S. S., Jang, T. Y., Kim, K. S., Chung, S. J., Shim, C. K., & Kim, D. D. (2011). Poloxamer/cyclodextrin/chitosan-based thermoreversible gel for intranasal delivery of fexofenadine hydrochloride. *Journal of pharmaceutical sciences*, 100(2), 681–691. <https://doi.org/10.1002/jps.22314>.
- Christine, R. N. (2018) “Ulkus Kornea dengan Penyebab Bakteri; Sebuah Laporan Kasus,” in, pp. 63–69.
- Colin, J., Simonpoli, S., Geldsetzer, K., & Ropo, A. (2003). Corneal penetration of levofloxacin into the human aqueous humour: a comparison with

- ciprofloxacin. *Acta ophthalmologica Scandinavica*, 81(6), 611–613. <https://doi.org/10.1111/j.1395-3907.2003.00173x>.
- Farahani, M., Patel, R. and Dwarakanathan, S. (2017). Infectious corneal ulcers, *Disease-a-Month*, 63(2), pp. 33–37.
- Farida, Y. (2015) “Corneal Ulcers Treatment,” *Corneal Ulcers Treatment J MAJORITY* |, 4(1), p. 119.
- Fish, D. N. and Chow, A. T. (1997) “The clinical pharmacokinetics of escitalopram,” *Clinical Pharmacokinetics*, 46(4), pp. 281–290. doi: 10.2165/00003088-200746040-00002.
- Gilani, C. J., Yang, A., Yonkers, M., & Boysen-Osborn, M. (2017). Differentiating Urgent and Emergent Causes of Acute Red Eye for the Emergency Physician. *The western journal of emergency medicine*, 18(3), 509–517. <https://doi.org/10.5811/westjem.2016.12.31798>.
- Gratieri, T., Gelfuso, G. M., Rocha, E. M., Sarmiento, V. H., de Freitas, O., & Lopez, R. F. (2010). A poloxamer/chitosan in situ forming gel with prolonged retention time for ocular delivery. *European journal of pharmaceutics and biopharmaceutics : official journal of Arbeitsgemeinschaft fur Pharmazeutische Verfahrenstechnik e.V*, 75(2), 186–193.
- Gupta, H., Aqil, M., Khar, R. K., Ali, A., Bhatnagar, A., & Mittal, G. (2015). An alternative in situ gel-formulation of levofloxacin eye drops for prolong ocular retention. *Journal of pharmacy & bioallied sciences*, 7(1), 9–14. <https://doi.org/10.4103/0975-7406.149810>.
- Gupta, H., Velpandian, T. and Jain, S. (2010). Ion-and pH-activated novel in-situ gel system for sustained ocular drug delivery. *Journal of Drug Targeting*, 18(7), pp. 499–505. doi: 10.3109/10611860903508788.
- Irimia, T., Dinu-Pîrvu, C. E., Ghica, M. V., Lupuleasa, D., Muntean, D. L., Udeanu, D. I., & Popa, L. (2018). Chitosan-Based In Situ Gels for Ocular Delivery of Therapeutics: A State-of-the-Art Review. *Marine drugs*, 16(10), 373. <https://doi.org/10.3390/md16100373>.
- JH Collett (2005). Poloxamer. In *Handbook of Pharmaceutical Excipient*, pp. 506–509.
- Kadam, A. T. (2017). Design and Evaluation of Modified Chitosan Based In Situ Gel For Ocular Drug Delivery. *International Journal of Pharmaceutical Sciences*, 9(11), pp. 87–91.
- Kanda, Y., Kayama, T., Okamoto, S., Hashimoto, M., Ishida, C., Yanai, T., Fukumoto, M., & Kunihiro, E. (2012). Post-marketing surveillance of levofloxacin 0.5% ophthalmic solution for external ocular infections. *Drugs in R&D*, 12(4), 177–185. <https://doi.org/10.2165/11636020-000000000-00000>.
- Kashikar, V. S. and Gonjari, I. D. (2014) “In situ gelling systems of ofloxacin: Comparative performance of in vivo precorneal drainage and pharmacokinetic study,” *Asian Journal of Pharmaceutics*, 7(1). Available at: <https://doi.org/10.22377/ajp.v7i1.35>.
- Kemenkes (2013) *Riset Kesehatan Dasar*. doi: 1 Desember 2013.
- Kemenkes (2020) *FARMAKOPE INDONESIA EDISI VI*.
- Khattab, A., Marzok, S. and Ibrahim, M. (2019). Development of optimized mucoadhesive thermosensitive pluronic based in situ gel for controlled

- delivery of Latanoprost: Antiglaucoma efficacy and stability approaches. *Journal of Drug Delivery Science and Technology*, 53(June). doi: 10.1016/j.jddst.2019.101134.
- King, D. E., Malone, R. and Lilley, S. H. (2000). New classification and update on the quinolone antibiotics. *American Family Physician*, 61(9), pp. 2741–2748.
- Kolawole, O. M. and Cook, M. T. (2023). In situ gelling drug delivery systems for topical drug delivery. *European Journal of Pharmaceutics and Biopharmaceutics*, 184(November 2022), pp. 36–49. doi: 10.1016/j.ejpb.2023.01.007.
- Ladas, J. G. and Mondino, B. J. (2000). Systemic disorders associated with peripheral corneal ulceration. *Current Opinion in Ophthalmology*, 11(6), pp. 468–471. doi: 10.1097/00055735-200012000-00014.
- Lin, A., Rhee, M. K., Akpek, E. K., Amescua, G., Farid, M., Garcia-Ferrer, F. J., Varu, D. M., Musch, D. C., Dunn, S. P., Mah, F. S., & American Academy of Ophthalmology Preferred Practice Pattern Cornea and External Disease Panel (2019). Bacterial Keratitis Preferred Practice Pattern®. *Ophthalmology*, 126(1), P1–P55.
- Maharini, I. (2021). Validation uv spectrophotometric method to determine entrapment efficiency of ocular polymeric nanoparticle levofloxacin hemihydrate. *Research Journal of Pharmacy and Technology*, 14(10), pp. 5479–5482. doi: 10.52711/0974-360X.2021.00956.
- Mandal, S., Thimmasetty, M. K., Prabhushankar, G., & Geetha, M. (2012). Formulation and evaluation of an in situ gel-forming ophthalmic formulation of moxifloxacin hydrochloride. *International journal of pharmaceutical investigation*, 2(2), 78–82. <https://doi.org/10.4103/2230-973X.100042>.
- Mohanty, D., Bakshi., Simharaju, Nandini., Haque, M Akiful., Sahoo, Chinmaya Keshari. (2018). A Review on in situ Gel: A Novel Drug Delivery System. *int.J.Pharm.Sci.Rev.Res*, 50(0976), pp. 175–181. Available at: <https://www.researchgate.net/publication/326246294>.
- Noreddin, A. M., Elkhatab, W. F., Cunnion, K. M., & Zhanel, G. G. (2011). Cumulative clinical experience from over a decade of use of levofloxacin in community-acquired pneumonia: critical appraisal and role in therapy. *Drug, healthcare and patient safety*, 3, 59–68. <https://doi.org/10.2147/DHPS.S15599>
- Permana, A. D., Utami, R. N., Layadi, P., Himawan, A., Juniarti, N., Anjani, Q. K., Utomo, E., Mardikasari, S. A., Arjuna, A., & Donnelly, R. F. (2021). Thermosensitive and mucoadhesive in situ ocular gel for effective local delivery and antifungal activity of itraconazole nanocrystal in the treatment of fungal keratitis. *International journal of pharmaceutics*, 602, 120623. <https://doi.org/10.1016/j.ijpharm.2021.120623>.
- Ren, Y., Qi, X.-J. and Cui, Y. (2015). Preparation of a Thermosensitive Guanidine-Chitosan/Poloxamer in situ Gel for Intranasal Delivery System. *Proceedings of the 5th International Conference on Information Engineering for Mechanics and Materials*, 21(Icimm), pp. 686–689. doi:

10.2991/icimm-15.2015.129.

- Sabra, R. and Billa, N. (2020). Gastrointestinal Delivery of APIs from Chitosan Nanoparticles. *IntechOpen*, p. 13. Available at: <http://dx.doi.org/10.1039/C7RA00172J%0Ahttps://www.intechopen.com/books/advanced-biometric-technologies/liveness-detection-in-biometrics%0Ahttp://dx.doi.org/10.1016/j.colsurfa.2011.12.014>.
- Sakai, T., Shinno, K., Kurata, M., & Kawamura, A. (2019). Pharmacokinetics of Azithromycin, Levofloxacin, and Ofloxacin in Rabbit Extraocular Tissues After Ophthalmic Administration. *Ophthalmology and therapy*, 8(4), 511–517. <https://doi.org/10.1007/s40123-019-00205-0>.
- Shelke, S., Shahi, Sadhana., Jalalpura, Sunil., Dhamecha, Dinesh., Shengule, Sushant. (2016). Formulation and evaluation of thermoreversible mucoadhesive in situ gel for intranasal delivery of Naratriptan hydrochloride. *Journal of Drug Delivery Science and Technology*.
- Sherif, A. Y., Mahrous, G. M. and Alanazi, F. K. (2018). Novel in-situ gel for intravesical administration of ketorolac. *Saudi Pharmaceutical Journal*, 26(6), pp. 845–851. doi: 10.1016/j.jsps.2018.03.014.
- Sheshala, R., Wai, N. Z., Said, I. D., Ashraf, K., Lim, S. M., Ramasamy, K., & Zeeshan, F. (2022). Poloxamer and Chitosan-Based In Situ Gels Loaded with Orthosiphon stamineus Benth. Extracts Containing Rosmarinic Acid for the Treatment of Ocular Infections. *Turkish journal of pharmaceutical sciences*, 19(6), 671–680. <https://doi.org/10.4274/tjps.galenos.2021.40121>
- Soliman, K. A. *et al.* (2019) “Poloxamer-based in situ gelling thermoresponsive systems for ocular drug delivery applications,” *Drug Discovery Today*, 24(8), pp. 1575–1586.
- Ur-Rehman, T., Tavelin, S., & Gröbner, G. (2011). Chitosan in situ gelation for improved drug loading and retention in poloxamer 407 gels. *International journal of pharmaceutics*, 409(1-2), 19–29.
- Üstündağ Okur, N., Yozgatlı, V., Okur, M., Yoltaş, A., Siafaka, P. (2019). Improving therapeutic efficacy of voriconazole against fungal keratitis: Thermo-sensitive in situ gels as ophthalmic drug carriers. *Journal of Drug Delivery Science and Technology*, 49(September 2018), pp. 323–333. doi: 10.1016/j.jddst.2018.12.005.
- Varshosaz, J., Tabbakhian, M. and Salmani, Z. (2008) “Designing of a Thermosensitive Chitosan/Poloxamer In Situ Gel for Ocular Delivery of Ciprofloxacin,” *The Open Drug Delivery Journal*, 2(1), pp. 61–70. doi: 10.2174/1874126600802010061.
- Verekar, R. R., Gurav, S. S. and Bolmal, U. (2020) “Thermosensitive mucoadhesive in situ gel for intranasal delivery of Almotriptan malate: Formulation, characterization, and evaluation,” *Journal of Drug Delivery Science and Technology*, 58(December 2019). doi: 10.1016/j.jddst.2020.101778.
- Vigani, B., Rossi, S., Sandri, G., Bonferoni, M. C., Caramella, C. M., & Ferrari, F. (2020). Recent Advances in the Development of In Situ Gelling Drug Delivery Systems for Non-Parenteral Administration Routes. *Pharmaceutics*, 12(9), 859.