

## DAFTAR PUSTAKA

- Al-Hussaini, A. *et al.* (2022) 'The Epidemiology and Outcome of Biliary Atresia: Saudi Arabian National Study (2000–2018)', *Frontiers in Pediatrics*, 10. Available at: <https://www.frontiersin.org/articles/10.3389/fped.2022.921948> (Accessed: 21 November 2022).
- Apostu, R.-C. *et al.* (2020) 'Biological and Histological Assessment of the Hepatoportoenterostomy Role in Biliary Atresia as a Stand-Alone Procedure or as a Bridge toward Liver Transplantation', *Medicina*, 57(1), p. 16. Available at: <https://doi.org/10.3390/medicina57010016>.
- Averbukh, L.D. and Wu, G.Y. (2018) 'Evidence for Viral Induction of Biliary Atresia: A Review', *Journal of Clinical and Translational Hepatology*, 6(4), pp. 1–10. Available at: <https://doi.org/10.14218/JCTH.2018.00046>.
- Baek, S.H. *et al.* (2020) 'The Epidemiology and Etiology of Cholangitis After Kasai Portoenterostomy in Patients With Biliary Atresia', *Journal of Pediatric Gastroenterology and Nutrition*, 70(2), pp. 171–177. Available at: <https://doi.org/10.1097/MPG.0000000000002555>.
- Betalli, P. and Davenport, M. (2019) 'Biliary Atresia and Other Congenital Disorders of the Extrahepatic Biliary Tree', in L. D'Antiga (ed.) *Pediatric Hepatology and Liver Transplantation*. Cham: Springer International Publishing, pp. 129–144. Available at: [https://doi.org/10.1007/978-3-319-96400-3\\_7](https://doi.org/10.1007/978-3-319-96400-3_7).
- Bijl, E.J. *et al.* (2013) 'The long-term outcome of the Kasai operation in patients with biliary atresia: a systematic review', 71(4), p. 4.
- Brahee, D.D. and Lampl, B.S. (2021) 'Neonatal diagnosis of biliary atresia: a practical review and update', *Pediatric Radiology* [Preprint]. Available at: <https://doi.org/10.1007/s00247-021-05148-y>.
- Calinescu, A.M. *et al.* (2022) 'Cholangitis Definition and Treatment after Kasai Hepatoportoenterostomy for Biliary Atresia: A Delphi Process and International Expert Panel', *Journal of Clinical Medicine*, 11(3), p. 494. Available at: <https://doi.org/10.3390/jcm11030494>.
- Capparelli, M.A. *et al.* (2021) 'Preoperative risk factors for the early failure of the Kasai portoenterostomy in patients with biliary atresia', *Pediatric Surgery International*, 37(9), pp. 1183–1189. Available at: <https://doi.org/10.1007/s00383-021-04933-y>.

Cavallo, L. *et al.* (2022) ‘The Epidemiology of Biliary Atresia: Exploring the Role of Developmental Factors on Birth Prevalence’, *The Journal of Pediatrics*, 246, pp. 89-94.e2. Available at: <https://doi.org/10.1016/j.jpeds.2022.03.038>.

Chen, S.-Y. *et al.* (2018) ‘Number of cholangitis episodes as a prognostic marker to predict timing of liver transplantation in biliary atresia patients after Kasai portoenterostomy’, *BMC pediatrics*, 18(1), p. 119. Available at: <https://doi.org/10.1186/s12887-018-1074-2>.

Chung, P.H.Y., Wong, K.K.Y. and Tam, P.K.H. (2015) ‘Predictors for failure after Kasai operation’, *Journal of Pediatric Surgery*, 50(2), pp. 293–296. Available at: <https://doi.org/10.1016/j.jpedsurg.2014.11.015>.

Davenport, M. (2017) ‘Adjuvant therapy in biliary atresia: hopelessly optimistic or potential for change?’, *Pediatric Surgery International*, 33(12), pp. 1263–1273. Available at: <https://doi.org/10.1007/s00383-017-4157-5>.

Dhamnetiya, D. *et al.* (2022) ‘How to Analyze the Diagnostic Performance of a New Test? Explained with Illustrations’, *Journal of Laboratory Physicians*, 14(01), pp. 090–098. Available at: <https://doi.org/10.1055/s-0041-1734019>.

Fawaz, R. *et al.* (2017) ‘Guideline for the Evaluation of Cholestatic Jaundice in Infants: Joint Recommendations of the North American Society for Pediatric Gastroenterology, Hepatology, and Nutrition and the European Society for Pediatric Gastroenterology, Hepatology, and Nutrition’, *Journal of Pediatric Gastroenterology and Nutrition*, 64(1), pp. 154–168. Available at: <https://doi.org/10.1097/MPG.0000000000001334>.

Feldman, A. and Mack, C.L. (2012) ‘Biliary Atresia: Cellular Dynamics and Immune Dysregulation’, *Seminars in Pediatric Surgery*, 21(3), pp. 192–200. Available at: <https://doi.org/10.1053/j.sempedsurg.2012.05.003>.

Fischler, B., Svensson, J.F. and Nemeth, A. (2009) ‘Early cytomegalovirus infection and the long-term outcome of biliary atresia’, *Acta Paediatrica (Oslo, Norway: 1992)*, 98(10), pp. 1600–1602. Available at: <https://doi.org/10.1111/j.1651-2227.2009.01416.x>.

Gad, E.H. *et al.* (2021) ‘Short- and long-term outcomes after Kasai operation for type III biliary atresia: Twenty years of experience in a single tertiary Egyptian center-A retrospective cohort study’, *Annals of Medicine and Surgery*, 62, pp. 302–314. Available at: <https://doi.org/10.1016/j.amsu.2021.01.052>.

Gunadi *et al.* (2018) ‘Liver transplant score for prediction of biliary atresia patients’ survival following Kasai procedure’, *BMC Research Notes*, 11(1), p. 381. Available at: <https://doi.org/10.1186/s13104-018-3498-z>.

- Gunadi *et al.* (2020) 'Histopathological findings for prediction of liver cirrhosis and survival in biliary atresia patients after Kasai procedure', *Diagnostic Pathology*, 15(1), p. 79. Available at: <https://doi.org/10.1186/s13000-020-00996-y>.
- Hahn, S.M. *et al.* (2013) 'Clinical Benefit of Liver Stiffness Measurement at 3 Months after Kasai Hepatopertoenterostomy to Predict the Liver Related Events in Biliary Atresia', *PLoS ONE*, 8(11), p. e80652. Available at: <https://doi.org/10.1371/journal.pone.0080652>.
- Hanalioglu, D. *et al.* (2019) 'Revisiting long-term prognostic factors of biliary atresia: A 20-year experience with 81 patients from a single center', *The Turkish Journal of Gastroenterology*, 30(5), pp. 467–474. Available at: <https://doi.org/10.5152/tjg.2019.18590>.
- Hill, R. *et al.* (2015) 'Th-17 cells infiltrate the liver in human biliary atresia and are related to surgical outcome', *Journal of Pediatric Surgery*, 50(8), pp. 1297–1303. Available at: <https://doi.org/10.1016/j.jpedsurg.2015.02.005>.
- Huang, C.-Y. *et al.* (2020) 'Bilirubin level 1 week after hepatopertoenterostomy predicts native liver survival in biliary atresia', *Pediatric Research*, 87(4), pp. 730–734. Available at: <https://doi.org/10.1038/s41390-019-0610-6>.
- Khayat, A., Alamri, A.M. and Saadah, O.I. (2021) 'Outcomes of late Kasai portoenterostomy in biliary atresia: a single-center experience', *The Journal of International Medical Research*, 49(5), p. 03000605211012596. Available at: <https://doi.org/10.1177/03000605211012596>.
- Kiriyama, S. *et al.* (2018) 'Tokyo Guidelines 2018: diagnostic criteria and severity grading of acute cholangitis (with videos)', *Journal of Hepato-Biliary-Pancreatic Sciences*, 25(1), pp. 17–30. Available at: <https://doi.org/10.1002/jhbp.512>.
- Kodo, K. *et al.* (2018) 'Cytomegalovirus-associated biliary atresia', *Journal of Pediatric Surgery Case Reports*, 35, pp. 17–20. Available at: <https://doi.org/10.1016/j.epsc.2018.02.002>.
- Lakshminarayanan, B. and Davenport, M. (2016) 'Biliary atresia: A comprehensive review', *Journal of Autoimmunity*, 73, pp. 1–9. Available at: <https://doi.org/10.1016/j.jaut.2016.06.005>.
- Lianti, H. *et al.* (2020) 'Analysis of knowledge regarding Biliary Atresia among healthcare providers and laypersons in East Jakarta after educational intervention', *ASEAN Journal of Community Engagement*, 4(1). Available at: <https://doi.org/10.7454/ajce.v4i1.1053>.

Liu, J. *et al.* (2019) ‘Risk factors and prognostic effects of cholangitis after Kasai procedure in biliary atresia patients: A retrospective clinical study’, *Journal of Pediatric Surgery*, 54(12), pp. 2559–2564. Available at: <https://doi.org/10.1016/j.jpedsurg.2019.08.026>.

Marsico, C. and Kimberlin, D.W. (2017) ‘Congenital Cytomegalovirus infection: advances and challenges in diagnosis, prevention and treatment’, *Italian Journal of Pediatrics*, 43(1), p. 38. Available at: <https://doi.org/10.1186/s13052-017-0358-8>.

McPherson, R.A. and Pincus, M.R. (eds) (2017) *Henry’s clinical diagnosis and management by laboratory methods*. 23rd edition. St. Louis, Missouri: Elsevier.

Moore, K.L., Persaud, T.V.N. and Torchia, M.G. (2013) *The developing human: clinically oriented embryology*. 9th ed. Philadelphia, PA: Saunders/Elsevier.

Nakajima, H. *et al.* (2018) ‘Does time taken to achieve jaundice-clearance influence survival of the native liver in post-Kasai biliary atresia?’, *World Journal of Pediatrics*, 14(2), pp. 191–196. Available at: <https://doi.org/10.1007/s12519-018-0139-5>.

Qisthi, S.A. *et al.* (2020) ‘Prognostic Factors for Survival of Patients with Biliary Atresia Following Kasai Surgery’, *The Kobe Journal of Medical Sciences*, 66(2), pp. E56–E60.

Ramachandran, P. *et al.* (2019) ‘Early Cholangitis after Portoenterostomy in Children with Biliary Atresia’, *Journal of Indian Association of Pediatric Surgeons*, 24(3), pp. 185–188. Available at: [https://doi.org/10.4103/jiaps.JIAPS\\_96\\_18](https://doi.org/10.4103/jiaps.JIAPS_96_18).

Ross, S.A. *et al.* (2011) ‘Diagnosis of Cytomegalovirus Infections’, *Infectious disorders drug targets*, 11(5), pp. 466–474.

Shehata, S. *et al.* (2018) ‘Prognostic factors for the outcome of Kasai portoenterostomy for infants with biliary atresia in Egypt’, *Alexandria Journal of Pediatrics*, 31(3), p. 112. Available at: [https://doi.org/10.4103/AJOP.AJOP\\_25\\_18](https://doi.org/10.4103/AJOP.AJOP_25_18).

Shen, C. *et al.* (2008) ‘Relationship between prognosis of biliary atresia and infection of cytomegalovirus’, *World journal of pediatrics: WJP*, 4(2), pp. 123–126. Available at: <https://doi.org/10.1007/s12519-008-0024-8>.

Shneider, B.L. *et al.* (2016) ‘Total Serum Bilirubin within Three Months of Hepatportoenterostomy Predicts Short-term Outcomes in Biliary Atresia’, *The Journal of pediatrics*, 170, pp. 211-217.e2. Available at: <https://doi.org/10.1016/j.jpeds.2015.11.058>.



Song, Z. *et al.* (2017) ‘Surgical outcome and etiologic heterogeneity of infants with biliary atresia who received Kasai operation less than 60 days after birth: A retrospective study’, *Medicine*, 96(26), p. e7267. Available at: <https://doi.org/10.1097/MD.00000000000007267>.

Vij, M. and Rela, M. (2020) ‘Biliary atresia: pathology, etiology and pathogenesis’, *Future Science OA*, 6(5), p. FSO466. Available at: <https://doi.org/10.2144/fsoa-2019-0153>.

Xu, Y. *et al.* (2012) ‘The Perinatal Infection of Cytomegalovirus Is an Important Etiology for Biliary Atresia in China’, *Clinical Pediatrics*, 51(2), pp. 109–113. Available at: <https://doi.org/10.1177/0009922811406264>.

Youden, W.J. (1950) ‘Index for rating diagnostic tests’, *Cancer*, 3(1), pp. 32–35. Available at: [https://doi.org/10.1002/1097-0142\(1950\)3:1<32::AID-CNCR2820030106>3.0.CO;2-3](https://doi.org/10.1002/1097-0142(1950)3:1<32::AID-CNCR2820030106>3.0.CO;2-3).

Zani, A. *et al.* (2015) ‘Cytomegalovirus-associated biliary atresia: An aetiological and prognostic subgroup’, *Journal of Pediatric Surgery*, 50(10), pp. 1739–1745. Available at: <https://doi.org/10.1016/j.jpedsurg.2015.03.001>.

Zhang, Y. *et al.* (2022) ‘A Novel Model for Predicting the Clearance of Jaundice in Patients With Biliary Atresia After Kasai Procedure’, *Frontiers in Pediatrics*, 10, p. 837247. Available at: <https://doi.org/10.3389/fped.2022.837247>.

Zhao, Y. *et al.* (2021) ‘Prognosis of Biliary Atresia Associated With Cytomegalovirus: A Meta-Analysis’, *Frontiers in Pediatrics*, 9. Available at: <https://www.frontiersin.org/article/10.3389/fped.2021.710450> (Accessed: 31 January 2022).