

## DAFTAR PUSTAKA

- Åberg, K. (2017). Neonatal complications following birth by vacuum extraction. Department of Womens and Childeern Health.
- Aslam, H. Muhammad, Saleem, S., Afzal, R., Iqbal, U., Saleem, S. M. uhammad, Shaikh, M. W. aqas A., & Shahid, N. (2014). "Risk factors of birth asphyxia." *Italian Journal of Pediatrics*, 40, 94. <https://doi.org/10.1186/s13052-014-0094-2>
- Bazzano, A. N., Var, C., Wilkosz, D., Duggal, R., & Oberhelman, R. A. (2019). Neonatal deaths in Cambodia: Findings from a community-based mortality review. In *BMC Research Notes* (Vol. 12, Issue 1). BioMed Central Ltd. <https://doi.org/10.1186/s13104-019-4265-5>
- Bello, F. A., Tsele, T. A., & Oluwasola, T. O. (2015). Decision-to-delivery intervals and perinatal outcomes following emergency cesarean delivery in a Nigerian tertiary hospital. *International Journal of Gynecology and Obstetrics*, 130(3), 279–283. <https://doi.org/10.1016/j.ijgo.2015.03.036>
- Berghella, V., Baxter, J. K., & Chauhan, S. P. (2005). Evidence-based surgery for cesarean delivery. In *American Journal of Obstetrics and Gynecology* (Vol. 193, Issue 5, pp. 1607–1617). <https://doi.org/10.1016/j.ajog.2005.03.063>
- Bogdanovic, G., Babovic, A., Rizvanovic, M., Ljuca, D., Grgic, G., & Djuranovic-Milicic, J. (2014). Cardiotocography in the prognosis of perinatal outcome. *Medical Archives (Sarajevo, Bosnia and Herzegovina)*, 68(2), 102–105. <https://doi.org/10.5455/medarh.2014.68.102-105>
- Brandt, J.A., Morgenstern, B., Thangarajah, F., Guttner, B., Ludwig, S., Eichler, C., Ratiu, J., Mallmann, P., Ratiu, D. (2020). Evaluating the Decision-to-Delivery Interval in Emergency Cesarean Sections and its Impact on Neonatal Outcome. *in vivo* 34: 3341-3347 (2020)
- Brucknerova I, Ujhazy E. (2014). Asphyxia in newborn-risk, prevention and identification of a hypoxic event. *Neuro Endocrinol. Lett*, 35 (2): 201-210.
- Chaudhari, S. N., Bhikane, D. B., & Gupta, P. (2017). A clinical study of postdated pregnancy. *International Journal of Reproduction, Contraception, Obstetrics and Gynecology*, 6(5), 2077. <https://doi.org/10.18203/2320-1770.ijrcog20171980>
- Chukwudi, O. E., & Okonkwo, C. A. (2014). Decision - Delivery interval and perinatal outcome of emergency caesarean sections at a tertiary institution. *Pakistan Journal of Medical Sciences*, 30(5). <https://doi.org/10.12669/pjms.305.5470>
- Cunningham et al. (2018). *Williams Obstetrics*. McGraw Hill Education. 25 Th Edition (30) : 1247
- Dur-e-Shahwar, Ahmed, I., Amerjee, A., & Hoodbhoy, Z. (2018). Comparison of neonatal outcomes between category-1 & non-category-1 primary emergency cesarean section: A retrospective record review in a tertiary care hospital. *Pakistan Journal of Medical Sciences*, 34(4), 823–827. <https://doi.org/10.12669/pjms.344.14496>
- Edipoglu, I. S., Celik, F., Marangoz, E. C., & Orcan, G. H. (2018). Effect of anaesthetic technique on neonatal morbidity in emergency caesarean section

- for foetal distress. *PLoS ONE*, 13(11).  
<https://doi.org/10.1371/journal.pone.0207388>
- Ekwochi, U., Asinobi, N. I., Osuorah, C. DI, Ndu, I. K., Ifediora, C., Amadi, O. F., Iheji, C. C., Orjioke, C. J., Okenwa, W. O., & Okeke, B. I. (2017). Incidence and Predictors of Mortality Among Newborns With Perinatal Asphyxia: A 4-Year Prospective Study of Newborns Delivered in Health Care Facilities in Enugu, South-East Nigeria. *Clinical Medicine Insights: Pediatrics*, 11, 117955651774664. <https://doi.org/10.1177/1179556517746646>
- Endale, T., Fentahun, N., Gemada, D., & Hussien, M. A. (2016). Maternal and fetal outcomes in term premature rupture of membrane. *World Journal of Emergency Medicine*, 7(2), 147. <https://doi.org/10.5847/wjem.j.1920-8642.2016.02.011>
- Grivell, R. M., Alfirevic, Z., Gyte, G. M. L., & Devane, D. (2015). Antenatal cardiotocography for fetal assessment. In *Cochrane Database of Systematic Reviews* (Vol. 2015, Issue 9, pp. 1–39). John Wiley and Sons Ltd. <https://doi.org/10.1002/14651858.CD007863.pub4>
- Gunawan, T., Attamimi, A., & Pradjatmo, H. (2018). Hubungan Response Time Seksio Sesarea Emergensi Kategori 1 dengan Luaran Perinatal di RSUP Dr.Sardjito. *Jurnal Kesehatan Reproduksi*, 5(1), 60. <https://doi.org/10.22146/jkr.37997>
- Gupta, S., Naithani, U., Madhanmohan, C., Singh, A., Reddy, P., Gupta, A. (2017). Evaluation of decision-to-delivery interval in emergency cesarean section: A 1-year prospective audit in a tertiary care hospital. *J Anaesthesiol Clin Pharmacol*. 2017 Jan-Mar; 33(1): 64–70.
- Hirani, B. A., Mchome, B. L., Mazuguni, N. S., & Mahande, M. J. (2017). The decision delivery interval in emergency caesarean section and its associated maternal and fetal outcomes at a referral hospital in northern Tanzania: a cross-sectional study. *BMC Pregnancy and Childbirth*, 17(1), 411. <https://doi.org/10.1186/s12884-017-1608-x>
- Hosea, R. O. (2020). KARAKTERISTIK GAMBARAN FOTO TORAKS PADA PASIEN YANG DIRAWAT DI NEONATAL INTENSIVE CARE UNIT (NICU) RSUP Dr. WAHIDIN SUDIROHUSODO PERIODE 2018 – 2019. Tersedia di <http://repository.unhas.ac.id/id/eprint/970>. Diakses pada 21 Maret 2021
- Hussein AH. Emergency Caesarean Section Turnaround Time and Its Effect on Maternal and Newborn Health Outcomes At University of Nairobi Teaching Hospitals. 2012. Available at <http://erepository.uonbi.ac.ke:8080/xmlui/handle/123456789/13758>. (Accessed 22 Maret 2021)
- Jackson, M., Holmgren, C. M., Esplin, M.S., Henry, E., Varner, M. W. (2011). Frequency of fetal heart rate categories and short-term neonatal outcome. *Obstet Gynecol* 2011;118:803–8. <https://doi.org/10.1097/AOG.0b013e31822f1b50>
- Kamala, B. A., Mgaya, A. H., Ngarina, M. M., & Kidanto, H. L. (2018). Predictors of low birth weight and 24-hour perinatal outcomes at Muhimbili national hospital in Dar es Salaam, Tanzania: A five-year retrospective analysis of

- obstetric records. *Pan African Medical Journal*, 29. <https://doi.org/10.11604/pamj.2018.29.220.15247>
- Khemworapong, K., Sompagdee, N., & Boriboonthirunsarn, D. (2018). Decision-to-delivery interval in emergency cesarean delivery in tertiary care hospital in Thailand. *Obstetrics and Gynecology Science*, 61(1), 48–55. <https://doi.org/10.5468/ogs.2018.61.1.48>
- Kuper, S. G., Sievert, R. A., Steele, R., Biggio, J. R., Tita, A. T., & Harper, L. M. (2017). Maternal and neonatal outcomes in indicated preterm births based on the intended mode of delivery. *Obstetrics and Gynecology*, 130(5), 1143–1151. <https://doi.org/10.1097/AOG.0000000000002320>
- Lipman, S. S., Carvalho, B., Cohen, S. E., Druzin, M. L., & Daniels, K. (2013). Response times for emergency cesarean delivery: Use of simulation drills to assess and improve obstetric team performance. *Journal of Perinatology*, 33(4), 259–263. <https://doi.org/10.1038/jp.2012.98>
- Locatelli, A., Lambicchi, L., Incerti, M., Bonati, F., Ferdico, M., Malguzzi, S., Torcasio, F., Calzi, P., Varisco, T., & Paterlini, G. (2020). Is perinatal asphyxia predictable? *BMC Pregnancy and Childbirth*, 20(1). <https://doi.org/10.1186/s12884-020-02876-1>
- Macones, G. A., Hankins, G. D. V., Spong, C. Y., Hauth, J., & Moore, T. (2008). The 2008 National Institute of Child Health and Human Development Workshop Report on Electronic Fetal Monitoring Update on Definitions, Interpretation, and Research Guidelines. [www.greenjournal.org/cgi/content/full/112/](http://www.greenjournal.org/cgi/content/full/112/)
- Manuaba. (2014). Ilmu kebidanan penyakit kandungan dan keluarga berencana. Untuk pendidikan kebidanan. Jakarta EGC.
- Nair, V. V., Nair, S. S., & Venugopalan, P. (2019). Decision to delivery interval in emergency LSCS and its impact on fetal outcome. *International Journal of Reproduction, Contraception, Obstetrics and Gynecology*, 8(9), 3679. <https://doi.org/10.18203/2320-1770.ijrcog20193797>
- National Institute for Health and Clinical Excellence. CG132 Cesarean Section. Clinical Guidelines for Emergency CS. Available from: <http://www.guidance.nice.org.uk/>
- Notoatmodjo, Soekidjo. (2010). *Metodologi Riset Kesehatan*. Jakarta: Rineka Cipta.
- Palmer, E., Ciechanowicz, S., Reeve, A., Harris, S., Wong, D. J. N., & Sultan, P. (2018). Operating room-to-incision interval and neonatal outcome in emergency caesarean section: a retrospective 5-year cohort study. *Anaesthesia*, 73(7), 825–831. <https://doi.org/10.1111/anae.14296>
- Pearson, G. A., Kelly, B., Russell, R., Dutton, S., Kurinczuk, J. J., & MacKenzie, I. Z. (2011). Target decision to delivery intervals for emergency caesarean section based on neonatal outcomes and three year follow-up. *European Journal of Obstetrics and Gynecology and Reproductive Biology*, 159(2), 276–281. <https://doi.org/10.1016/j.ejogrb.2011.07.044>
- Pinto, V. L., Sharma, S. (2020). Continuous Positive Airway Pressure. Tersedia di <https://pubmed.ncbi.nlm.nih.gov/29489216/>. Diakses pada 22 maret 2021

- Plucinska, A., Hajduczenia, M., Pastusiak, M., Kowalik, M., Miechowicz, I., Szymankiewicz, M. (2010). The impact of premature rupture of membranes (PROM) on neonatal outcome. *Ginekol Pol.* 2010 Apr;81(4):277-82.
- Pramanik. (2020). Respiratory Distress Syndrome. Medscape. Tersedia di <https://emedicine.medscape.com/article/976034-overview>, diakses pada 21 Maret 2021
- Pratiwi, S. R., Prasetya, H., & Murti, B. (2021). The Effect of Asphyxia on Neonatal Death: A Meta-Analysis. *Journal of Maternal and Child Health*, 5(4), 413–421. <https://doi.org/10.26911/thejmch.2020.05.04.08>
- Preterm Labor and Preterm Birth [internet]. American College of Obstetricians and Gynecologists; 2013
- Purwaningsih, Y., Dewi, Y. L. R., Indarto, D., & Murti, B. (2018). Factors Associated with Newborn Asphyxia at Dr. Harjono Hospital, Ponorogo, East Java. *Journal of Maternal and Child Health*, 03(04), 287–293. <https://doi.org/10.26911/thejmch.2018.03.04.06>
- Qureshi, D. A., Rather, D. S., Muzaffar, D. U., & Shah, D. U. (2020). Role of Cardiotocography in Predicting Perinatal Outcome. *International Journal of Scientific and Research Publications (IJSRP)*, 10(05), 599–603. <https://doi.org/10.29322/ijsrp.10.05.2020.p10168>
- Radhakrishnan, G., Yadav, G., Vaid, N., & Ali, H. (2013). Factors affecting decision to delivery interval; in emergency caesarean sections in a tertiary care hospital: a cross sectional observational study. *International Journal of Reproduction, Contraception, Obstetrics and Gynecology*, 2(4), 651. <https://doi.org/10.5455/2320-1770.ijrcog20131231>
- Rocha, G., Soares, P., Gonçalves, A., Silva, A. I., Almeida, D., Figueiredo, S., Pissarra, S., Costa, S., Soares, H., Flôr-De-Lima, F., & Guimarães, H. (2018). Respiratory Care for the Ventilated Neonate. In *Canadian Respiratory Journal* (Vol. 2018). Hindawi Limited. <https://doi.org/10.1155/2018/7472964>
- Royal College of Obstetricians & Gynaecologists, the Royal College of Anaesthetists. Classification of urgency of caesarean section - a continuum of risk (Good Practice No. 11)
- Sayed Ahmed, W. A., & Hamdy, M. A. (2018). Optimal management of umbilical cord prolapse. In *International Journal of Women's Health* (Vol. 10, pp. 459–465). Dove Medical Press Ltd. <https://doi.org/10.2147/IJWH.S130879>
- Seelbach-Goebel, B. (2013). Antibiotic therapy for premature rupture of membranes and preterm labor and effect on fetal outcome. *Geburtshilfe Und Frauenheilkunde*, 73(12), 1218–1227. <https://doi.org/10.1055/s-0033-1360195>
- Seikku, L., Gissler, M., Andersson, S., Rahkonen, P., Stefanovic, V., Tikkanen, M., Paavonen, J., & Rahkonen, L. (2016). Asphyxia, Neurologic Morbidity, and Perinatal Mortality in Early-Term and Postterm Birth. In *PEDIATRICS* (Vol. 137, Issue 6).
- Shahwar, D., Ahmed, I., Amerjee, A., Hoodbhoy, Z. (2018). Comparison of Neonatal Outcome Between Category-1 and Non-Category-1 Primary Emergency Caesarean Section: A Retrospective Record Review In A Tertiary

- Care Hospital. *Pak J Med Sci* (Vol 34, No.4).  
<https://doi.org/10.12669/pjms.344.14496>
- Sharma, D., Choudhary, M., Lamba, M., & Shastri, S. (2016). Correlation of Apgar Score with Asphyxial Hepatic Injury and Mortality in Newborns: A Prospective Observational Study from India. *Clinical Medicine Insights: Pediatrics*, 10, CMPed.S38503. <https://doi.org/10.4137/cmped.s38503>
- Sharma, D., Sharma, P., & Shastri, S. (2016). Postnatal Complications of Intrauterine Growth Restriction. *Journal of Neonatal Biology*, 05(04).  
<https://doi.org/10.4172/2167-0897.1000232>
- Singh, R., Deo, S., & Pradeep, Y. (2012). The decision-to-delivery interval in emergency Caesarean sections and its correlation with perinatal outcome: Evidence from 204 deliveries in a developing country. *Tropical Doctor*, 42(2), 67–69. <https://doi.org/10.1258/td.2012.110315>
- Smith, J., Murphy, M., & Kandasamy, Y. (2013). The IUGR infant: A case study and associated problems with IUGR infants. In *Journal of Neonatal Nursing* (Vol. 19, Issue 2, pp. 46–53). <https://doi.org/10.1016/j.jnn.2012.12.005>
- Stark, A. R., Adamkin, D. H., Batton, D. G., Bell, E. F., Bhutani, V. K., Denson, S. E., Engle, W. A., Martin, G. I., Blackmon, L. R., Barrington, K. J., Hankins, G. D. V., Raju, T. N. K., Tomashek, K. M., Wallman, C., Riley, L. E., Couto, J., Kilpatrick, S. J., Bell, A. L., Coulehan, J. M., ... Hawks, D. (2006). The Apgar score. In *Pediatrics* (Vol. 117, Issue 4, pp. 1444–1447).  
<https://doi.org/10.1542/peds.2006-0325>
- Techane, M. A., Alemu, T. G., Wubneh, A. A., Belay, G. M., Tamir, T. T., et al. (2022). The effect of gestational age, low birth weight and parity on birth asphyxia among neonates in sub-Saharan Africa: systematic review and meta-analysis: 2021. *Italian Journal of Pediatrics* (2022) 48:114
- Temesgen, M. M., Gebregzi, A. H., Kasahun, H. G., Ahmed, S. A., & Woldegerima, Y. B. (2020). Evaluation of decision to delivery time interval and its effect on fetomaternal outcomes and associated factors in category-1 emergency caesarean section deliveries: Prospective cohort study. *BMC Pregnancy and Childbirth*, 20(1). <https://doi.org/10.1186/s12884-020-2828-z>
- The AAP Committee on Fetus and Newborn, ACOG Committee on Obstetric Practice. *Guidelines for Perinatal Care*. 7th ed. American Academy of Pediatricians; 2012:192.
- The American College of Obstetricians and Gynecologists, (2015). The Apgar Score. Committee Opinion No. 644. *American College of Obstetricians and Gynecologists. Obstet Gynecol* 2015;126:e52–5.
- Thomas, J., Paranjothy, S., & James, D. (2004). National cross sectional survey to determine whether the decision to delivery interval is critical in emergency caesarean section. In *British Medical Journal* (Vol. 328, Issue 7441, pp. 665–668). <https://doi.org/10.1136/bmj.38031.775845.7c>
- Tri Anjaswarni, Naomi Sih Bingah. (2015). The Relationship Between Gestational Age and Asphyxia Among Newborn Baby. Preceding Book the 11 International Nursing Conference.

- Tri Harjati, Eny Sendra, Lumastari Ajeng Wijayanti, & Ririn Indriani. (2022). Correlation between the Duration of Premature Rupture of Membranes during the Labor and the Asphyxia of Newborn Babies at Aura Syifa Hospital Kediri Regency. *STRADA Jurnal Ilmiah Kesehatan*, 11(2), 133–140. <https://doi.org/10.30994/sjik.v11i2.965>
- Wong, T. C. T., Lau, C. Q. H., Tan, E. L., Kanagalingam, D. (2017). Decision-to-delivery intervals and total duration of surgery for Caesarean sections in a tertiary general hospital. *Singapore Med J* 2017; 58(6): 332-337
- World Health Organization. (2006). Neonatal and Perinatal Mortality. Tersedia di [https://www.who.int/maternal\\_child\\_adolescent/topics/maternal/maternal\\_perinat al/en/](https://www.who.int/maternal_child_adolescent/topics/maternal/maternal_perinat al/en/), diakses pada 21 Maret 2021.
- Yakasai, I. A., Ahmed, Z. D., Okonofua, F. E. (2012). Decision delivery interval in emergency cesarean section in tertiary centre in Northern Nigeria. *Orient J Sci Res*;1:16–23
- Yun Sung, T., Seok Jae, Y., Ju You, H., Kyu Cho, C. (2021). Comparison of the effect of general and spinal anesthesia for elective cesarean section on maternal and fetal outcomes: a retrospective cohort study. *Anesth Pain Med (Seoul)*. 16(1): 49–55