



## DAFTAR PUSTAKA

- American Society of Anesthesiologists Task Force on Acute Pain Management 2012. Practice Guidelines for Acute Pain Management in the Perioperative Setting: An Updated Report by the American Society of Anesthesiologists Task Force on Acute Pain Management. *Anesthesiology*. 116(2). hal. 248–273.
- Arzola, C. dan Wieczorek, P. M. 2011. Efficacy of low-dose bupivacaine in spinal anaesthesia for Caesarean delivery: Systematic review and meta-analysis. *British Journal of Anaesthesia*. 107(3). hal. 308–318. doi: 10.1093/bja/aer200.
- Basbaum, A. I., Bautista, D. M., Scherrer, G. dan Julius, D. 2009. Cellular and Molecular Mechanisms of Pain. *Cell*. 139(2). hal. 267–284. doi: 10.1016/j.cell.2009.09.028.
- Benevides, M. L., Nuchi, R. J., Solcia, C. E., Júnior, V. M. X. dan Lima, A. F. B. 2018. Three Different Doses of Intrathecal Morphine as Part of a Multimodal Regimen for Pasca-Cesarean Delivery Analgesia: A Randomized Double-Blinded Trial. *Open Journal of Obstetrics and Gynecology*. 08(09). hal. 780–789. doi: 10.4236/ojog.2018.89081.
- Boekel, R. L. M. ., Warlé, M. C., Nielen, R. G. C., Vissers, K. C. P., Van Der Sande, R., Bronkhorst, E. M., Lerou, J. G. C. dan Steegers, M. A. H. 2019. Relationship between Pascaoperative Pain and Overall 30-Day Complications in a Broad Surgical Population: An Observational Study. *Annals of Surgery*. 269(5). hal. 856–865. doi: 10.1097/SLA.0000000000002583.
- Bogra, J., Arora, N. dan Srivastava, P. 2005. Synergistic effect of intrathecal fentanyl and bupivacaine in spinal anesthesia for cesarean section. *BMC Anesthesiology*. 5(5). hal. 1–6. doi: 10.1186/1471-2253-5-5.
- Brennan, T. J. 2011. Pathophysiology of pascaoperative pain. *Pain*. 152(3). hal. 33–40. doi: 10.1016/b978-0-12-809324-5.24249-1.
- Butterworth, J. F., Mackey, D. C. dan Wasnick, J. D. 2018. Anesthesia for Neurosurgery.in Butterworth, J. F., Mackey, D. C., dan Wasnick, J. D. (ed.)



*Morgan & Mikhail's Clinical Anesthesiology.* 6th ed. New York: McGraw Hill Education. hal. 961–989.

Castillo Sardi, A. S. Del, Olivadía, A. dan Martín, E. S. 2014. Comparison of a Standard Dose with a Low Dose of Levobupivacaine in Spinal Anesthesia for Caesarean Section. *Open Journal of Anesthesiology.* 04(12). hal. 318–323. doi: 10.4236/ojanes.2014.412045.

Chan, J. J. I., Tan, C. W., Yeam, C. T., Sultana, R., Sia, A. T. H., Habib, A. S. dan Sng, B. L. 2020. Risk factors associated with development of acute and subacute pasca-cesarean pain: A prospective cohort study. *Journal of Pain Research.* 13. hal. 2317–2328. doi: 10.2147/JPR.S257442.

Chandra, S., Nugroho, A. M., Agus, H. dan Susilo, A. P. 2019. How low can we go? A double-blinded randomized controlled trial to compare bupivacaine 5 mg and bupivacaine 7.5 mg for spinal anesthesia in cesarean delivery in Indonesian population. *Anesthesiology and Pain Medicine.* 9(2). hal. 1–6. doi: 10.5812/aapm.91275.

Chen, J. S., Kandle, P. F., Murray, I., Fitzgerald, L. A. dan Sehdev, J. S. 2021. *Physiology, Pain.* Treasure Islands: StatPearls Publishing.

Chiu, L. Y. L., Sun, T., Ree, R., Dunsmuir, D., Dotto, A., Ansermino, J. M. dan Yarnold, C. 2019. The evaluation of smartphone versions of the visual analogue scale and numeric rating scale as pascaoperative pain assessment tools: a prospective randomized trial. *Canadian Journal of Anesthesia.* 66(6). hal. 706–715. doi: 10.1007/s12630-019-01324-9.

Chooi, C. S. L., White, A. M., Tan, S. G. M., Dowling, K. dan Cyna, A. M. 2013. Pain vs comfort scores after Caesarean section: A randomized trial. *British Journal of Anaesthesia.* 110(5). hal. 780–787. doi: 10.1093/bja/aes517.

Chowdhury, N. I., Turner, J. H., Dorminy, C., Wu, J. dan Chandra, R. K. 2019. Preoperative quality-of-life measures predict acute pascaoperative pain in endoscopic sinus surgery. *Laryngoscope.* 129(6). hal. 1274–1279. doi: 10.1002/lary.27763.

Cummings, A., Orgil, B. D. dan Fitzgerald, B. M. 2021. *Intrathecal Morphine.* Treasure Islands: StatPearls Publishing.



- Cummings, K. dan Naguib, M. A. 2015. Opioid agonists and antagonists. In Flood, P., Rathmell, J. P., dan Shafer, S. (ed.) *Stoelting's Pharmacology and Physiology in Anesthetic Practice*. 5 ed. Philadelphia. hal. 217–256. doi: 10.1002/0470845015.coa002m.
- Ebrie, A. M., Woldeyohanis, M., Abafita, B. J., Ali, S. A., Zemedkun, A., Yimer, Y., Ashebir, Z. dan Mohammed, S. 2022. Hemodynamic and analgesic effect of intrathecal fentanyl with bupivacaine in patients undergoing elective cesarean section; a prospective cohort study. *PLoS ONE*. 17(7 July). hal. 1–12. doi: 10.1371/journal.pone.0268318.
- Erol, D. D. dan Aytac, I. 2018. Current anesthesia for Cesarean Section. *Clinical Journal of Obstetrics and Gynecology*. (1). hal. 61–66. doi: 10.11261/iryo1946.19.143.
- Ferrarezi, W. P. P., Braga, A. de F. de A., Ferreira, V. B., Mendes, S. Q., Brandão, M. J. N., Braga, F. S. da S. dan Carvalho, V. H. 2021. Spinal anesthesia for elective cesarean section. Bupivacaine associated with different doses of fentanyl: randomized clinical trial. *Brazilian Journal of Anesthesiology (English Edition)*. 71(6). hal. 642–648. doi: 10.1016/j.bjane.2021.03.030.
- Freys, S. M. dan Pogatzki-Zahn, E. 2019. Pain therapy to reduce perioperative complications. *Innovative Surgical Sciences*. 4(4). hal. 158–166. doi: 10.1515/iss-2019-0008.
- Gan, T. J. 2017. Poorly controlled postoperative pain: Prevalence, consequences, and prevention. *Journal of Pain Research*. 10. hal. 2287–2298. doi: 10.2147/JPR.S144066.
- Gan, T. J., Epstein, R. S., Leone-Perkins, M. L., Salimi, T., Iqbal, S. U. dan Whang, P. G. 2018. Practice Patterns and Treatment Challenges in Acute Postoperative Pain Management: A Survey of Practicing Physicians. *Pain and Therapy*. 7(2). hal. 205–216. doi: 10.1007/s40122-018-0106-9.
- Gauchan, S., Thapa, C., Prasai, A., Pyakurel, K., Joshi, I. dan J., T. 2014. Effects of intrathecal fentanyl as an adjunct to hyperbaric bupivacaine in spinal anesthesia for elective caesarean section. *Nepal Medical College Journal*. 16(1). hal. 5–8.



- Goel, S., Bhardwaj, N. dan Grover, V. K. 2003. Intrathecal fentanyl added to intrathecal bupivacaine for day case surgery: A randomized study. *European Journal of Anaesthesiology*. 20(4). hal. 294–297. doi: 10.1017/S0265021503000462.
- Gupta, N., Gupta, S., Agarwal, A., Agarwal, S. dan Mahto, R. 2016. A comparison of visual analog scale and Wong Baker facial pain scale for pain measurement in pasca lower caesarean section case. *International Journal of Reproduction, Contraception, Obstetrics and Gynecology*. 5(9). hal. 3032–3035. doi: 10.18203/2320-1770.ijrcog20162979.
- Hawker, G. A., Mian, S., Kendzerska, T. dan French, M. 2011. Measures of adult pain: Visual Analog Scale for Pain (VAS Pain), Numeric Rating Scale for Pain (NRS Pain), McGill Pain Questionnaire (MPQ), Short-Form McGill Pain Questionnaire (SF-MPQ), Chronic Pain Grade Scale (CPGS), Short Form-36 Bodily Pain Scale (SF. *Arthritis Care and Research*. 63(SUPPL. 11). hal. 240–252. doi: 10.1002/acr.20543.
- Hu, B., Zhang, X., Xu, G., Zhang, Q., Qian, P., Liu, S., Zhu, J. dan Shen, R. 2019. Association between COMT Polymorphism Val158Met and Opioid Consumption in Patients with Pascaoperative Pain: A Meta-Analysis. *NeuroSignals*. 26(1). hal. 11–21. doi: 10.1159/000487038.
- Hu, J., Zhang, C., Yan, J., Wang, R., Wang, Y. dan Xu, M. 2016. Sufentanil and bupivacaine combination versus bupivacaine alone for spinal anesthesia during cesarean delivery: A meta-analysis of randomized trials. *PLoS ONE*. 11(3). hal. 1–15. doi: 10.1371/journal.pone.0152605.
- Huang, B., Huang, Q., Hai, C., Zheng, Z., Li, Y. dan Zhang, Z. 2019. Height-based dosing algorithm of bupivacaine in spinal anaesthesia for decreasing maternal hypotension in caesarean section without prophylactic fluid preloading and vasopressors: Study protocol for a randomised controlled non-inferiority trial. *BMJ Open*. 9(5). hal. 1–8. doi: 10.1136/bmjopen-2018-024912.
- Imani, F., Rahimzadeh, P., Faiz, H. R., Nowruzina, S., Shakeri, A. dan Ghahremani, M. 2018. Comparison of the pasca-caesarean analgesic effect of adding dexmedetomidine to paracetamol and ketorolac: A randomized clinical trial.



*Anesthesiology and Pain Medicine.* 8(5). doi: 10.5812/aapm.85311.

- Jasim, H. H., Sulaiman, S. A. B. S., Khan, A. H. dan Rajah, U. A. S. 2017. Factors affecting pasca caesarean pain intensity among women in the northern peninsular of Malaysia. *Journal of Clinical and Diagnostic Research.* 11(9). hal. 7–11. doi: 10.7860/JCDR/2017/25364.10630.
- Kang, S. dan Brennan, T. 2016. Mechanisms of pascaoperative pain. *Anesthesia and Pain Medicine.* 120(22). hal. 236–248. doi: 10.1097/00029330-200711020-00002.
- Karaman, S., Günüsén, I., Uyar, M., Biricik, E. dan Firat, V. 2011. The effects of morphine and fentanyl alone or in combination added to intrathecal bupivacaine in spinal anesthesia for cesarean section. *Agri.* 23(2). hal. 57–63. doi: 10.5505/agri.2011.44127.
- Karcioglu, O., Topacoglu, H., Dikme, Ozgur dan Dikme, Ozlem 2018. A systematic review of the pain scales in adults: Which to use? *American Journal of Emergency Medicine.* 36(4). hal. 707–714. doi: 10.1016/j.ajem.2018.01.008.
- Kashani, I., Gilani, M. T., Keshtan, F. G. dan Sedaghat, A. 2018. Low-dose bupivacaine versus high-dose bupivacaine along with fentanyl for spinal anesthesia in opioid-dependent mothers. *International Journal of High Risk Behaviors and Addiction.* 7(3). hal. 7–11. doi: 10.5812/ijhrba.67694.
- Kehlet, H. 2018. Pascaoperative pain, analgesia, and recovery—bedfellows that cannot be ignored. *Pain.* 159(1). hal. S11–S16. doi: 10.1097/j.pain.0000000000001243.
- Kia, E. A., Tafish, R. dan Hs, Z. 2018. Morphine versus Fentanyl Used Spinally for Pasca Cesarean Section Analgesia : A Randomized Clinical Trial. *Journal of Anesthesia and Therapeutics.* 1(1). hal. 1–6.
- Kintu, A., Abdulla, S., Lubikire, A., Nabukenya, M. T., Igaga, E., Bulamba, F., Semakula, D. dan Olufolabi, A. J. 2019. Pascaoperative pain after cesarean section: Assessment and management in a tertiary hospital in a low-income country. *BMC Health Services Research.* 19(1). hal. 1–6. doi: 10.1186/s12913-019-3911-x.
- Korczeniewska, O. A., Kuo, F., Huang, C. Y., Nasri-Heir, C., Khan, J., Benoliel,



- R., Hirschberg, C., Eliav, E. dan Diehl, S. R. 2021. Genetic variation in catechol-O-methyltransferase is associated with individual differences in conditioned pain modulation in healthy subjects. *Journal of Gene Medicine.* 23(11). hal. 1–7. doi: 10.1002/jgm.3374.
- Laxmikanth, J., Segaran, S., George, S., John, A., Johns, J. dan Mani, N. 2021. To determine the efficacy of intrathecal dexmedetomidine and fentanyl as adjuvants to spinal anesthesia for lower abdominal surgeries below the level of umbilicus - A prospective randomized controlled double blinded study. *Bali Journal of Anesthesiology.* 5(1). hal. 15–20. doi: 10.4103/BJOA.BJOA\_160\_20.
- Lee, H. J., Cho, Y., Joo, H., Jeon, J. Y., Jang, Y. E. dan Kim, J. T. 2021. Comparative study of verbal rating scale and numerical rating scale to assess pascaoperative pain intensity in the pasca anesthesia care unit: A prospective observational cohort study. *Medicine.* 100(6). hal. e24314. doi: 10.1097/MD.00000000000024314.
- Mardani, I., Kresnoadi, E. dan Setyorini, R. H. 2017. Comparison Duration of Analgesia between Hyperbaric Bupivacaine + Morphine Intrathecal with Hyperbaric Bupivacaine + NaCl in Patients Undergoing Surgery with Spinal Anesthesia. *Jurnal Kedokteran Universitas Mataram.* 6(2.1). hal. 1–4.
- Marfuah, D., Nurhayati, N., Mutiar, A., Sumiati, M. dan Mardiani, R. 2019. Pain Intensity among Women with Pasca-Caesarean Section: A Descriptive Study. *KnE Life Sciences.* 2019. hal. 657–663. doi: 10.18502/kls.v4i13.5322.
- Meissner, W., Coluzzi, F., Fletcher, D., Huygen, F., Morlion, B., Neugebauer, E., Pérez, A. M. dan Pergolizzi, J. 2015. Improving the management of pasca-operative acute pain: Priorities for change. *Current Medical Research and Opinion.* 31(11). hal. 2131–2143. doi: 10.1185/03007995.2015.1092122.
- Moshiri, E., Malekianzadeh, B., Modir, H. dan Moghadam, V. D. 2017. Comparison of spinal anesthesia with low and conventional doses of bupivacaine on hemodynamic changes in severe preeclampsic patients candidate for cesarean section. *Biomedical and Pharmacology Journal.* 10(2). hal. 627–632. doi: 10.13005/bpj/1149.



- Myles, P. S., Myles, D. B., Galagher, W., Boyd, D., Chew, C., MacDonald, N. dan Dennis, A. 2017. Measuring acute pascaoperative pain using the visual analog scale: The minimal clinically important difference and patient acceptable symptom state. *British Journal of Anaesthesia*. 118(3). hal. 424–429. doi: 10.1093/bja/aew466.
- Nofal, W. H., Abdelaal, W. A. dan Elfawal, S. M. 2016. Minimum effective volume of bupivacaine in spinal anesthesia for elective cesarean section. Does it differ with height? A non-randomized parallel study. *Egyptian Journal of Anaesthesia*. 33(1). hal. 67–72. doi: 10.1016/j.ejga.2016.10.008.
- Olawin, A. M. dan Das, J. M. 2021. *Spinal Anesthesia*. Treasure Islands: StatPearls Publishing.
- Palada, V., Kaunisto, M. A. dan Kalso, E. 2018. Genetics and genomics in pascaoperative pain and analgesia. *Current Opinion in Anaesthesiology*. 31(5). hal. 569–574. doi: 10.1097/ACO.0000000000000633.
- Pathak A, Sharma S dan Jensen M 2018. The utility and validity of pain intensity rating scales for use in developing countries. *Pain Reports*. 3. hal. 1–8.
- Perry, M., Baumbauer, K., Young, E. E., Dorsey, S. G., Taylor, J. Y. dan Starkweather, A. R. 2019. The Influence of Race, Ethnicity and Genetic Variants on Pascaoperative Pain Intensity: An Integrative Literature Review. *Pain Management Nursing*. 20(3). hal. 198–206. doi: 10.1016/j.pmn.2018.11.002.
- Pogatzki-Zahn, E. M., Segelcke, D. dan Schug, S. A. 2017. Pascaoperative pain—from mechanisms to treatment. *Pain Reports*. 2(2). hal. 1–16. doi: 10.1097/PR9.0000000000000588.
- Rani, K., Zutshi, V., Patel, M. dan Marwah, S. 2016. Analgesic efficacy of intravenous paracetamol versus intravenous tramadol after caesarean section: a single blind randomized controlled study. *International Journal of Reproduction, Contraception, Obstetrics and Gynecology*. 5(12). hal. 4285–4289. doi: 10.18203/2320-1770.ijrcog20164329.
- Rindarto, R. dan Istanto, W. 2014. Pengaruh Penambahan Morfin dan Klonidin pada Bupivakain Dosis Rendah pada Anestesi Spinal untuk Bedah Besar



Ditinjau dari Perubahan Hemodinamik dan Kadar Glukosa Darah. *Medica Hospitalia: Journal of Clinical Medicine.* 2(1). hal. 6–13. doi: 10.36408/mhjcm.v2i1.81.

Rooftshoof, E., Joshi, G. P., Rawal, N., Van de Velde, M., Joshi, G. P., Pogatzki-Zahn, E., Van de Velde, M., Schug, S., Kehlet, H., Bonnet, F., Rawal, N., Delbos, A., Lavand'homme, P., Beloeil, H., Raeder, J., Sauter, A., Albrecht, E., ... Freys, S. 2021. PROSPECT guideline for elective caesarean section: updated systematic review and procedure-specific pascaoperative pain management recommendations. *Anaesthesia.* 76(5). hal. 665–680. doi: 10.1111/anae.15339.

Shafiei, F. T., McAllister, R. K. dan Lopez, J. 2021. *Bupivacaine Mechanism of Action Contraindications.* Treasure Islands: StatPearls Publishing.

Shembekar, M. C., Shembekar, C. A., Shembekar, S. C. dan Upadhye, J. J. 2019. Evaluation of pain relief and satisfaction in women using epidural analgesia in labour. *International Journal of Reproduction, Contraception, Obstetrics and Gynecology.* 8(3). hal. 1016–1021. doi: 10.18203/2320-1770.ijrcog20190873.

Small, C. dan Laycock, H. 2020. Acute pascaoperative pain management. *British Journal of Surgery.* 107(2). hal. 70–80. doi: 10.1002/bjs.11477.

Stamenkovic, D. M., Geric, V., Djordjevic, M., Raskovic, J., Slavkovic, Z., Randjelovic, T. dan Karanikolas, M. 2009. Subarachnoid morphine, bupivacaine and fentanyl as part of combined spinal-epidural analgesia for low anterior resection. A prospective, randomised, double-blind clinical trial. *Anaesthesia and Intensive Care.* 37(4). hal. 552–560. doi: 10.1177/0310057x0903700410.

Suyuthi, M. E. Z. 2012. *Comparison Between Spinal Anesthesia Combination of 0,5% 10 mg Bupivacaine, 0.05 Mg of Morphine Sulphate and 40% 50 mg Magnesium Sulfate, and 0.5% 10 mg Bupivacaine and 0,1 mg Morphine Sulphate on Pasca-Surgery Caesarian Section.* Universitas Hasanuddinn.

Thornton, P., Hanumanthaiah, D., O'Leary, R. A. dan Iohom, G. 2015. Effects of fentanyl added to a mixture of intrathecal bupivacaine and morphine for spinal



anaesthesia in elective caesarean section. *Romanian journal of anaesthesia and intensive care*. 22(2). hal. 97–102.

Tsen, L. C. dan Bateman, B. T. 2019. Anesthesia for cesarean delivery.in Chestnut, D., Wong, C., Tsen, L., Kee, W. D. N., Beilin, Y., Mhyree, J., et al. (ed.) *Chestnut's Obstetric Anesthesia: Principles and Practice*. Philadelphia: Elsevier. hal. 568–613. doi: 10.1007/978-3-319-74766-8\_51.

Uppal, V., Retter, S., Casey, M., Sancheti, S., Matheson, K. dan McKeen, D. M. 2020. Efficacy of Intrathecal Fentanyl for Cesarean Delivery: A Systematic Review and Meta-analysis of Randomized Controlled Trials With Trial Sequential Analysis. *Anesthesia and analgesia*. 130(1). hal. 111–125. doi: 10.1213/ANE.0000000000003975.

Van De Velde, M. 2019. Low-dose spinal anesthesia for cesarean section to prevent spinal-induced hypotension. *Current Opinion in Anaesthesiology*. 32(3). hal. 268–270. doi: 10.1097/ACO.0000000000000712.

Weigl, W., Bieryło, A., Wielgus, M., Krzemień-Wiczyńska, Ś., Kołacz, M. dan Dąbrowski, M. J. 2017. Perioperative analgesia after intrathecal fentanyl and morphine or morphine alone for cesarean section. *Medicine*. 96(48). hal. 1–7. doi: 10.1097/MD.0000000000008892.

Wikström, L., Nilsson, M., Broström, A. dan Eriksson, K. 2019. Patients' self-reported nausea: Validation of the Numerical Rating Scale and of a daily summary of repeated Numerical Rating Scale scores. *Journal of Clinical Nursing*. 28(5–6). hal. 959–968. doi: 10.1111/jocn.14705.

Willingham, M., Rangrass, G., Curcuru, C., Ben Abdallah, A., Wildes, T. S., McKinnon, S., Kronzer, A., Sharma, A., Helsten, D., Hall, B., Avidan, M. S. dan Haroutounian, S. 2020. Association between pascaoperative complications and lingering pasca-surgical pain: an observational cohort study. *British Journal of Anaesthesia*. 124(2). hal. 214–221. doi: 10.1016/j.bja.2019.10.012.

World Health Organization 2015. *WHO Statement on Caesarean Section Rates*. Geneva. doi: 10.1016/S0140-6736(80)91104-6.

Yang, M. M. H., Hartley, R. L., Leung, A. A., Ronksley, P. E., Jetté, N., Casha, S.



dan Riva-Cambrin, J. 2019. Preoperative predictors of poor acute postoperative pain control: A systematic review and meta-analysis. *BMJ Open*. 9(4). hal. 1–11. doi: 10.1136/bmjopen-2018-025091.

Yesuf, K. A., Gebremedhn, E. G. dan Melkie, T. B. 2017. Analgesic Effect of Intrathecal Fentanyl as an Adjuvant to Spinal Anaesthesia in Comparison with Spinal Anaesthesia with Bupivacaine Only for Mothers Delivered by Emergency Cesarean Section. *Journal of Anesthesia & Critical Care: Open Access*. 7(5). hal. 1–9. doi: 10.15406/jaccoa.2017.07.00278.