



DAFTAR PUSTAKA

- Al-Hasani, R. and Bruchas, M.R. (2011) ‘Molecular mechanisms of opioid receptor-dependent signaling and behavior’, *The Journal of the American Society of Anesthesiologists*, 115(6), pp. 1363–1381.
- Arzola, C. and Wieczorek, P.M. (2011) ‘Efficacy of low-dose bupivacaine in spinal anaesthesia for Caesarean delivery: systematic review and meta-analysis’, *British Journal of Anaesthesia*, 107(3), pp. 308–318.
- Bernards, C.M., McDonald, S.B. and Salinas, F. V (2009) *A practical approach to regional anesthesia*. Lippincott Williams & Wilkins.
- Betrán, A.P. et al. (2016) ‘WHO statement on caesarean section rates’, *Bjog*, 123(5), p. 667.
- Bhatia, P. and Chhabra, S. (2018) ‘Physiological and anatomical changes of pregnancy: Implications for anaesthesia’, *Indian journal of anaesthesia*, 62(9), p. 651.
- Białowolska, K. et al. (2020) ‘Fixed Dose versus Height-Adjusted Conventional Dose of Intrathecal Hyperbaric Bupivacaine for Caesarean Delivery: A Prospective, Double-Blinded Randomised Trial’, *Journal of Clinical Medicine*, 9(11), p. 3600.
- Bishop, D.G. (2014) ‘Predicting spinal hypotension during Caesarean section’, *Southern African Journal of Anaesthesia and Analgesia*, 20(4), pp. 14–17.
- Bollag, L. et al. (2021) ‘Society for obstetric anesthesia and perinatology: consensus statement and recommendations for enhanced recovery after cesarean’, *Anesthesia & Analgesia*, 132(5), pp. 1362–1377.
- Bower, J.R. and Kinsella, S.M. (2020) ‘Preventing and treating hypotension during spinal anaesthesia for caesarean section’, *BJA education*, 20(11), p. 360.
- Butterworth, J.F., Mackey, D.C. and Wasnick, J.D. (2018) *Morgan and Mikhail's clinical anesthesiology*. McGraw-Hill Education.
- Caughey, A.B. et al. (2014) ‘Safe prevention of the primary cesarean delivery’, *American journal of obstetrics and gynecology*, 210(3), pp. 179–193.
- Cenkowski, M.J. et al. (2019) ‘Hemodynamic effects of low-dose bupivacaine spinal anesthesia for cesarean section: A randomized controlled trial’, *Saudi Journal of Anaesthesia*, 13(3), p. 208.
- Chandra, S. et al. (2019) ‘How low can we go? A double-blinded randomized controlled trial to compare bupivacaine 5 mg and bupivacaine 7.5 mg for spinal anesthesia in cesarean delivery in Indonesian population’, *Anesthesiology and Pain Medicine*, 9(2).



Chestnut, D.H. *et al.* (2014) *Chestnut's obstetric anesthesia: principles and practice e-book*. Elsevier Health Sciences.

Dahl, J.B. *et al.* (1999) 'Intraoperative and Postoperative Analgesic Efficacy and Adverse Effects of Intrathecal Opioids in Patients Undergoing Cesarean Section with Spinal Anesthesia', *Anesthesiology*, 91(6), pp. 1919–1919. Available at: <https://doi.org/10.1097/00000542-199912000-00045>.

Dahlan, M.S. (2011) *Statistik untuk kedokteran dan kesehatan*. Penerbit Salemba.

Ferré, F. *et al.* (2020) 'Control of spinal anesthesia-induced hypotension in adults', *Local and regional anesthesia*, pp. 39–46.

Gropper, M.A. *et al.* (2019) *Miller's anesthesia, 2-volume set E-book*. Elsevier Health Sciences.

Hadzic, A. (2017) *Hadzic's textbook of regional anesthesia and acute pain management*. McGraw-Hill Education New York.

Hasanin, A. *et al.* (2017) 'Post-spinal anesthesia hypotension during cesarean delivery, a review article', *Egyptian Journal of Anaesthesia*, 33(2), pp. 189–193.

Huang, Q. *et al.* (2022) 'A Height-Based Dosing Algorithm of Bupivacaine in Spinal Anesthesia for Decreasing Maternal Hypotension in Cesarean Section Without Prophylactic Fluid Preloading and Vasopressors: A Randomized-Controlled Non-Inferiority Trial', *Frontiers in Medicine*, 9, p. 858115.

Iddrisu, M. and Khan, Z.H. (2021) 'Anesthesia for cesarean delivery: general or regional anesthesia—a systematic review', *Ain-Shams Journal of Anesthesiology*, 13(1), pp. 1–7.

Indradata, F. *et al.* (2021) 'Perbandingan Efektivitas Anestesi Spinal dengan Bupivacain 12, 5 Mg dan Bupivacain 5 Mg yang ditambah Fentanyl 50 Mcg pada Seksio Sesarea', *Jurnal Anestesi Obstetri Indonesia*, 4(1), pp. 11–17.

Karaman, S. *et al.* (2011) 'The effects of morphine and fentanyl alone or in combination added to intrathecal bupivacaine in spinal anesthesia for cesarean section', *Agri*, 23(2), pp. 57–63.

Kaye, A.D., Urman, R.D. and Vadivelu, N. (2012) *Essentials of regional anesthesia*. Springer.

Kinsella, S.M. *et al.* (2018) 'International consensus statement on the management of hypotension with vasopressors during caesarean section under spinal anaesthesia', *Obstetric Anesthesia Digest*, 38(4), pp. 171–172.

Kitaguchi, M. *et al.* (2022) 'Associated factors with umbilical arterial pH after cesarean delivery under spinal anesthesia: a retrospective cohort study', *Brazilian Journal of Anesthesiology*, 72, pp. 466–471.



Langesæter, E., Rosseland, L.A. and Stubhaug, A. (2008) ‘Continuous Invasive Blood Pressure and Cardiac Output Monitoring during Cesarean Delivery: A Randomized, Double-blind Comparison of Low-dose versus High-dose Spinal Anesthesia with Intravenous Phenylephrine or Placebo Infusion’, *The Journal of the American Society of Anesthesiologists*, 109(5), pp. 856–863.

Lanowski, J.-S. and von Kaisenberg, C.S. (2018) ‘The Surgical Technique of Caesarean Section: What is Evidence Based’, *Caesarean Section*, 4, pp. 63–77.

Laporan Nasional Riskesdas (2018). Jakarta.

MacLennan, K., O’Brien, K. and Macnab, W.R. (2015) *Core topics in obstetric Anaesthesia*. Cambridge University Press.

Manomayangkul, K. et al. (2016) ‘Reference values for umbilical cord blood gases of newborns delivered by elective cesarean section’, *J Med Assoc Thai*, 99(5), pp. 611–617.

Martin, J.A. et al. (2021) ‘Births: final data for 2019’.

Massoth, C., Töpel, L. and Wenk, M. (2020) ‘Hypotension after spinal anesthesia for cesarean section: how to approach the iatrogenic sympathectomy’, *Current Opinion in Anesthesiology*, 33(3), pp. 291–298.

Mehta, N. (2020) ‘Adjuvant drugs to local anesthetics’, in *Topics in Local Anesthetics*. IntechOpen.

Michelsen, T.M., Tronstad, C. and Rosseland, L.A. (2021) ‘Blood pressure and cardiac output during caesarean delivery under spinal anaesthesia: a prospective cohort study’, *BMJ open*, 11(6), p. e046102.

Moshiri, E., Malekianzadeh, B. and Modir, H. (2017) ‘Comparison of spinal anesthesia with low and conventional doses of bupivacaine on hemodynamic changes in severe Preeclampsic patients candidate for cesarean section’, *Biomedical and Pharmacology Journal*, 10(2), pp. 627–632.

Mylonas, I. and Friese, K. (2015) ‘Indications for and risks of elective cesarean section’, *Deutsches Ärzteblatt International*, 112(29–30), p. 489.

Ni, T. et al. (2018) ‘Intra-abdominal pressure, vertebral column length, and spread of spinal anesthesia in parturients undergoing cesarean section: An observational study’, *PLOS ONE*, 13(4), p. e0195137. Available at: <https://doi.org/10.1371/journal.pone.0195137>.

Olang, P.R., Wamalwa, D.C. and Ogutu, O. (2012) ‘Maternal hypotension and neonatal acidemia during Caesarean delivery under spinal anaesthesia’.

Olapour, A. et al. (2020) ‘Comparing the effect of bupivacaine and ropivacaine in cesarean delivery with spinal anesthesia’, *Anesthesiology and pain medicine*, 10(1).



Osorio-Gutiérrez, C.I. *et al.* (2021) ‘Comparison among subarachnoid opioid mix for cesarean section-An observational study’, *Colombian Journal of Anesthesiology*, 49(1).

Pérez, M.L.M., Hernandez Garre, J.M. and Pérez, P.E. (2021) ‘Analysis of factors associated with variability and acidosis of the umbilical artery pH at birth’, *Frontiers in pediatrics*, 9, p. 650555.

Pinto, S.L. *et al.* (2017) ‘Pregnant Women Positioning in Spinal Anesthesia for Cesarean Section: Integrative Review’, *International Archives of Medicine*, 10. Available at: <https://doi.org/10.3823/2449>.

Powlovich, L.G., Nemergut, E.C. and Collins, S.R. (2018) ‘Barash’s Clinical Anesthesia’, *Anesthesia & Analgesia*, 127(3), p. e41.

Regmi, B.D.S. *et al.* (2019) ‘Low dose hyperbaric bupivacaine with fentanyl in spinal anesthesia in elective cesarean section’, *Journal of Chitwan Medical College*, 9(2), pp. 79–82.

Reynolds, F. and Seed, P.T. (2005) ‘Anaesthesia for Caesarean section and neonatal acid-base status: a meta-analysis’, *Anaesthesia*, 60(7), pp. 636–653.

Ryu, C. *et al.* (2019) ‘Vasopressors for the management of maternal hypotension during cesarean section under spinal anesthesia: A Systematic review and network meta-analysis protocol’, *Medicine*, 98(1).

Shitemaw, T. *et al.* (2020) ‘Incidence and associated factors for hypotension after spinal anesthesia during cesarean section at Gandhi Memorial Hospital Addis Ababa, Ethiopia’, *PloS one*, 15(8), p. e0236755.

Šklebar, I., Bujas, T. and Habek, D. (2019) ‘Spinal anaesthesia-induced hypotension in obstetrics: prevention and therapy’, *Acta Clinica Croatica*, 58(Suppl 1), p. 90.

Soxhuku-Isufi, A., Shpata, V. and Sula, H. (2016) ‘Maternal and neonatal effects of vasopressors used for treating hypotension after spinal anesthesia for caesarean section: a randomized controlled study’, *Open Access Macedonian Journal of Medical Sciences*, 4(1), p. 54.

Surana, R.N. and Jain, T. (2020) ‘Evaluation of adjunctive analgesia with intrathecal fentanyl along with hyperbaric bupivacaine in spinal anesthesia for elective cesarean section’, *International Journal of Research in Medical Sciences*, 8(8), p. 2924.

Suresh, M., Segal, S. and Preston LR (2013) *Shnider and Levinson’s Anesthesia for Obstetrics*. 5th edn. Philadelphia: Wolters Kluwer Health/Lippincott Williams & Wilkins Health. Available at: <https://books.google.co.id/books?id=QMrjnQEACAAJ>.



Thornton, P. *et al.* (2015) ‘Effects of fentanyl added to a mixture of intrathecal bupivacaine and morphine for spinal anaesthesia in elective caesarean section’, *Romanian journal of anaesthesia and intensive care*, 22(2), p. 97.

Van de Velde, M. (2019) ‘Low-dose spinal anesthesia for cesarean section to prevent spinal-induced hypotension’, *Current Opinion in Anesthesiology*, 32(3). Available at: https://journals.lww.com/co-anesthesiology/Fulltext/2019/06000/Low_dose_spinal_anesthesia_for_cesarean_section_to.4.aspx.

Verma, V.K. *et al.* (2020) ‘Low dose spinal anaesthesia for caesarean section in parturients having pregnancy induced hypertension: Effect on haemodynamic response and vasopressor requirement: A prospective randomized double blind case control study’, *Indian Journal of Clinical Anaesthesia*, 7(1), pp. 104–111.

Weigl, W. *et al.* (2017) ‘Perioperative analgesia after intrathecal fentanyl and morphine or morphine alone for cesarean section: A randomized controlled study’, *Medicine*, 96(48).

Yu, C. *et al.* (2021) ‘Prediction of spinal anesthesia-induced hypotension during elective cesarean section: a systematic review of prospective observational studies’, *International Journal of Obstetric Anesthesia*, 47, p. 103175.