UNIVERSITAS GADJAH MADA
Neutrophil to Lymphocyte Ratio (NLR) Sebagai Prediktor Luaran Pasien Abses Leher Dalam Di RSUP Dr Sardjito
Okky Irawan, dr. Dian Paramita Wulandari, M.Sc., Sp. T.H.T.B.K.L.SubSp LF.(K); Dr. dr. Siswanto Sastrowijoto, Sp. T
Universitas Gadjah Mada, 2023 | Diunduh dari http://etd.repository.ugm.ac.id/

DAFTAR PUSTAKA

Aguslia, SD. & Farokah 2016, 'Pola Kuman dan Sensitivitas Terhadap Antibiotik Pasien Abses leher dalam di RSUP Dr. Kariadi Semarang', *Medica Hospitalia*, vol.3, pp. 164–169.

Alias, K. 2013, 'Diagnosis and Management of Deeper Neck Infections - A Review', *IOSR Journal of Dental and Medical Sciences*, vol. 9, no.5, pp. 36–41. doi: 10.9790/0853-0953641.

Altamimi, AS., Alshammari, RR., Banan, T., Alshammari, ZA., Alduheim, MA. & Alnais, IA. 2020, 'Deep neck infection treatment and prognosis : a systematic review', *Int J Med Dev Ctries*, vol 4, no.3, pp. 932–937. doi: 10.24911/ijmdc.511584390348

Arliando, MA., Adelien, A. & Utama, DS. 2017. 'Prevalensi Abses Leher Dalam di RSUP dr. Mohammad Hoesin Palembang Periode 1 Januari 2012 – 31 Desember 2015'. *Majalah Kedokteran Sriwijaya*, vol. 49, no.3, pp. 124–133.

Baglam, T., Binnetoglu, A., Cemal, A., Gerin, F., Demir, B. & Sari, M. 2015. 'Predictive value of the neutrophil-to-lymphocyte ratio in patients with deep neck space infection secondary to acute bacterial tonsillitis', *Int J Pediatr Otorhinolaryngol*, pp. 4–7. http://dx.doi.org/10.1016/j.ijporl.2015.06.016.

Barber, BR., Dziegielewski, PT., Biron, VL., Ma, A. & Seikaly, H. 2014. 'Factors associated with severe deep neck space infections : targeting multiple fronts'. *J Otolaryngol*, vol. 43, no. 35, pp. 1–7.

Bigcas, JLM. & Jain, KS. 2017. 'Management of parapharyngeal and retropharyngeal space infections', *Oper Tech Otolaryngol*, vol. 28, no.4, pp. 229–237. doi: 10.1016/j.otot.2017.08.007.

Cheepcharoenrat, C. 2019. 'The Result of Treatment of Deep Neck Infection in Patients Referred According to Public Health System'. *SAGE*, vol. 99, no. 10, pp. 627–632.

Colbert, K.R. & Devakumari, S. 2013. 'Diagnosis and Management of Deeper Neck Infections - A Review'. *IOSR - JDMS*, vol. 9, no.5, pp. 36–41.

CÖMERT, E. & ŞİMŞEK, B. 2022. 'Prognostic Factors in Deep Neck Infection'. *KU Tip Fak Derg*, vol. 24, no. 1, pp. 171–182.

Famous, KR., Delucchi, K., Ware, LB., Kangelaris, KN., Liu, KD., Thompson, B. T., Calfee, CS. 2016. 'ARDS Subphenotypes Respond Differently to Randomized Fluid Management Strategy Running title : Response to Fluid Management Strategy by ARDS Subphenotype Authors and Affiliations', *Am J Resp Crit Care Medicine*, pp. 1–49.

UNIVERSITAS GADJAH MADA
Neutrophil to Lymphocyte Ratio (NLR) Sebagai Prediktor Luaran Pasien Abses Leher Dalam Di RSUP Dr Sardjito
Okky Irawan, dr. Dian Paramita Wulandari, M.Sc., Sp. T.H.T.B.K.L.SubSp LF.(K); Dr. dr. Siswanto Sastrowijoto, Sp. T
Universitas Gadjah Mada, 2023 | Diunduh dari http://etd.repository.ugm.ac.id/

Fiorella, ML., Greco, P., Madami, LM., Giannico, OV., Pontillo, V. & Quaranta, N. 2020. 'New laboratory predictive tools in deep neck space infections'. *Acta otorhinolaryngol*, vol. 40, pp. 332–337.

Gleeson, M., Browning, GG. Burton, MJ., Clarke, R., John, H., Jones, NS., Lund, VJ., Luxon, LM., Watkinson, JC. 2008. *Scott-Brown's Otorhinolaryngology, Head and Neck Surgery*. London: Edward Arnold.

Grisaru-Soen, G., Komisar, O., Aizenstaein, O., Soudack, M., Schwartz, D. 2010. Retropharyngeal and parapharyngeal abscess in children-Epidemiology, clinical features and treatment', *Int J Pediatr Otorhinolaryngol*, vol. 74, no. 9, pp. 1016–1020. doi: 10.1016/j.ijporl.2010.05.030.

Gyawali, B., Ramakrishna, K. & Dhamoon, AS. 2019. 'Sepsis: The evolution in definition, pathophysiology, and management'. *SAGE Open Medicine*, vol. 7, doi: 205031211983504.

Huang, TT., Liu, TC., Chen, PR., Tseng, FU, Yeh, TH., Chen, YS. 2004. 'Deep neck infection: analysis of 185 cases', *Head Neck*, 26(10), pp. 854–860. doi: 10.1002/hed.20014.

Hwang, SY., Shin, TG., Jo, IJ., Jeon, K., Suh, GY., Lee, TR., Yoon, H., Cha, WC. & Sim, MS. 2017. 'Neutrophil-to-lymphocyte ratio as a prognostic marker in critically-ill septic patients'. *Am J Emerg Med*, vol. 35, no. 2, pp. 234–239. http://dx.doi.org/10.1016/j.ajem.2016.10.055.

Hynes, WL. & Walton, SL. 2000. 'Hyaluronidases of Gram-positive bacteria', *FEMS Microbiology Letters*, vol. 183, no. 2, pp. 201–207. doi: 10.1016/S0378-1097(99)00669-2.

Indrayani, LW. & Putra, IDA. 2019. 'Pola kuman dan sensitifitas antibiotika pada pasien abses leher dalam di RSUP Sanglah Denpasar periode 1 Januari - 31 Desember 2014'. *MEDICINA*, vol. 50, no. 1, pp. 143–147.

Jayagandhi, S., Cheruvu, SC., Manimaran, V., Mohanty, S. 2019. 'Deep Neck Space Infection: Study of 52 Cases', Indian *Journal of Otolaryngology and Head and Neck Surgery*, vol. 71, pp. 923–926. doi: 10.1007/s12070-019-01592-3.

Johnson, JT. & Rosen, CA. 2014 'Deep Neck infections', in *Bailey's Head & Neck Surgery Otolaryngology.* 5th edn. Philadelphia: Lippincott Williams & Wilkins, pp. 794–805.

Kameshwar, P., Rawangban, W. & Tangjaturonrasme, N. 2019. 'Serum Procalcitonin as Outcome Predictors in Deep Neck Infections'. *Biomed J Sci & Tech Res*, vol. 22, no. 2, pp. 16555–16559.

Kobayashi, SD., Malachowa, N. & Deleo, FR. 2015 'Pathogenesis of Staphylococcus aureus abscesses', *American Journal of Pathology*, vol. 185, no. 6, pp. 1518–1527. doi: 10.1016/j.ajpath.2014.11.030.

Lora, PS., Martins, EC., Da Fe Silveira, L., Viegas, K., Beck, AD., Júnior, GF. & Cremonese, RV. 2019. 'Neutrophil-lymphocyte ratio in the early diagnosis of sepsis in an intensive care unit: A case-control study'. *Rev Bras Ter Intensive*, vol. 31, no. 1, pp. 63–70.

Mahapatra, S. & Heffner, AC. 2021. 'Septic shock'. *StatPearls*, pp. 1–8.

UNIVERSITAS GADJAH MADA
Neutrophil to Lymphocyte Ratio (NLR) Sebagai Prediktor Luaran Pasien Abses Leher Dalam Di RSUP Dr Sardjito
Okky Irawan, dr. Dian Paramita Wulandari, M.Sc., Sp. T.H.T.B.K.L.SubSp LF.(K); Dr. dr. Siswanto Sastrowijoto, Sp. T
Universitas Gadjah Mada, 2023 | Diunduh dari http://etd.repository.ugm.ac.id/

https://www.ncbi.nlm.nih.gov/books/NBK430939/?report=printable.

Maharaj, S., Ahmed, S. & Pillay, P. 2019. 'Deep Neck Space Infections: A Case Series and Review of the Literature'. *SAGE*, vol. 12, pp. 1–5.

Naess, A., Saervold, S., Mo, R. & Eide, G.E. 2017. 'Role of neutrophil to lymphocyte and monocyte to lymphocyte ratios in the diagnosis of bacterial infection in patients with fever'. *Infection*. Vol. 45, pp. 299-307

Norton, NS. 2012. *Netter's Head and Neck Anatomy for Dentistry*. 2nd edn. Philadelphia: Elsevier Saunders.

Nugroho, A., Suwarman & Nawawi, AM. 2013. 'Hubungan antara Rasio Neutrofil-Limfosit dan Skor Sequencial Organ Failure Assesment pada Pasien yang Dirawat di Ruang Intensive Care Unit'. *Jurnal anestesi perioperatif*, vol. 1, no. 3, pp. 189–96.

Nurdin, N., Kalma, K., Hasnawati, H. & Nasir, H. 2021. 'PROFIL NILAI Neutrophil Lymphocyte RATIO (NLR) PADA PENDERITA DIABETES MELITUS TIPE-2'. *Jurnal Media Analis Kesehatan*, vol. 12, no.1, pp. 64–70.

Pineda-Alvarado, A., Lugo-Machado, JA., Canché-Martin, E., Quintero, JZ., Arellano-Ridriguez, I. & Lizárraga, LL. 2021. 'Prevalence, morbidity and mortality of deep neck abscess in a tertiary hospital from Northwestern Mexico'. *Rom J Rhinol*, vol. 11, no. 41, pp. 41–46.

Priyamvada, S. & Motwani, G. 2019. 'A Study on Deep Neck Space Infections', *Indian Journal of Otolaryngology and Head and Neck Surger*y, vol. 71, no.1, pp. 912–917. doi: 10.1007/s12070-019-01583-4.

Rijal, S. & Romdhoni, AC. 2018. 'Bacteria Pattern, Results of Antibiotic Sensitivity Test, and Complications of Deep Neck Abscess Patients in Dr. Soetomo General Hospital', *Biomolecular and Health Science Journal*, vol. 1, no. 2, pp. 124. doi: 10.20473/bhsj.v1i2.9832.

Ruop, M., Perkins, N., Whitcomb, B. & Schisterman, E. 2008. Youden Index and Optimal Cut-Point Estimated from Observation Affected by a Lower Limit Detection. *Biom J*, vol. 50, no. 3, pp. 419–430.

Russell, CD., Parajuli, A., Gale, HJ., Bulteel, NS., Schuetz, P., de Jager, CPC., Loonen, AJM., Merekoulias, GI. & Baillie, JK. 2019. 'The utility of peripheral blood leucocyte ratios as biomarkers in infectious diseases: A systematic review and meta-analysis'. *Journal of Infection*, vol. 78, no. 5, pp. 339–348. doi: 10.1016/j.jinf.2019.02.006.

Salciccioli, JD., Marshall, DC., Pimentel, MAF., Santos, MD., Pollard, T., Celi, AA. & Shalhoub, J. 2015. 'The association between the neutrophil-to-lymphocyte ratio and mortality in critical illness: An observational cohort study'. *Crit Care*, vol. 19, no.1, pp. 1–8.

Sataloff, RT. 2016, *Sataloff'S Comprehensive Textbook of Otolaryngology Head and Neck Surgery*. 1st edn. Daryaganj: Jaypee Brothers Medical Publisher.

Sebastian, GP. & Ramalingam, K. 2021. 'Deep Neck Space Infections: Otolaryngologist Perspectives in Triage and Challenges'. *Am J Otolaryngol*, vol. 4, no. 2, pp. 1125.

UNIVERSITAS GADJAH MADA
Neutrophil to Lymphocyte Ratio (NLR) Sebagai Prediktor Luaran Pasien Abses Leher Dalam Di RSUP Dr Sardjito
Okky Irawan, dr. Dian Paramita Wulandari, M.Sc., Sp. T.H.T.B.K.L.SubSp LF.(K); Dr. dr. Siswanto Sastrowijoto, Sp. T
Universitas Gadjah Mada, 2023 | Diunduh dari http://etd.repository.ugm.ac.id/

Seer Yee, ML., Rahim, NA., Ngah, NA., Aziz, YFA., Subha, S. 2014. 'Predicting Neck Abscess with Contrast-Enhanced Computed Tomography', *Advances in Otolaryngology*, pp. 1–8. doi: 10.1155/2014/896831.

Şentürk, M., Azgın, İ., Övet, G., Alataş, N., Ağırgöl, B. & Yılmaz, E. 2016. 'The role of the mean platelet volume and neutrophil-to-lymphocyte ratio in peritonsillar abscesses'. *Braz J Otorhinolaryngol*, vol. 82, no.6, pp. 662–667.

Singer, M., Deutschman, CS., Seymour, C., Shankar-Hari, M., Annane, D., Bauer, M., Bellomo, R., Bernard, G., Chiche, J., Coopersmith, C., Hotchkiss, R., Levy, M., Marshall, J., Martin, G., Opal, S., Rubenfeld, GTP., Vincent, J. & Angus, D. 2016. 'The Third International Consensus Definitions for Sepsis and Septic Shock (Sepsis-3)'. *JAMA*, vol. 315, no. 8, pp. 801–810.

Suehara, AB., Rodrigues, AAN., Kavabata, NK., Menezes, MB., Ramos, E. de A., Kawamukai, JN. & Gonçalves, AJ. 2020. 'Predictive factors of lethality and complications of deep fascial space infections of the neck'. *Rev Col Bras Cir*, vol. 47, pp. 1–8.

Treviño-gonzalez, JL., Maldonado-chapa, F., Alejandra, J., Germán, MA. & García-villanueva, JMZ. 2021. 'Deep Neck Infections : Demographic and Clinical Factors Associated with Poor Outcomes'. *ORL*, pp. 1–9.

Umihanic, S., Umihamic, S., Ramic, N., Kamenjakovic, S., Tihic, N. Mahmutovic, E. 2018. 'Factors associated with poor prognosis in deep neck infections', *Medeniyet medical journal*, vol. 33, no. 1, pp. 28–32. doi: 10.5222/MMJ.2018.49140.

Vaikjärv, R., Mändar, R. & Kasenõmm, P. 2019. 'Peritonsillar abscess is frequently accompanied by sepsis symptoms'. *Eur Archives of OtoRhinoLaryngol*, vol. 276, no.6, pp. 1721–1725. https://doi.org/10.1007/s00405-019-05424-6.

Vatdianto & Mavarenia.2015.Blog Post.*Matched Case Control Study*.Dilihat pada 5 April 2023

Velhonoja, J., Lääveri, M., Soukka, T., Irjala, H. & Kinnunen, I. 2020. 'Deep neck space infections: an upward trend and changing characteristics.' . *Eur Archives of OtoRhinoLaryngol*, vol. 277, no. 3, pp. 863–872. https://doi.org/10.1007/s00405-019-05742-9.

Viakhane, B.A., Kandhi, P.W. & Primadewi, N. 2021.'RASIO NEUTROFIL LIMFOSIT SEBAGAI PREDIKTOR DERAJAT KEPARAHAN ABSES LEHER DALAM'. Universitas Sebelas Maret.

Vieira, F., Allen, SM., Stocks, RMS. & Thompson, J.W. 2008. 'Deep Neck Infection.' , vol. 41, pp. 459–483. doi: 10.1016/j.otc.2008.01.002.

Wang, LF., Kuo, WR., Shai, SM., Huang, KJ. 2003. 'Characterizations of life-threatening deep cervical space infections: A review of one hundred ninety-six cases', *American Journal of Otolaryngology - Head and Neck Medicine and Surgery*, vol. 24, no.2, pp. 111–117. doi: 10.1053/ajot.2003.31.

UNIVERSITAS
GADJAH MADA
**Neutrophil to Lymphocyte Ratio (NLR) Sebagai Prediktor Luaran Pasien Abses Leher Dalam Di RSUP Dr
Sardjito**
Okky Irawan, dr. Dian Paramita Wulandari, M.Sc., Sp. T.H.T.B.K.L.SubSp LF.(K); Dr. dr. Siswanto Sastrowijoto, Sp. T
Universitas Gadjah Mada, 2023 | Diunduh dari http://etd.repository.ugm.ac.id/

WHO. 2001. Health Research Methodology.A Guide for Training in ResearchMethods. 2nd Edition. World Health Organization, Regional Office for the Western Pacific Manila.

Wiargitha, IK. 2017. 'Hubungan Rasio Neutrofil Limfosit dalam Memprediksi Mortalitas pada Pasien Peritonitis yang Dioperasi di RSUP Sanglah Denpasar'. Universitas Udayana.

World Union of Wound Healing Societies 2019, 'World Union of Wound Healing Societies (WUWHS) Consensus Document. Wound exudate: effective assessment and management', Wounds International. Available at: https://www.woundsinternational.com/download/resource/7732.

Youden, W. 1950. 'Index for rating diagnostic tests'. *Cancer*, vol.3, pp.32–35.

Zamrini, B., Hashemi, SB., Hashemi, SH., Rafiee, Z. & Ehsani, S. 2012. 'Prevalence of Odontogenic Deep Head and Neck Spaces Infection and its Correlation with Length of Hospital Stay'. *Dent J*, vo. 13, no.1, pp. 29–35