

DAFTAR PUSTAKA

- Al-Kawaz, S.A., 2014. Factor That Predict Mortality Rate in Biliary Atresia. *J. Fac. Med. Baghdad* 56: 136–140.
- Arnon, R., Annunziato, R.A., D'Amelio, G., Chu, J., & Shneider, B.L., 2016. Liver transplantation for Biliary Atresia: Is there a difference in outcome for infants? *J. Pediatr. Gastroenterol. Nutr.* 62: 220–225. doi:10.1097/MPG.0000000000000986
- Averbukh, L.D., & Wu, G.Y., 2018. Evidence for viral induction of biliary atresia: A review. *J. Clin. Transl. Hepatol.* 6: 410–419. doi:10.14218/JCTH.2018.00046
- Baek, S.H., Kang, J.M., Ihn, K., Han, S.J., Koh, H., & Ahn, J.G., 2020. The Epidemiology and Etiology of Cholangitis after Kasai Portoenterostomy in Patients with Biliary Atresia. *J. Pediatr. Gastroenterol. Nutr.* 70: 171–177. doi:10.1097/MPG.0000000000002555
- Centers for Disease Control and Prevention, 2020. Cytomegalovirus (CMV) and Congenital CMV Infection: Transmission. *CDC.gov* 1–2.
- Chan, K.W.E., Lee, K.H., Wong, H.Y.V., Tsui, S.Y.B., Mou, J.W.C., & Tam, Y.H., 2021. Impact of Age of Patient and Experience of Surgeon on the Outcome after Kasai Portoenterostomy: Can We Delay the Surgery? *Eur. J. Pediatr. Surg.* 31: 335–340. doi:10.1055/s-0040-1713934
- Chen, S.Y., Lin, C.C., Tsan, Y.T., Chan, W.C., Wang, J. Der, Chou, Y.J., et al., 2018. Number of cholangitis episodes as a prognostic marker to predict timing of liver transplantation in biliary atresia patients after Kasai portoenterostomy. *BMC Pediatr.* 18: 1–7. doi:10.1186/s12887-018-1074-2
- Chiopris, G., Veronese, P., Cusenza, F., Procaccianti, M., Perrone, S., Daccò, V., et al., 2020. Congenital cytomegalovirus infection: Update on diagnosis and treatment. *Microorganisms* 8: 1–17. doi:10.3390/microorganisms8101516
- Chusilp, S., Sookpotarom, P., Tepmalai, K., Rajatapiti, P., Chongsrisawat, V., Poovorawan, Y., et al., 2016. Prognostic values of serum bilirubin at 7th day post-Kasai for survival with native livers in patients with biliary atresia. *Pediatr. Surg. Int.* 32: 927–931. doi:10.1007/s00383-016-3951-9
- Davenport, M., Puricelli, V., Farrant, P., Hadzic, N., Mieli-Vergani, G., Portmann, B., et al., 2004. The Outcome of the Older (≥ 100 Days) Infant with Biliary Atresia. *J. Pediatr. Surg.* 39: 575–581. doi:10.1016/j.jpedsurg.2003.12.014
- De Lédinghen, V., Le Bail, B., Rebouissoux, L., Fournier, C., Foucher, J., Miette, V., et al., 2007. Liver stiffness measurement in children using fibroscan: Feasibility study and comparison with fibrotest, aspartate transaminase to platelets ratio index, and liver biopsy. *J. Pediatr. Gastroenterol. Nutr.* 45: 443–450. doi:10.1097/MPG.0b013e31812e56ff
- Derusso, P.A., Ye, W., Shepherd, R., Haber, B.A., Benjamin, L., Whittington, P.F., et al., 2014. Growth Failure and Outcome in Infant with Biliary Atresia: A Report from the Biliary Atresia Research Consortium 46: 1–17. doi:10.1002/hep.21923.Growth
- Du, M., Wang, J., Tang, Y., Jiang, J., Chen, G., Huang, Y., et al., 2020. Evaluation

- of Perioperative Complications in the Management of Biliary Atresia. *Front. Pediatr.* 8: 1–10. doi:10.3389/fped.2020.00460
- Elhassan, A.B.E., Mohamed, S., Ali, A., Elbathani, M., Abdallah, O., Mohamed, A., et al., 2021. 388 Detection of Cytomegalovirus Infection in Infants with Biliary Atresia: A Meta-analysis. *Br. J. Surg.* 108: 3–9. doi:10.1093/bjs/znab134.268
- Fischler, B., Czubkowski, P., Dezsofi, A., Liliemark, U., Socha, P., Sokol, R.J., et al., 2022. Incidence, Impact and Treatment of Ongoing CMV Infection in Patients with Biliary Atresia in Four European Centres. *J. Clin. Med.* 11: 4–9. doi:10.3390/jcm11040945
- Fischler, B., Svensson, J.F., & Nemeth, A., 2009. Early cytomegalovirus infection and the long-term outcome of biliary atresia. *Acta Paediatr. Int. J. Paediatr.* 98: 1600–1602. doi:10.1111/j.1651-2227.2009.01416.x
- Fischler, B., Woxenius, S., Nemeth, A., & Papadogiannakis, N., 2005. Immunoglobulin deposits in liver tissue from infants with biliary atresia and the correlation to cytomegalovirus infection. *J. Pediatr. Surg.* 40: 541–546. doi:10.1016/j.jpedsurg.2004.11.035
- Gad, E.H., Kamel, Y., Salem, T.A.H., Ali, M.A.H., & Sallam, A.N., 2021. Short- and long-term outcomes after Kasai operation for type III biliary atresia: Twenty years of experience in a single tertiary Egyptian center-A retrospective cohort study. *Ann. Med. Surg.* 62: 302–314. doi:10.1016/j.amsu.2021.01.052
- Gallo, A., & Esquivel, C.O., 2013. Current options for management of biliary atresia. *Pediatr. Transplant.* 17: 95–98. doi:10.1111/petr.12040
- Goda, T., Kawahara, H., Kubota, A., Hirano, K., Umeda, S., Tani, G., et al., 2013. The most reliable early predictors of outcome in patients with biliary atresia after Kasai's operation. *J. Pediatr. Surg.* 48: 2373–2377. doi:10.1016/j.jpedsurg.2013.08.009
- Grieve, A., Makin, E., & Davenport, M., 2013. Aspartate Aminotransferase-to-Platelet Ratio index (APRI) in infants with biliary atresia: Prognostic value at presentation. *J. Pediatr. Surg.* 48: 789–795. doi:10.1016/j.jpedsurg.2012.10.010
- Grizelj, R., Vuković, J., Novak, M., & Batinica, S., 2010. Biliary atresia: The Croatian experience 1992-2006. *Eur. J. Pediatr.* 169: 1529–1534. doi:10.1007/s00431-010-1266-8
- Gunkel, J., Wolfs, T.F.W., De Vries, L.S., & Nijman, J., 2014. Predictors of severity for postnatal cytomegalovirus infection in preterm infants and implications for treatment. *Expert Rev. Anti. Infect. Ther.* 12: 1345–1355. doi:10.1586/14787210.2014.966080
- Jonas, M.M., Fischler, B., Casswall, T., Malmberg, P., & Nemeth, A., 2002. Re: Ganciclovir treatment in infants with cytomegalovirus infection and cholestasis [2] (multiple letters). *J. Pediatr. Gastroenterol. Nutr.* 35: 399–400. doi:10.1097/00005176-200209000-00035
- Khayat, A., Alamri, A.M., & Saadah, O.I., 2021. Outcomes of late Kasai portoenterostomy in biliary atresia: a single-center experience. *J. Int. Med. Res.* 49. doi:10.1177/03000605211012596
- Kim, S.Y., Seok, J.Y., Han, S.J., & Koh, H., 2010. Assessment of Liver Fibrosis

- and Cirrhosis by Aspartate Aminotransferase-to-Platelet Ratio Index in Children With Biliary Atresia. *J. Pediatr. Gastroenterol. Nutr.* 51: 198–202. doi:10.1097/MPG.0b013e3181da1d98
- Kliegman, R.M., Geme III St, J.W., Blum, N.J., Shah, S.S., Tasker, R.C., Wilson, karen M., et al., 2020. Nelson Textbook of Pediatrics, in: Edition 21ft. pp. 6908–6926.
- Lazim, H.H., Arif, H.S., Kadhim, H.S., Al-Kahafaji, K.R., & Abdulghafour, K.H., 2015. The association between biliary atresia and cytomegalovirus hepatitis. *J. Nepal Paediatr. Soc.* 35: 269–273. doi:10.3126/jnps.v35i3.13547
- Lee, W.S., Chai, P.F., Lim, K.S., Lim, L.H., Looi, L.M., & Ramanujam, T.M., 2009. Outcome of biliary atresia in Malaysia: A single-centre study. *J. Paediatr. Child Health* 45: 279–285. doi:10.1111/j.1440-1754.2009.01490.x
- Leevan, E., Matsuoka, L., Cao, S., Groshen, S., & Alexopoulos, S., 2019. Biliary-Enteric Drainage vs Primary Liver Transplant as Initial Treatment for Children with Biliary Atresia. *JAMA Surg.* 154: 26–32. doi:10.1001/jamasurg.2018.3180
- Li, D., Chen, X., Fu, K., Yang, J., & Feng, J., 2017. Preoperative nutritional status and its impact on cholangitis after Kasai portoenterostomy in biliary atresia patients. *Pediatr. Surg. Int.* doi:10.1007/s00383-017-4118-z
- Liu, F., Yeung, F., & Chung, P.H.Y., 2022. The outcome of Kasai portoenterostomy after day 70 of life. *Front. Pediatr.* 10: 1–6. doi:10.3389/fped.2022.1015806
- Luo, Y., & Zheng, S., 2008. Current concept about postoperative cholangitis in biliary atresia. *World J. Pediatr.* 4: 14–19. doi:10.1007/s12519-008-0003-0
- Lykavieris, P., Chardot, C., Sokhn, M., Gauthier, F., Valayer, J., & Bernard, O., 2005. Outcome in adulthood of biliary atresia: A study of 63 patients who survived for over 20 years with their native liver. *Hepatology* 41: 366–371. doi:10.1002/hep.20547
- Mangunkusumo, C., Rahayatri, T.H., Siahaan, K.B.P., Ulima, R.P., Stephanie, M., Oswari, H., et al., 2019. Evaluasi Pasien Praoperasi Transplantasi Hati Anak Evaluation of Preoperative Pediatric Liver Transplantation 7: 53–60. doi:10.23886/ejki.7.10182.1
- Min, C., Song, J.Y., & Jeong, S.J., 2017. Characteristics and prognosis of hepatic cyto - megalovirus infection in children : 10 years of experience at a university hospital in Korea 60: 261–265.
- Muthukanagarajan, S.J., Karnan, I., Srinivasan, P., Sadagopan, P., & Manickam, S., 2016. Diagnostic and prognostic significance of various histopathological features in extrahepatic biliary atresia. *J. Clin. Diagnostic Res.* 10: EC23–EC27. doi:10.7860/JCDR/2016/19252.8035
- Nakase, H., & Herfarth, H., 2016. Cytomegalovirus Colitis, Cytomegalovirus Hepatitis and Systemic Cytomegalovirus Infection: Common Features and Differences. *Inflamm. Intest. Dis.* 1: 15–23. doi:10.1159/000443198
- Namasamayam, D., & Nallusamy, M., 2017. Factors influencing outcome after hepatic portoenterostomy among extrahepatic bile duct atresia patients in hospital Sultanah Bahiyah, Alor Setar. *Med. J. Malaysia* 72: 329–332.
- Nguyen, A.P., Hai, Y., Pham, T., Vu, G.H., & Ha, M., 2021. Biliary atresia liver histopathological determinants of early post-Kasai outcome 56: 1169–1173.

- Nightingale, S., Stormon, M.O., O'Loughlin, E. V., Shun, A., Thomas, G., Benchimol, E.I., et al., 2017. Early Posthepatoportoenterostomy Predictors of Native Liver Survival in Biliary Atresia. *J. Pediatr. Gastroenterol. Nutr.* 64: 203–209. doi:10.1097/MPG.0000000000001289
- Ohhama, Y., Shinkai, M., Fujita, S., Nishi, T., & Yamamoto, H., 2000. Early prediction of long-term survival and the timing of liver transplantation after the Kasai operation. *J. Pediatr. Surg.* 35: 1031–1034. doi:10.1053/jpsu.2000.7765
- Oliveira, N.L.G. De, Kanawaty, F.R., Costa, S.C.B., & Hessel, G., 2002. Infection By Cytomegalovirus in Patients With Neonatal Cholestasis. *Arq. Gastroenterol.* 39: 132–136.
- Pratiwi, F.P., Widowati, T., & Prawirohartono, E.P., 2011. Comparison of nutritional status among children with biliary atresia according to age at the time of Kasai procedure. *Paediatr. Indones.* 51: 207–212. doi:10.14238/pi
- Qi, Q., Wang, Y., Wu, Q., SU, P., Wang, D., Li, T., et al., 2022. Predicting the outcomes of Kasai portoenterostomy for biliary atresia: a cohort study. *Yale J. Biol. Med.* 1–17.
- Rauschenfels, S., Krassmann, M., Al-Masri, A.N., Verhagen, W., Leonhardt, J., Kuebler, J.F., et al., 2009. Incidence of hepatotropic viruses in biliary atresia. *Eur. J. Pediatr.* 168: 469–476. doi:10.1007/s00431-008-0774-2
- Reni, W., Endang, W., & Badriul, H., 2011. *Paediatrica Indonesiana* 51: 207–212. doi:10.14238/pi
- Rokhayati, E., Setyoboedi, B., & Arief, S., 2021. Aktivitas Serum Bilirubin pada Pasien Atresia Bilier Sebelum dan Sesudah Operasi Kasai 4: 66–72. doi:10.13057/smj.v4i2.
- Sanghai, S.R., Shah, I., Bhatnagar, S., & Murthy, A., 2009. Incidence and prognostic factors associated with biliary atresia in western India. *Ann. Hepatol.* 8: 120–122. doi:10.1016/s1665-2681(19)31789-2
- Schreiber, R.A., Harpavat, S., Hulscher, J.B.F., & Wildhaber, B.E., 2022. Biliary Atresia in 2021: Epidemiology, Screening and Public Policy. *J. Clin. Med.* 11. doi:10.3390/jcm11040999
- Serinet, M.O., Wildhaber, B.E., Broué, P., Lachaux, A., Sarles, J., Jacquemin, E., et al., 2009. Impact of age at Kasai operation on its results in late childhood and adolescence: A rational basis for biliary atresia screening. *Pediatrics* 123: 1280–1286. doi:10.1542/peds.2008-1949
- Shehata, S., Waheeb, S., Osman, M., Mahfouz, A., & Elrouby, A., 2018. Prognostic factors for the outcome of Kasai portoenterostomy for infants with biliary atresia in Egypt. *Alexandria J. Pediatr.* 31: 112. doi:10.4103/ajop.ajop_25_18
- Shen, C., Zheng, S., Wang, W., & Xiao, X.-M., 2008. Relationship between prognosis of biliary atresia and infection of cytomegalovirus. *World J. Pediatr.* 4: 123–126. doi:10.1007/s12519-008-0024-8
- Shneider, B.L., Magee, J.C., Karpen, S.J., Rand, E.B., Narkewicz, M.R., Bass, L.M., et al., 2016. Total Serum Bilirubin within Three Months of Heptopartoenterostomy Predicts Short-term Outcomes in Biliary Atresia. *Physiol. Behav.* 176: 139–148. doi:10.1016/j.jpeds.2015.11.058.Total

- Sira, M.M., Salem, T.A.-H., & Sira, A.M., 2013. Biliary Atresia: A Challenging Diagnosis. *Glob. J. Gastroenterol. Hepatol.* doi:10.12970/2308-6483.2013.01.01.6
- Situmorang, L., Setyoboedi, B., Mastutik, G., & Arief, S., 2020. Infection of Cytomegalovirus in Cholestasis Infant with Biliary Atresia. *Clin. Pathol. Med. Lab.* 26: 175–181.
- Smiechura, M., Struzycka, M., & Konopka, W., 2014. Congenital and acquired cytomegalovirus infection and hearing evaluation in children. *Otolaryngol. Pol.* 68: 303–307. doi:10.1016/j.otopol.2014.04.003
- Song, Z., Dong, R., Shen, Z., Chen, G., Yang, Y., & Zheng, S., 2017. Surgical outcome and etiologic heterogeneity of infants with biliary atresia who received Kasai operation less than 60 days after birth: A retrospective study. *Med. (United States)* 96: 1–5. doi:10.1097/MD.00000000000007267
- Spengler, U., 2019. Liver Disease Associated with Non-Hepatitis Viruses, 2nd ed, Encyclopedia of Gastroenterology, Second Edition. Elsevier Inc. doi:10.1016/B978-0-12-801238-3.65782-3
- Suchy, F.J., Sokol, R.J., & Balistreri, W.F., 2007. Liver Disease in Children, Third Edit. ed, Cambridge University.
- Vig, A., Elhence, P., Rathod, K.J., Nayak, S., Jadhav, A.S., Pathak, M., et al., 2023. Effect of cytomegalovirus Infection on Initial Presentation and Overall Prognosis of Biliary Atresia Patients. *J. Indian Assoc. Pediatr. Surg.* 28: 5–8. doi:10.4103/jiaps.JIAPS
- Volpert, D., White, F., Finegold, M.J., Molleston, J., DeBaun, M., & Perlmutter, D.H., 2001. Outcome of early hepatic portoenterostomy for biliary atresia. *J. Pediatr. Gastroenterol. Nutr.* 32: 265–269. doi:10.1097/00005176-200103000-00006
- Wai, C.T., Greenson, J.K., Fontana, R.J., Kalbfleisch, J.D., Marrero, J.A., Conjeevaram, H.S., et al., 2003. A simple noninvasive index can predict both significant fibrosis and cirrhosis in patients with chronic hepatitis C. *Hepatology* 38: 518–526. doi:10.1053/jhep.2003.50346
- Waiman, E., & Oswari, H., 2010. Peran Operasi Kasai pada Pasien Atresia Bilier yang Datang Terlambat. *Sari P* 11: 463–470.
- Wildhaber, B.E., 2012. Biliary Atresia: 50 Years after the First Kasai. *ISRN Surg.* 2012: 1–15. doi:10.5402/2012/132089
- Williams, A.L.B., & Hoofnagle, J.H., 1988. Ratio of Serum Aspartate to Alanine Aminotransferase in Chronic Hepatitis Relationship to Cirrhosis. *Gastroenterology* 95: 734–739. doi:10.1016/S0016-5085(88)80022-2
- Wolkoff, A.W., & Berk, P.D., 2011. Bilirubin Metabolism and Jaundice. *Schiff's Dis. Liver* 120–151. doi:10.1002/9781119950509.ch6
- Wu, E.-T., Chen, H.-L., Ni, Y.-H., Lee, H.-Y., & Chang, M.-H., 2001. Bacterial cholangitis in patients with biliary atresia : impact on short-term outcome. *Pediatr. Surg. Int.* 17: 390–395.
- Xu, Y., Yu, J., Zhang, R., Yin, Y., Ye, J., Tan, L., et al., 2012. The perinatal infection of cytomegalovirus is an important etiology for biliary atresia in China. *Clin. Pediatr. (Phila)*. 51: 109–113. doi:10.1177/0009922811406264
- Yang, L.Y., Fu, J., Peng, X.F., Pang, S.Y., Gao, K.K., Chen, Z.R., et al., 2015.



- Validation of aspartate aminotransferase to platelet ratio for diagnosis of liver fibrosis and prediction of postoperative prognosis in infants with biliary atresia. *World J. Gastroenterol.* 21: 5893–5900. doi:10.3748/wjg.v21.i19.5893
- Yassin, N.A., El-Tagy, G., Abdelhakeem, O.N., Asem, N., & El-Karakasy, H., 2020. Predictors of short-term outcome of kasai portoenterostomy for biliary atresia in infants: A single-center study. *Pediatr. Gastroenterol. Hepatol. Nutr.* 23: 266–275. doi:10.5223/PGHN.2020.23.3.266
- Zabiegaj-Zwick, C., Nel, E., & Moore, S.W., 2012. Problems related to cytomegalovirus infection and biliary atresia. *South African Med. J.* 102: 890–892. doi:10.7196/SAMJ.6163
- Zani, A., Quaglia, A., Hadzić, N., Zuckerman, M., & Davenport, M., 2015. Cytomegalovirus-associated biliary atresia: An aetiological and prognostic subgroup. *J. Pediatr. Surg.* 50: 1739–1745. doi:10.1016/j.jpedsurg.2015.03.001
- Zhao, Y., Xu, X., Liu, G., Yang, F., & Zhan, J., 2021. Prognosis of Biliary Atresia Associated With Cytomegalovirus: A Meta-Analysis. *Front. Pediatr.* 9: 1–9. doi:10.3389/fped.2021.710450