

DAFTAR PUSTAKA

- Adhikari, K.C., Malla, R., Maskey, A., Rajbhandari, S., Baral, B.R., Budhathoki, A., *et al.* (2021). Contrast Induced Nephropathy and its predictors after Primary Percutaneous Intervention. *NHJ* 18 : 27–31.
- Ali, A., Bhan, C., Malik, M.B., Ahmad, Q., Sami, S.A., & Ahmad, M.Q. (2018). The Prevention and Management of Contrast-induced Acute Kidney Injury: A Mini-review of the Literature. *Cureus* 10(9):e328.
- Andreucci, M., Faga, T., Pisani, A., Sabbatini, M., Russo, D., & Michael, A. (2014). The Choice of the Iodinated Radiographic Contrast Media to Prevent Contrast-Induced Nephropathy. *Adv. Nephrol.* 2014 : 1–11.
- Andreucci, M., Faga, T., Riccio, E., Sabbatini, M., Pisani, A., & Michael, A. (2016). The potential use of biomarkers in predicting contrast-induced acute kidney injury. *Int J Nephrol Renov. Dis* 9 : 205–221.
- Aurelio, A., & Durante, A. (2014). Contrast-Induced Nephropathy in Percutaneous Coronary Interventions : Pathogenesis , Risk Factors , Outcome , Prevention and Treatment 62–72.
- Chillo, P., Malaja, N., & Kisenge, P. (2021). Magnitude and associated factors of contrast induced nephropathy among patients undergoing coronary angiography and interventions at a cardiac referral hospital in Tanzania - a cross-sectional study. *Pan Afr Med J* 38 : 311.
- Hidayati L dan Maradhona Y (2018). Six Sigma For Evaluation Of Quality Control in Clinical Laboratory. *Int. J. Public Heal. Clin. Sceciences* 5 : 144–150.
- Kalantar-Zadeh, K., & Amin, A.N. (2012). Toward more accurate detection and risk stratification of chronic kidney disease. *Jama* 307 : 1976–1977.
- Kandarini, Y., Wulandari, P.J., & Mahadita, G.W. (2021). Faktor resiko contrast induced nephropathy pada pasien yang menjalani prosedur coronary intervention. *Udayana J. Intern. Med.* 5 : 20–25.
- Kang, F., Zhang, C., Wang, W., & Wang, Z. (2016). Sigma metric analysis for performance of creatinine with fresh frozen serum. *Scand. J. Clin. Lab. Invest.* 76 : 40–44.
- Karkalousos, P., & Evangelopoulos, A. (2011). Quality Control in Clinical Laboratories <http://www.intechopen.com/books/applications-and-e>.
- Kashani, K., Cheungpasitporn, W., & Ronco, C. (2017). Biomarkers of acute kidney injury: The pathway from discovery to clinical adoption. *Clin. Chem. Lab. Med.* 55 : 1074–1089.
- Korolov, Y., Nogués, I., Gambarte, M.J., Donato, M.S., Ciambrone, M.G., Novo, F., *et al.* (2021). Microalbuminuria predicts contrast-induced nephropathy in patients with acute coronary syndrome. *REC Interv. Cardiol.* 3 : 21–35.

- Kumar, B.V., & Mohan, T. (2018). Sigma metrics as a tool for evaluating the performance of internal quality. *J. Lab. Physicians* 10 : 194–199.
- Kurniati, A., & Tahono, T. (2016). ANGKA BANDING ALBUMIN KREATININ AIR KEMIH DAN HbA1C SERTA ESTIMASI LAJU FILTRASI GLOMERULUS PADA PASIEN DIABETES MELITUS TIPE 2. *Indones. J. Clin. Pathol. Med. Lab.* 21 : 250.
- Levey, A., Stevens, L., & Schmid, C. (2009). A New Equation to Estimate Glomerular Filtration Rate. *J. Nephrol. Ther.* 06 : 604–612.
- Levey, A.S., Becker, C., & Inker, L.A. (2015). Glomerular filtration rate and albuminuria for detection and staging of acute and chronic kidney disease in adults: A systematic review. *JAMA - J. Am. Med. Assoc.* 313 : 837–846.
- Levey, A.S., Stevens, L.A., Schmid, C.H., Zhang, Y. (Lucy), Castro, A.F., Feldman, H.I., *et al.* (2009). Evaluation of the CKD-EPI Equation in Multiple Races and Ethnicities. *Ann. Intern. Med.* 150 : 604.
- Li, Y., Ma, K., Shen, G., Zheng, D., Xuan, Y., & Li, W. (2021). Impact of small and dense low-density lipoprotein (sd-LDL) on contrast-induced acute kidney injury in patients with acute ST-segment elevation myocardial infarction undergoing primary percutaneous coronary intervention. *Int. Urol. Nephrol.* 53 : 2611–2617.
- Marenzi Giancarlo , Emilio Assanelli, Jeness Campodonico, Gianfranco Lauri, Ivana Marana, Monica De Metrio, Marco Moltrasio, Marco Grazi, Mara Rubino, Fabrizio Veglia, Franco Fabbiochi, and A.L.B. (2009). Article Annals of Internal Medicine Contrast Volume During Primary Percutaneous Coronary Intervention. *Development D* : 170–177.
- McCullough, P. (2006). Outcomes of contrast-induced nephropathy: Experience in patients undergoing cardiovascular intervention. *Catheter. Cardiovasc. Interv.* 67 : 335–343.
- McCullough, P.A., Choi, J.P., Feghali, G.A., Schussler, J.M., Stoler, R.M., Vallabahn, R.C., *et al.* (2016). Contrast-Induced Acute Kidney Injury. *J. Am. Coll. Cardiol.* 68 : 1465–1473.
- Mehran R, Dangas, G.D., & Weisbord, S.D. (2019). Contrast-Associated Acute Kidney Injury. *N Engl J Med* 380 : 2146–2155.
- National Kidney Foundation - Kidney Disease Improving Global Outcome (2012). KDIGO Clinical Practice Guideline for Acute Kidney Injury. *Kidney Int. Suppl.* 2 : 69–88.
- Ningrum, N.R., & Yuniadi, Y. (2009). Contrast Induced Nephropathy Tinjauan Pustaka Contrast Induced Nephropathy 30 : 71–79.
- Nunes, M.B.G., Filho, A.C., Alvares, V.R.C., Meneguz-Moreno, R., Lamas, E., Loures, V., *et al.* (2018). CKD-EPI versus Cockcroft-Gault formula for

predicting contrast-induced nephropathy following percutaneous coronary intervention in patients without significant renal impairment. *Rev Port Cardiol.* 37 : 25–33.

Ontario Association of Medical Laboratories (2015). Community Laboratory Guidelines 1–6.

Oweis, A.O., Alshelleh, S.A., Saadeh, N., Jarrah, M.I., Ibdah, R., & Alzoubi, K.H. (2020). Long-Term Follow-Up of Contrast-Induced Acute Kidney Injury: A Study from a Developing Country. *Int. J. Vasc. Med.* 2020 : 15–20.

Park, H.S., Kim, C.J., Hwang, B.H., Kim, T.H., Koh, Y.S., Park, H.J., *et al.* (2016). HDL Cholesterol Level Is Associated with Contrast Induced Acute Kidney Injury in Chronic Kidney Disease Patients Undergoing PCI. *Sci. Rep.* 6 : 2–7.

PERKI (2021). Pedoman Nasional Pelayanan Kedokteran Intervensi Koroner Perkutan. *J. Kardiol. Indones.* 22–28.

PERKI (2018). Pedoman Tata Laksana Sindrom Koroner Akut. *J. Kardiol. Indones.* 44–59.

PERKI (2016). Panduan praktik klinis (ppk) dan clinical pathway (cp) penyakit jantung dan pembuluh darah.

Rakowski, T., Dziewierz, A., Węgiel, M., Siudak, Z., Zasada, W., Jąkała, J., *et al.* (2022). Risk factors of contrast-induced nephropathy in patients with acute coronary syndrome. *Kardiol. Pol.* 80 : 760–764.

Rakowski, T., Dziewierz, A., Węgiel, M., Siudak, Z., Zasada, W., Jąkała, J., *et al.* (2021). Risk factors of contrast induced nephropathy in patients with acute coronary syndromes. *Kardiol Pol.* 0–4.

Rear, R., Bell, R.M., & Hausenloy, D.J. (2016). Contrast-induced nephropathy following angiography and cardiac interventions. *Heart* 102 : 638–648.

Ricos, C., Alvarez, V., & Cava, F. (2014). Desirable Biological Variation Database specifications [WWW Document]. *Westgard QC*.

Roche Diagnostics Cobas (2010). Creatinine Jaffé Gen.2 1–5.

Ronco, F., Tarantini, G., & McCullough, P.A. (2020). Contrast induced acute kidney injury in interventional cardiology: An update and key guidance for clinicians. *Rev. Cardiovasc. Med.* 21 : 9–23.

Stevens, L.A., Li, S., Kurella Tamura, M., Chen, S.C., Vassalotti, J.A., Norris, K.C., *et al.* (2011). Comparison of the CKD Epidemiology Collaboration (CKD-EPI) and Modification of Diet in Renal Disease (MDRD) study equations: Risk factors for and complications of CKD and mortality in the Kidney Early Evaluation Program (KEEP). *Am. J. Kidney Dis.* 57 : S9.

Toprak, O., & Cirit, M. (2006). Risk factors for contrast-induced nephropathy. *Kidney Blood Press. Res.* 29 : 84–93.

Wang, C., Ma, S., Deng, B., Lu, J., Shen, W., Jin, B., *et al.* (2017). The predictive value of the product of contrast medium volume and urinary

albumin/creatinine ratio in contrast-induced acute kidney injury. *Ren. Fail.* 39 : 555–560.

Watanabe, M., Saito, Y., Aonuma, K., Hirayama, A., Tamaki, N., Tsutsui, H., *et al.* (2016). Prediction of contrast-induced nephropathy by the serum creatinine level on the day following cardiac catheterization. *J. Cardiol.* 68 : 412–418.

Weisbord, S.D. (2018). Contrast-Associated Acute Kidney Injury. *Springer Nature, Core Concepts Acute Kidney Inj.*

Westgard, S., & Westgard, Q. (2020). Six Sigma Metric Analysis for Analytical Testing Processes. *Diagnostics* 1–9.

Yuniadi, Y., & Ningrum, N.R. (2008). Risk factors and incidence of contrast induced nephropathy following coronary intervention. *Med. J. Indones.* 17 : 131–137.

Zhang, F., Lu, Z., & Wang, F. (2020). Advances in the pathogenesis and prevention of contrast-induced nephropathy. *Life Sci.* 259 : 118379.

Zhou, S., Yuan, G., Huang, B., Geng, D., Lei, J., Zhang, S., *et al.* (2020). Feasibility of Using eGFR to Diagnose Contrast-Induced Nephropathy in Patients Undergoing Coronary Angiography. *J. Integr. Cardiol. Open Access* 8620813324 : 1–6.