

DAFTAR PUSTAKA

- Abdo, R. A., Mosa, H., Sultan, M., Helill, S. E., Assefa, B., Abdu, M., Ahmed, U., Abose, S., Nuramo, A. A., Alemu, A., Demelash, M., & Delil, R. (2021). Prevalence and risk factors associated with birth asphyxia among neonatus delivered in Ethiopia: A systematic review and meta-analysis. *PLOS ONE*, *16*(8), e0255488. <https://doi.org/10.1371/journal.pone.0255488>
- Admasu, F. T., Melese, B. D., Amare, T. J., Zewude, E. A., Denku, C. Y., & Dejenie, T. A. (2022). The magnitude of neonatal asphyxia and its associated factors among newborns in public hospitals of North Gondar Zone, Northwest Ethiopia: A cross-sectional study. *PLoS ONE*, *17*(3 March), 1–10. <https://doi.org/10.1371/journal.pone.0264816>
- Agbeko, F., Kétévi, A. A., Fiawoo, M., Tata, B. B.-L., Abalo, K. E., Takassi, E. O., Douaguibe, B., Akolly, D. A. E., Batalia, H. D., Djomaleu, R. A., Kérékou, R. B., Hemou, M., Pakoudjare, M., Nzonou, M., Sewu, E. K., Talboussouma, S., Saka, B., Azoumah, D. K., Djadou, E. K., ... Atakouma, Y. D. (2021). Risk Factors for Birth Asphyxia in Togo: A Case-Control Study. *Open Journal of Pediatrics*, *11*(04), 816–831. <https://doi.org/10.4236/ojped.2021.114077>
- Ahmad, ZF., Nurdin, SSI. -. Faktor Risiko Kejadian Preeklampsia di RSIA Siti Khadijah Gorontalo
- Ahmed, S. (2020). Risk Factors of Birth Asphyxia Among Neonatus Born in Public Hospitals of Tigray , Northern. 13–20.
- Aini, F. N., Widyawati, M. N., & Suryono, S. (2019). Early Detection of Preeclampsia using a Rule-Based System Information System. *Journal of Physics: Conference Series*, *1179*, 012139. doi:10.1088/1742-6596/1179/1/012139
- American College of Obstetricians and Gynecologists, & American Academy of Pediatrics. (2004). Neonatal Encephalopathy and Cerebral Palsy: Executive Summary*. *Obstetris & Gynecology*, *103*(4), 780–781. <https://doi.org/10.1097/01.AOG.0000120142.83093.30>
- Ameyaw, E. K., Njue, C., Tran, N. T., & Dawson, A. (2020). Quality and women’s satisfaction with maternal referral practices in sub-Saharan African low and lower-middle income countries: a systematic review. *BMC Pregnancy and Childbirth*, *20*(1). doi:10.1186/s12884-020-03339-3
- Aslam, H. M. uhamma., Saleem, S., Afzal, R., Iqbal, U., Saleem, S. Muhammad, Shaikh, M. W. aqa. A., & Shahid, N. (2014). “Risk factors of birth asphyxia.” *Italian journal of pediatrics*, *40*, 94. <https://doi.org/10.1186/s13052-014-0094-2>
- Asres, A. W., Daga, W. B., Samuel, S., Adella, G. A., Workie, S. B., & Desalegn, A. (2022). Determinants of Preeclampsia among Pregnant Women in Public Hospitals of Wolaita and Dawuro Zones, Southern Ethiopia: A Case-Control

Study. *Women's Health Reports*, 3(1), 964–970. <https://doi.org/10.1089/whr.2022.0043>

- Ayele, A. D., & Tilahun, Z. A. (2022). Determinants of pre-eclampsia among women attending delivery services in public health institutions of Debre Tabor Town: a case-control study. *Reproductive Health*, 19(1), 1–8. <https://doi.org/10.1186/s12978-022-01463-1>
- Azizah, I., & Handayani, O. K. (2017). Kematian Neonatal di Kabupaten Grobogan. *HIGEIA*, 1(4). <http://journal.unnes.ac.id/sju/index.php/higeia>
- Azwar, Azrul. (1996). *Menjaga Mutu Pelayanan Kesehatan*. Jakarta: Pustaka Sinar Harapan.
- Azwar, Azrul. (1996). *Pengantar Administrasi Kesehatan*. Edisi Kedua Jakarta: Pt. Binariya Aksara.
- Babore GO, Aregago TG, Ermolo TI, Nunemo MH, Habebo TT. (2021). Determinants of Pregnancy-Induced Hypertension on Maternal and Foetal Outcomes in Hossana Town Administration, Hadiya Zone, Southern Ethiopia: Unmatched Case-Control Study. *Plos One* 16(5): E0250548. <https://doi.org/10.1371/Journal.Pone.0250548>
- Badan Pusat Statistik. (2023). *Angka Kematian Bayi/AKB (Infant Mortality Rate/IMR) Menurut Provinsi, 1971-2020*. <https://www.bps.go.id/statictable/2023/03/31/2216/angka-kematian-bayi-akb-infant-mortality-rate-imr-menurut-provinsi-1971-2020.html>
- Bailit, J. L., Garrett, J. M., Miller, W. R., McMahan, M. T., & Cefalo, R. C. (2002). Hospital primary cesarean delivery rates and the risk of poor neonatal outcomes. *American Journal of Obstetrics and Gynecology*, 187(3), 721–727. <https://doi.org/10.1067/mob.2002.125886>
- Bayih, W. A., Yitbarek, G. Y., Aynalem, Y. A., Abate, B. B., Tesfaw, A., Ayalew, M. Y., Belay, D. M., Hailemeskel, H. S., & Alemu, A. Y. (2020). Prevalence and associated factors of birth asphyxia among live births at Debre Tabor General Hospital, North Central Ethiopia. *BMC Pregnancy and Childbirth*, 20(1), 1–12. <https://doi.org/10.1186/s12884-020-03348-2>
- BPS Kepulauan Riau. (2012). *Angka Kematian Neonatal (AKN) (Per 1.000 Kelahiran Hidup), 2012*. <https://kepri.bps.go.id/indicator/30/293/1/angka-kematian-neonatal-akn-.html>
- Brooks, M. I., Thabrany, H., Fox, M. P., Wirtz, V. J., Feeley, F. G., & Sabin, L. (2017). Health facility and skilled birth deliveries among poor women with Jamkesmas health insurance in Indonesia: a mixed-methods study. *BMC Health Services Research*, 17(1). <https://doi.org/10.1186/s12913-017-2028-3>
- Budihardjo, S.N., & Suryawan, I.W. (2019). Relationship Between Parity with Neonatal Asphyxia in Wangaya General Regional Hospital Denpasar.
- Beketie, E. D., Tafese, W. T., Shiferaw, B. Z., Tilahun, G. A., Gebretsadik, M. A., Suraj, K. D., Mengistie, B. T., Ali, F. Y., Assefa, Z. M., Berriera, F. W., & Teke, N. E. (2022). Determinants of preeclampsia among mothers attending perinatal care in Gurage zone public hospitals, Ethiopia, matched case control

- study. *International Journal of Africa Nursing Sciences*, 17, 100453. <https://doi.org/10.1016/j.ijans.2022.100453>
- Creanga, A. A., Gillespie, D., Karklins, S., & Tsui, A. O. (2011). Low use of contraception among poor women in Africa: an equity issue. *Bulletin of the World Health Organization*, 89(4), 258–266. <https://doi.org/10.2471/blt.10.083329>
- Diba, F., Ichsan, I., Muhsin, M., Marthoenis, M., Sofyan, H., Andalas, M., Monfared, I. G., Richert, K., Kaplan, L., Rogge, L., Doria, S., Samadi, S., & Vollmer, S. (2019). Healthcare providers' perception of the referral system in maternal care facilities in Aceh, Indonesia: a cross-sectional study. *BMJ Open*, 9(12), e031484. <https://doi.org/10.1136/bmjopen-2019-031484>
- Dinas Kesehatan Provinsi Kepulauan Riau. (2020). *Profil Kesehatan Provinsi Kepulauan Riau 2019*.
- Dinas Kesehatan Provinsi Kepulauan Riau. (2022). *Profil Kesehatan Provinsi Kepulauan Riau 2021*.
- Dinas Kesehatan Provinsi Kepulauan Riau. (2021). *Rencana Strategis (RENSTRA) Dinas Kesehatan Provinsi Kepulauan Riau Tahun 2021 – 2026*.
- Ekasari, W. U. (2015). Pengaruh umur ibu, paritas, usia kehamilan, dan berat lahir bayi terhadap asfiksia bayi pada ibu pre eklamsia berat. Universitas Sebelas Maret.
- Elmusharaf, K., Byrne, E., & O'Donovan, D. (2015). Strategies to increase demand for maternal health services in resource-limited settings: challenges to be addressed. *BMC Public Health*, 15(1). doi:10.1186/s12889-015-2222-3
- Eyler, A. (2021). *Research Methods for Public Health*. Springer Publishing Company.
- Faiqoh E, Hendrati Ly. (2014). Hubungan Karakteristik Ibu, Anc Dan Kepatuhan Perawatan Ibu Hamil Dengan Terjadinya Preeklampsia, 2014. *Jurnal Berkala Epidemiologi*. Vol. 2. No. 2 Mei. Pp.: 216–226
- Fatma, Yulia, Martalena Br. Purba. (2010)._ .
- Fatmawati, L., Sulistyono, A., & Notobroto, H. B. (2017). Pengaruh Status Kesehatan Ibu Terhadap Derajat Preeklampsia/Eklampsia di Kabupaten Gresik. *Buletin Penelitian Sistem Kesehatan*, 20(2), 52–58.
- Fitramaya. Pohan, I. S. (2006). *Jaminan Mutu Layanan Kesehatan*. Jakarta: EGC.
- Fitriana, Y., M, A. I., Mallongi, A., Mappajanci, M., Seweng, A., Hidayanty, H., Nur, R., & Syam, A. (2021). Risk factors for asphyxia neonatorum in public health centres of Nosarara and Pantoloan, Palu City. *Gaceta Sanitaria*, 35, S131–S134. <https://doi.org/10.1016/j.gaceta.2021.07.009>
- Foeh, A. Y., Tibuludji, P., & Ludji, I. D. R. (2021). Risk Factors for Neonatal Mortality Due to Birth Asphyxia in East Sumba, East Nusa Tenggara. *The 4th International Conference on Public Health*, 121(5), 26911. <https://doi.org/10.26911/theicph.2018.03.01>

- Fondjo, L. A., Amoah, B., Tashie, W., & Annan, J. J. (2022). Risk factors for the development of new-onset and persistent postpartum preeclampsia: A case-control study in Ghana. *Women's Health, 18*. <https://doi.org/10.1177/17455057221109362>
- Give, C., Ndima, S., Steege, R., Ormel, H., McCollum, R., Theobald, S., ... Sidat, M. (2019). Strengthening referral systems in community health programs: a qualitative study in two rural districts of Maputo Province, Mozambique. *BMC Health Services Research, 19*(1). doi:10.1186/s12913-019-4076-3
- Goodman, M. S., & Thompson, V. S. (2017). Public Health Research Methods for Partnerships and Practice. In *Public Health Research Methods for Partnerships and Practice*. <https://doi.org/10.1201/9781315155722>
- Grum, T., Seifu, A., Abay, M., Angesom, T., & Tsegay, L. (2017). Determinants of pre-eclampsia/Eclampsia among women attending delivery Services in Selected Public Hospitals of Addis Ababa, Ethiopia: A case control study. *BMC Pregnancy and Childbirth, 17*(1), 1–7. <https://doi.org/10.1186/s12884-017-1507-1>
- Guduri, G. B., Bhimarasetty, D. M., Sreegiri, S., & B, T. N. (2015). Socio-Demographic Determinants for Hypertension in Pregnancy A Case-Control Study in A Tertiary Care Hospital of Visakhapatnam, Andhra Pradesh. (1), 93–98.
- Hagaman, A. K., Singh, K., Abate, M., Alemu, H., Kefale, A. B., Bitewulign, B., ... Magge, H. (2020). The impacts of quality improvement on maternal and newborn health: preliminary findings from a health system integrated intervention in four Ethiopian regions. *BMC Health Services Research, 20*(1). doi:10.1186/s12913-020-05391-3
- Hamzah, S. T. R., Aminuddin, Idris, I., & Rachmat, M. (2021). Antenatal care parameters that are the risk factors in the event of preeclampsia in primigravida. *Gaceta Sanitaria, 35*, S263–S267. <https://doi.org/10.1016/j.gaceta.2021.10.073>
- Harrisma. (2018). *Pengaruh Antenatal Care (ANC) dan Rujukan Terhadap Luaran Maternal dan Perinatal Ibu Preeklampsia-Eklampsia di RSUD Waluyo Jati Kabupaten Probolinggo*. Poltekkes Kemenkes Semarang: Prodi DIV Kebidanan Semarang Poltekkes Kemenkes Semarang.
- Haslan, H., & Trisutrisno, I. (2022). Dampak Kejadian Preeklampsia dalam Kehamilan Terhadap Pertumbuhan Janin Intrauterine. *Jurnal Ilmiah Kesehatan Sandi Husada, 11*, 445–454. <https://doi.org/10.35816/jiskh.v11i2.810>
- Hidayati, R., & Setyorini, D. (2020). Multi Level Education Katoga Improve The Competence of Health Cadres, Public Figure, and Family in Preventing, Early Detection and Handling Pregnancy Emergency. *Indonesian Nursing Journal of Education and Clinic (Jurnal Pendidikan Dan Praktik Keperawatan Indonesia)*. <https://doi.org/10.24990/injec.v4i2.242>
- Hijazi, H., Alyahya, M. S., Sindiani, A., Saqan, R. S., & Okour, A. (2018). Determinants of antenatal care attendance among women residing in highly

- disadvantaged communities in northern Jordan: a cross-sectional study. *Reproductive Health*, 15(1). <https://doi.org/10.1186/s12978-018-0542-3>
- Hyre, A., Caiola, N., Amelia, D., Gandawidjaja, T., Markus, S., & Baharuddin, M. (2019). Expanding Maternal and Neonatal Survival in Indonesia: A program overview. *International Journal of Gynecology & Obstetrics*, 144, 7–12. doi:10.1002/ijgo.12730
- Ibrahim, N. A., Muhye, A., & Abdulie, S. (2017). Prevalence of Birth Asphyxia and Associated Faktors among Neonatus Delivered in Dilchora Referral Hospital, in Dire Dawa, Eastern Ethiopia. *Clinics in Mother and Child Health*, 14(4). <https://doi.org/10.4172/2090-7214.1000279>
- Jafar, N., & Lestari, R. (2017). Faktor Risiko Kejadian Preeklampsia Di Rsia Pertiwi Makassar. *Health Care Media*, 3(1), 25 - 31.
- Juban, N., Reyes, J., & Ongkeko, A. (2018). *Inter-Island Health Service Boat Project*. Social Innovation in Health Initiative Case Collection. [Online] WHO, Geneva: Social Innovation in Health Initiative, Available at: www.socialinnovationinhealth.org/case-studies/inter-island-health-service-boat-project/
- Katore Fh, Gurara Am, Beyen Tk. (2021). Determinants of Preeclampsia Among Pregnant Women in Chiro Referral Hospital, Oromia Regional State, Ethiopia: Unmatched Case–Control Study. *Integr Blood Press Control*. 14:163-172. <https://doi.org/10.2147/Ibpc.S336651>
- Kawuryan, S. L. (2004). Pengaruh kadar trombosit, hematokrit, hemoglobin darah dan protein urin pada ibu preeklamsi /eklamsi terhadap nilai APGAR bayi yang dilahirkan. *Jurnal Kedokteran Brawijaya*, XX(2), 78–81.
- Kemendes RI. (2015). *Rencana strategies Kementerian Kesehatan tahun 2015–2019*. Jakarta, Indonesia: Kemendes RI.
- Kemendes RI. (2015). *Peraturan Menteri Kesehatan Nomor 33 Tahun 2015 tentang Pedoman Penyusunan Perencanaan Kebutuhan Sumber Daya Manusia Kesehatan*. Jakarta, Indonesia: Kemendes RI.
- Kemendes RI. (2019). Keputusan Menteri Kesehatan Republik Indonesia Tentang Pedoman Nasional Pelayanan Kedokteran Tata Laksana Asfiksia. Jakarta, Indonesia: Kemendes RI.
- Klugman, J., Li, L., Barker, K. M., Parsons, J., & Dale, K. (2019). How are the domains of women’s inclusion, justice, and security associated with maternal and infant mortality across countries? Insights from the Women, Peace, and Security Index. *SSM-Population Health*, 9, 100486. <https://doi.org/10.1016/j.ssmph.2019.100486>
- Kusumaningrum, R. Y., Murti, B., & Prasetya, H. (2019). Low Birth, Prematurity, and Pre-Eclampsia as Risk Faktors of Neonatal Asphyxia. *Journal of Maternal and Child Health*, 04(01), 49–54. <https://doi.org/10.26911/thejmch.2019.04.01.07>

- Laksono, A. D., Rukmini, R., & Wulandari, R. D. (2020). Regional disparities in antenatal care utilization in Indonesia. *PloS one*, *15*(2), e0224006. <https://doi.org/10.1371/journal.pone.0224006>
- Lawn, J. E., Blencowe, H., Oza, S., You, D., Lee, A. C. C., Waiswa, P., Lalli, M., Bhutta, Z., Barros, A. J. D., Christian, P., Mathers, C., & Cousens, S. N. (2014). Every newborn: Progress, priorities, and potential beyond survival. *The Lancet*, *384*(9938), 189–205. [https://doi.org/10.1016/S0140-6736\(14\)60496-7](https://doi.org/10.1016/S0140-6736(14)60496-7)
- Lee, A. C. C., Mullany, L. C., Tielsch, J. M., Katz, J., Khatri, S. K., Leclercq, S. C., Adhikari, R. K., Shrestha, S. R., & Darmstadt, G. L. (2008). Risk factors for neonatal mortality due to birth asphyxia in Southern Nepal: A prospective, community-based cohort study. *Pediatrics*, *121*(5), 1–18. <https://doi.org/10.1542/peds.2007-1966>
- Lestary, H., Sugiharti, S., & Mujiati, M. (2018). Sistem Rujukan Maternal dan Neonatal di Daerah Kepulauan (Studi Kasus di Provinsi Papua dan Maluku). *Media of Health Research and Development*. <https://doi.org/10.22435/mpk.v28i2.177>
- Locatelli, A., Lambicchi, L., Incerti, M., Bonati, F., Ferdico, M., Malguzzi, S., Torcasio, F., Calzi, P., Varisco, T., & Paterlini, G. (2020). Is perinatal asphyxia predictable? *BMC Pregnancy and Childbirth*, *20*(1). <https://doi.org/10.1186/s12884-020-02876-1>
- Luti, I., Hasanbasri, M., & Lazuardi, L. (2012). Kebijakan Pemerintah Daerah Dalam Meningkatkan Sistem Rujukan Kesehatan Daerah Kepulauan di Kabupaten Lingga Provinsi Kepulauan Riau. *Jurnal Kebijakan Kesehatan Indonesia*. <http://jurnal.ugm.ac.id/jkki/article/download/3072/2728>
- Mamo, S. A., Teshome, G. S., Tesfaye, T., & Tibebu, A. (2022). Perinatal asphyxia and associated factors among neonatus admitted to a specialized public hospital in South Central Ethiopia: A retrospective cross-sectional study. *PLOS ONE*, *17*(1), e0262619. <https://doi.org/10.1371/journal.pone.0262619>
- Mareg, M., Molla, A., Dires, S., Mamo, Z. B., & Hagos, B. (2020). Determinants of preeclampsia among pregnant mothers attending antenatal care (ANC) and delivery service in Gedeo Zone, Southern Ethiopia: Case control-study. *International Journal of Women's Health*, *12*, 567–575. <https://doi.org/10.2147/IJWH.S251342>
- Masruroh, N. (2021). Determinan Maternal Kejadia Preeklampsia Pada Ibu Hamil Trimester III di RS Prima Husada Sidoarjo. *Jurnal Bidang Ilmu Kesehatan*, *11*(1), 94–104. <https://doi.org/10.52643/jbik.v11i1.1072>
- Matyas, M., Hășmășanu, M. G., Silaghi, C., Samasca, G., Lupan, I., Orsolya, K., & Zaharie, G. (2022). Early preeclampsia effect on preterm newborns outcome. *Journal of Clinical Medicine*, *11*(2), 452. <https://doi.org/10.3390/jcm11020452>
- McCarthy, J., & Maine, D. (1992). A Framework for Analyzing the Determinants of Maternal Mortality. *Studies in Family Planning*, *23*(1), 23. <https://doi.org/10.2307/1966825>

- MacDonald, T. M., Walker, S., Hannan, N. J., Tong, S., & Kaitu'u-Lino, T. J. (2022). Clinical tools and biomarkers to predict preeclampsia. *EBioMedicine*, 75, 103780. <https://doi.org/10.1016/j.ebiom.2021.103780>
- Mekie, M., Bezie, M., Melkie, A., Addisu, D., Chanie, E. S., Bayih, W. A., Biru, S., Hailie, M., Seid, T., Dagne, E., Muche, T., & Alemu, E. M. (2022). Perception towards preeclampsia and perceived barriers to early health-seeking among pregnant women in selected Hospitals of South Gondar Zone, Northwest Ethiopia: A qualitative study. *PLoS ONE*, 17(8 August), 1–12. <https://doi.org/10.1371/journal.pone.0271502>
- Meshram, R. M., & Bokade, C. M. (2019). Meshram RM, Bokade CM. Risk factors for mortality in birth asphyxia of outborn neonatus: A prospective observational study. *Sri Lanka Journal of Child Health*, 48(1), 26–32.
- Mikat, B., Gellhaus, A., Wagner, N. N., Birdir, C., Kimmig, R., & Köninger, A. (2012). Early detection of maternal risk for preeclampsia. *ISRN Obstetris and Gynecology (Print)*, 2012, 1–7. <https://doi.org/10.5402/2012/172808>
- Mirzakhani, K., Khadivzadeh, T., Faridhosseini, F., & Ebadi, A. (2020). Pregnant Women's Experiences of the Conditions Affecting Marital Well-Being in High-Risk Pregnancy: A Qualitative Study. *PubMed*, 8(4), 345–357. <https://doi.org/10.30476/ijcbnm.2020.85666.1285>
- Moshiro, R., Mdoe, P., & Perlman, J. M. (2019). A Global View of Neonatal Asphyxia and Resuscitation. *Frontiers in Pediatrics*, 7(November), 1–6. <https://doi.org/10.3389/fped.2019.00489>
- Mou, A. D., Barman, Z., Hasan, M., Miah, R., Hafsa, J. M., Das Trisha, A., & Ali, N. (2021). Prevalence of preeclampsia and the associated risk factors among pregnant women in Bangladesh. *Scientific Reports*, 11(1), 1–9. <https://doi.org/10.1038/s41598-021-00839-w>
- Mulugeta, T., Sebsibe, G., Fenta, F. A., & Sibhat, M. (2020). Risk factors of perinatal asphyxia among newborns delivered at public hospitals in Addis Ababa, Ethiopia: Case-control study. *Pediatric Health, Medicine and Therapeutics*, 11, 297–306. <https://www.dovepress.com/getfile.php?fileID=61023>
- Muninjaya, A. A. G. (2011). *Manajemen Mutu Pelayanan Kesehatan*. Jakarta: EGC.
- Mutiara, A., Apriyanti, F., & Alini, A. (2020). Hubungan Jenis Persalinan dan Berat Badan Lahir Dengan Kejadian Asfiksia Pada Bayi Baru Lahir di RSUD Selasih Kabupaten Pelalawan Tahun 2019. *Jurnal Kesehatan Tambusai*, 1(2), 42–49. <https://doi.org/10.31004/jkt.v1i2.1104>
- Negrini, R., Dalbuquerque, I., De Cássia Sanchez E Oliveira, R., Da Silva Ferreira, R. D., De Stefani, L. F. B., & Podgaec, S. (2021). Strategies to reduce the caesarean section rate in a private hospital and their impact. *BMJ Open Quality*, 10(3), e001215. <https://doi.org/10.1136/bmjopen-2020-001215>

- Nguyen, R. H. N., & Wilcox, A. J. (2005). Terms in reproductive and perinatal epidemiology: 2. Perinatal terms. *Journal of Epidemiology and Community Health*, 59(12), 1019–1021. <https://doi.org/10.1136/jech.2004.023465>
- Nur, A. F., & Arifuddin, A. (2017). Faktor Risiko Kejadian Preeklampsia Pada Ibu Hamil di RSUD Anutapura Kota Palu. *Healthy Tadulako*, 3(2), 69–75.
- Nur, R., Mamar, S., Demak, I. P. K., Fadhliah, Patui, N. S., Radiah, & Rusydi, M. (2019). Husban Behavior in Pregnancy-Postpartum Care and Wife's Reproductive Health. *KnE Life Sciences*. <https://doi.org/10.18502/cls.v4i13.5265>
- Nuridin, S. S. I., Dunggio, R., Batiti, S., Sondakh, L., & Ahmad, Z. F. (2021). Determinan Kejadian Preeklampsia pada Ibu Hamil di RSUD dr. Mohammad Hosein Palembang. *Journal Midwifery Jurusan Kebidanan Politeknik Kesehatan Gorontalo*, 7(1), 31. <https://doi.org/10.52365/jm.v7i1.318>
- Nurmawati. (2010). *Mutu Pelayanan Kebidanan*. Jakarta: Trans Info Media.
- Nursal, D. G. A., Tamela, P., & Fitrayeni, F. (2017). Faktor Risiko Kejadian Preeklampsia Pada Ibu Hamil di RSUD Dr. M. Djamil Padang Tahun 2014. *Jurnal Kesehatan Masyarakat Andalas*, 10(1), 38–44.
- Nurunnayah, S., & Nurhayati, A. S. (2013). *Mutu Pelayanan Kebidanan*. Yogyakarta:
- Olusanya, B. O., & Solanke, O. A. (2011). Perinatal correlates of delayed childbearing in a developing country. *Archives of Gynecology and Obstetrics*, 285(4), 951–957. <https://doi.org/10.1007/s00404-011-2105-5>
- Opiyo, N., Kingdon, C., Oladapo, O. T., Souza, J. P., Vogel, J. P., Bonet, M., Bucagu, M., Portela, A., McConville, F., Downe, S., Gülmezoglu, A. M., & Betrán, A. P. (2019). Non-clinical interventions to reduce unnecessary caesarean sections: WHO recommendations. *Bulletin of the World Health Organization*, 98(1), 66–68. <https://doi.org/10.2471/blt.19.236729>
- Otchere, S. A., Omunoyidde, S., Rosales, A., Ochieng, J. A., Chebon, L., Agordoh, S. W., & Allison, A. (2023). Faith Leaders Improve Healthy Timing and Spacing of Pregnancy: Results of Operations Research on the Channels of Hope Methodology in Kenya and Ghana. *Annals of Global Health*, 89(1), 19. <https://doi.org/10.5334/aogh.3944>
- Park, H., Shim, S. S., & Hyun, D. C. (2015). Combined screening for early detection of Pre-Eclampsia. *International Journal of Molecular Sciences*, 16(8), 17952–17974. <https://doi.org/10.3390/ijms160817952>
- Poma, P. (1998). Effect of Departmental Policies on Cesarean Delivery Rates: A Community Hospital Experience. *Obstetrics & Gynecology*, 91(6), 1013–1018. doi:10.1016/s0029-7844(98)00077-5
- Poon, L. C., Shennan, A., Hyett, J., Kapur, A., Hadar, E., Divakar, H., McAuliffe, F. M., Da Silva Costa, F., Von Dadelszen, P., McIntyre, H. D., Kihara, A. B., Di Renzo, G. C., Romero, R., D'Alton, M. E., Berghella, V., Nicolaides, K. H., & Hod, M. (2019). The International Federation of Gynecology and Obstetrics (FIGO) initiative on pre-eclampsia: A pragmatic guide for first-

- trimester screening and prevention. *International Journal of Gynecology and Obstetrics*, 145(S1), 1–33. <https://doi.org/10.1002/ijgo.12802>
- Poon, L. C., Staboulidou, I., Maiz, N., Plasencia, W., & Nicolaides, K. H. (2009). Hypertensive disorders in pregnancy: screening by uterine artery Doppler at 11-13 weeks. *Ultrasound in Obstetrics & Gynecology*, 34(2), 142–148. <https://doi.org/10.1002/uog.6452>
- Prawirohardjo S. (2012). *Ilmu Kebidanan*. Jakarta: Yayasan Bina Pustaka Sarwono Prawirohardjo.
- Pratama, S. D., Hanum, L., & Handoyo, Y. (2018). Angka Kejadian Asfiksia Neonatorum Pada Bayi Dengan Berat Badan Lahir Rendah di RSUD Goeteng Taroenadibrata Purbalingga. *Herb-medicine Journal*, 1(2). <https://doi.org/10.30595/hmj.v1i2.3098>
- Prayoga, B., Prawitasari, S., & Claramita, M. (2018). Training Detection of Preeclampsia Risk Factors for Physicians in Primary Care with Experience-Based Learning Methods: A Quasi Experimental Study. *Review of Primary Care Practice and Education (Kajian Praktik Dan Pendidikan Layanan Primer)*, 1(2), 82. <https://doi.org/10.22146/rpcpe.36266>
- Rahayuningsih, Fb. & Trisnawati, Ag. (2012). *Hubungan Jarak Kelahiran Dengan Kejadian Pre-Eklamsia*. Yogyakarta: Prosiding Seminar Nasional "Kiat Menjadi Perawat Peneliti Yang Handal
- Rainaldi, M. A., & Perlman, J. M. (2016). Pathophysiology of Birth Asphyxia. *Clinics in Perinatology*, 43(3), 409–422. <https://doi.org/10.1016/j.clp.2016.04.002>
- Sacks, E., Brizuela, V., & Perrotta, C. (2021). It's the Destination and the Journey— A mapping of the challenges in transport and referral for maternal and newborn health in pandemics and beyond. *Frontiers in Public Health*, 9. <https://doi.org/10.3389/fpubh.2021.612409>
- Saifuddin, A. B., Rachimhadhi, T., & Wiknjastro, G. H. (Ed.). (2010). *Ilmu Kebidanan Sarwono Prawirohardjo (3 ed.)*. Bina Pustaka Sarwono Prawirohardjo.
- Saifuddin, A.B. (2006). *Buku Acuan Nasional Pelayanan Kesehatan Maternal Dan Neonatal*. Jakarta: Yayasan Bina Pustaka Sarwono Prawirohardjo
- Saifuddin. (2008). *Ilmu Kebidanan*. Jakarta: Yayasan Bina Pustaka Sarwono Prawirohardjo,
- Sari, P. W., Budihastuti, U. R., & Pamungkasari, E. P. (2019). Path Analysis on the Determinants of Severe Preeclampsia in Surakarta, Central Java. *Journal of Maternal and Child Health*, 4(2), 126–135. <https://doi.org/10.26911/thejmch.2019.04.02.08>
- Satrianegara, M. F. & Saleha, S. (2012). *Buku Ajar Organisasi dan Manajemen Pelayanan Kesehatan Serta Kebidanan*. Jakarta: Salemba Medika.
- Schreier, M. (2012). *Qualitative Content Analysis in Practice*. SAGE Publications, Inc.

- Sekarwati, T. A., Darsini, N., & Husada, D. (2020). Risk Faktors for Neonatal Asphyxia Occurrence at General Hospital Dr. M. Soewandhie, Surabaya. *Indian Journal of Public Health Research & Development*, *11(04)*, 866–871. <https://doi.org/10.37506/ijphrd.v11i4.7757>
- Silva Lm, Coolman M, Steegers Ea, Jaddoe Vw, Moll Ha, Hofman A, *et al.* (2008). Low Socioeconomic Status is A Risk Faktor for Preeclampsia: The Generation R Study. *J Hypertens*. *26(6):1200-8*.
- Soleman, S. R. (2020). The Trends of Neonatal Mortality Rate Among South East Asia Countries from 2000-2017. *Disease Prevention and Public Health Journal*, *14(2)*, 90. <https://doi.org/10.12928/dpphj.v14i2.1912>
- Sondakh, J. J. S., Marjati, & Pipitcahyani, T. I. (2013). *Mutu Pelayanan Kesehatan Dan Kebidanan*. Jakarta: Salemba Medika.
- Sriwandoko, H., Purnomo, W., Trijanto, B., & Darmawan, E. S. (2019). The Effect of Referral and Treatment of Severe Preeclampsia on Maternal Death at Sultan Imanudin General Hospital Pangkalan Bun, Central Kalimantan. 174–182. <https://doi.org/10.26911/the6thicph-fp.03.01>
- Stockwell, E. G., Bedard, M., Swanson, D. A., & Wicks, J. W. (1987). Public policy and the socioeconomic mortality differential in infancy. *Population Research and Policy Review*, *6(2)*, 105–121. <https://doi.org/10.1007/BF00149203>
- Sudarman, Tendean, H. M. M., & Wagey, F. W. (2021). Faktor-Faktor yang Berhubungan dengan Terjadinya Preeklampsia. *e-CliniC*, *9(1)*, 68–80. <https://doi.org/10.35790/ecl.v9i1.31960>
- Suryatini, E., Mamlukah, M., & Wahyuniar, L. (2022). Faktor-Faktor Yang Berhubungan Dengan Kejadian Preeklampsia Pada Ibu Hamil Di Rumah Sakit Umum Daerah (RSUD) Dr. Soekardjo Kota Tasikmalaya Pada Masa Pandemi Covid-19 Tahun 2022. *Journal of Public Health Innovation*, *3(01)*, 1–12. <https://doi.org/10.34305/jphi.v3i01.564>
- Syafrudin, Masitoh S., & Rosyanawati, T. (2011). *Manajemen Mutu Pelayanan Kesehatan Untuk Bidan*. Jakarta: Trans Info Media.
- Taha, I. D., Herini, E. S., & Ismail, D. (2017). Hubungan kejadian asfiksia neonatorum dengan perkembangan anak usia 6 – 24 bulan di RSUD Luwuk Kabupaten Banggai Sulawesi Tengah. *Berita Kedokteran Masyarakat*, *33(5)*, 237. <https://doi.org/10.22146/bkm.17051>
- Tambunan, L. N., Arsesiana, A., & Paramita, A. (2020). Determinant Of Preeclampsia Occurrence In General Hospital Dr . Doris Sylvanus Palangka Raya. *Jurnal Surya Medika*, *6(1)*, 101–111.
- Techane, M. T., Alemu, T. G., Wubneh, C. A., Belay, G. M., Tamir, T. T., Muhye, A., Kassie, D. G., Wondim, A., Terefe, B., Tarekegn, B. G., Ali, M. K., Fentie, B., Gonete, A. T., Tekeba, B., Kassa, S. F., Desta, B. K., Ayele, A. D., Dessie, M. T., Atalell, K. A., & Assimamaw, N. T. (2022). The effect of gestational age, low birth weight and parity on birth asphyxia among neonatus in sub-Saharan Africa: systematic review and meta-analysis: 2021. *Italian Journal of Pediatrics*, *48(1)*. <https://doi.org/10.1186/s13052-022-01307-5>

- Titaley, C.R, Dibley, Michael. J, *et al.* (2008). Determinant of Neonatal Mortality in Indonesia. *UK: BMC Public Health.*
- Titaley, C. R., Hunter, C., Dibley, M. J., & Heywood, P. (2010). Why do some women still prefer traditional birth attendants and home delivery?: a qualitative study on delivery care services in West Java Province, Indonesia. *BMC Pregnancy and Childbirth*, 10(1). <https://doi.org/10.1186/1471-2393-10-43>
- Tolley, E. E., Ulin, P. R., Mack, N., Robinson, E. T., & Succop, S. M. (2016). *Qualitative methods in public health: a field guide for applied research.* Josey Bass.
- Torres-Muñoz, J., Rojas, C., Mendoza-Urbano, D., Marín-Cuero, D., Orobio, S., & Echandía, C. (2017). Risk factors associated with the development of perinatal asphyxia in neonatus at the Hospital Universitario del Valle, Cali, Colombia, 2010-2011. *Biomedica*, 37, 51–56. <https://doi.org/10.7705/biomedica.v37i1.2844>
- UN Inter-agency Group for Child Mortality Estimation. (2022). *Levels & trends in child mortality: report 2022.*
- UNICEF. (2023a). *Countdown to 2030 Country Profile: Indonesia.* Countdown to 2030: Women's Children & Adolescent's Health. <https://data.unicef.org/countdown-2030/country/Indonesia/4/>
- UNICEF. (2023b). *Neonatal mortality.* UNICEF Data: Monitoring the situation of children and women. <https://data.unicef.org/topic/child-survival/neonatal-mortality/#:~:text=The first 28 days of,1%2C000 live births in 1990.>
- Usta, I. M., & Nassar, A. H. (2008). Advanced maternal Age. part I: Obstetri complications. *American Journal of Perinatology*, 25(08), 521–534. <https://doi.org/10.1055/s-0028-1085620>
- Warren, C. E., Hossain, S. M. I., Ishaku, S. M., Armbruster, D., & Hillman, E. (2020). A primary health care model for managing pre-eclampsia and eclampsia in low- and middle- income countries. *Reproductive Health*, 17(1). <https://doi.org/10.1186/s12978-020-0897-0>
- Welembuntu, M., & Gobel, I. (2022). Gambaran Pelaksanaan Sistem Rujukan Puskesmas di Daerah Kepulauan. *Jurnal Ilmiah Sesebanua*, 5(2), 54–59. <https://doi.org/10.54484/jis.v5i2.471>
- WHO. (2012). *Guidelines on basic newborn resuscitation.*
- WHO. (2022). *Newborn Mortality.* Fact Sheets. <https://www.who.int/news-room/fact-sheets/detail/levels-and-trends-in-child-mortality-report-2021>
- WHO. (2023). *SDG Target 3.2 End preventable deaths of newborns and children under 5 years of age.* The Global Health Observatory. https://www.who.int/data/gho/data/themes/topics/sdg-target-3_2-newborn-and-child-mortality
- Workineh, Y., Kasa, A. S., Ayalew, E., Temesgen, W. A., Tirfie, M., & Birhanu, M. (2020). Prevalence of perinatal asphyxia in East and Central Africa:

systematic review and meta-analysis. *Heliyon*, 6(4), e03793.
<https://doi.org/10.1016/j.heliyon.2020.e03793>

World Health Organization, UNDP/UNFPA/UNICEF/WHO/World Bank Special Programme of Research, Development and Research Training in Human Reproduction & Kunnskapscenteret. (2014). *Using auxiliary nurse midwives to improve access to key maternal and newborn health interventions*. World Health Organization. <https://apps.who.int/iris/handle/10665/128037>

Youssef, A. E.-D. A., Mohamed, M. H., Moussa, S. S. A., & Habib, D. M. E.-S. (2018). Effect of Socioeconomic Status on Preeclampsia Cross Sectional Study. *The Medical Journal of Cairo University*, 86(12), 4227–4234. <https://doi.org/10.21608/mjcu.2018.62808>

Yu, Y., Gao, J., Liu, J., Tang, Y., Zhong, M., He, J., Liao, S., Wang, X., Liu, X., Cao, Y., Liu, C., & Sun, J. (2022). Perinatal maternal characteristics predict a high risk of neonatal asphyxia: A multi-center retrospective cohort study in China. *Frontiers in medicine*, 9, 944272. <https://doi.org/10.3389/fmed.2022.944272>

Zakiyya, A., Rusmini, R., & Sartika, Q. L. (2019). Enhancing the pregnant women knowledge through health education about high risk pregnancy. *Jurnal Kesehatan Ibu Dan Anak*, 13(1), 67–73. <https://doi.org/10.29238/kia.v13i1.415>