

## REFERENCE

- Albero, B., Tadeo, J. L., & Pérez, R. A. (2019). Ultrasound-assisted extraction of organic contaminants. In *TrAC - Trends in Analytical Chemistry* (Vol. 118, pp. 739–750). Elsevier B.V. <https://doi.org/10.1016/j.trac.2019.07.007>
- Albuquerque, B. R., Pinela, J., Barros, L., Oliveira, M. B. P. P., & Ferreira, I. C. F. R. (2020). Anthocyanin-rich extract of jabuticaba epicarp as a natural colorant: Optimization of heat- and ultrasound-assisted extractions and application in a bakery product. *Food Chemistry*, 316. <https://doi.org/10.1016/j.foodchem.2020.126364>
- Aliaño-González, M. J., Espada-Bellido, E., Ferreiro-González, M., Carrera, C., Palma, M., Ayuso, J., Álvarez, J. Á., & Barbero, G. F. (2020). Extraction of anthocyanins and total phenolic compounds from Açai (*Euterpe oleracea* Mart.) using an experimental design methodology. Part 2: Ultrasound-assisted extraction. *Agronomy*, 10(3). <https://doi.org/10.3390/agronomy10030326>
- Aliaño-González, M. J., Jarillo, J. A., Carrera, C., Ferreiro-González, M., Álvarez, J. Á., Palma, M., Ayuso, J., Barbero, G. F., & Espada-Bellido, E. (2020). Optimization of a novel method based on ultrasound-assisted extraction for the quantification of anthocyanins and total phenolic compounds in blueberry samples (*Vaccinium corymbosum* L.). *Foods*, 9(12). <https://doi.org/10.3390/foods9121763>
- AOAC. (2016). Guidelines for Standard Method Performance Requirements.
- Berger, P. D., Maurer, R. E., & Celli, G. B. (2018). *Introduction to Response-Surface Methodology*. <https://doi.org/10.1007/978-3-319>
- Blando, F., Calabriso, N., Berland, H., Maiorano, G., Gerardi, C., Carluccio, M. A., & Andersen, Ø. M. (2018). Radical scavenging and anti-inflammatory activities of representative anthocyanin groupings from pigment-rich fruits and vegetables. *International Journal of Molecular Sciences*, 19(1). <https://doi.org/10.3390/ijms19010169>
- Carrera, C., Aliaño-González, M. J., Valaityte, M., Ferreiro-González, M., Barbero, G. F., & Palma, M. (2021). A novel ultrasound-assisted extraction method for the analysis of anthocyanins in potatoes (*Solanum tuberosum* L.). *Antioxidants*, 10(9). <https://doi.org/10.3390/antiox10091375>

- Celli, G. B., Ghanem, A., & Brooks, M. S. L. (2015). Optimization of ultrasound-assisted extraction of anthocyanins from haskap berries (*Lonicera caerulea* L.) using Response Surface Methodology. *Ultrasonics Sonochemistry*, 27, 449–455. <https://doi.org/10.1016/j.ultsonch.2015.06.014>
- Chayati, I., Marsono, Y., & Astuti, M. (2020). The Effect of Varieties, Sifting Factions, and Solvents on Total Anthocyanins, Total Phenolic Contents, and Antioxidant Activities of Purple Corn Extracts. *Jurnal Riset Teknologi Industri*, 14(1).
- Chen, L., Yang, M., Mou, H., & Kong, Q. (2018). Ultrasound-assisted extraction and characterization of anthocyanins from purple corn bran. *Journal of Food Processing and Preservation*, 42(1). <https://doi.org/10.1111/jfpp.13377>
- Chen, X. Q., Li, Z. H., Wang, Z. J., Liu, L. L., Sun, T. T., Ma, J. Z., & Zhang, Y. (2020). Ultrasound-assisted extraction of total anthocyanins from *Rubia sylvatica* Nakai fruit and radical scavenging activity of the extract. *Industrial Crops and Products*, 150. <https://doi.org/10.1016/j.indcrop.2020.112420>
- Colombo, F., Di Lorenzo, C., Petroni, K., Silano, M., Pilu, R., Falletta, E., Biella, S., & Restani, P. (2021). Pigmented corn varieties as functional ingredients for gluten-free products. *Foods*, 10(8). <https://doi.org/10.3390/foods10081770>
- Espada-Bellido, E., Ferreiro-González, M., Carrera, C., Palma, M., Álvarez, J. A., Barbero, G. F., & Ayuso, J. (2019). Extraction of Antioxidants from Blackberry (*Rubus ulmifolius* L.): Comparison between ultrasound- And microwave-assisted extraction techniques. *Agronomy*, 9(11). <https://doi.org/10.3390/agronomy9110745>
- Fernandez-Aulis, F., Hernandez-Vazquez, L., Aguilar-Orsorio, G., Arrieta-Baez, D., & Navarro-Ocana, A. (2019). Extraction and Identification of Anthocyanins in Corn Cob and Corn Husk from Cacahuacintle Maize. *Journal of Food Science*, 84(5), 954–962. <https://doi.org/10.1111/1750-3841.14589>
- Ferreira, S. L. C., Bruns, R. E., Ferreira, H. S., Matos, G. D., David, J. M., Brandão, G. C., da Silva, E. G. P., Portugal, L. A., dos Reis, P. S., Souza, A. S., & dos Santos, W. N. L. (2007). Box-Behnken design: An alternative for the optimization of analytical methods. In *Analytica Chimica Acta* (Vol. 597, Issue 2, pp. 179–186). <https://doi.org/10.1016/j.aca.2007.07.011>

- González, M. J. A., Carrera, C., Barbero, G. F., & Palma, M. (2022). A comparison study between ultrasound-assisted and enzyme-assisted extraction of anthocyanins from blackcurrant (*Ribes nigrum* L.). *Food Chemistry: X*, 13. <https://doi.org/10.1016/j.fochx.2021.100192>
- González-De-Peredo, A. V., Vázquez-Espinosa, M., Espada-Bellido, E., Ferreiro-González, M., Carrera, C., Palma, M., Álvarez, J. Á., Barber, G. F., & Ayuso, J. (2020). Optimization of analytical ultrasound-assisted methods for the extraction of total phenolic compounds and anthocyanins from sloes (*Prunus spinosa* L.). *Agronomy*, 10(7). <https://doi.org/10.3390/agronomy10070966>
- Hong, H. T., Netzel, M. E., & O'Hare, T. J. (2020). Optimisation of extraction procedure and development of LC–DAD–MS methodology for anthocyanin analysis in anthocyanin-pigmented corn kernels. *Food Chemistry*, 319. <https://doi.org/10.1016/j.foodchem.2020.126515>
- Hu, Q. P., & Xu, J. G. (2011). Profiles of carotenoids, anthocyanins, phenolics, and antioxidant activity of selected color waxy corn grains during maturation. *Journal of Agricultural and Food Chemistry*, 59(5), 2026–2033. <https://doi.org/10.1021/jf104149q>
- ICH. (2005). International Conference On Harmonisation Of Technical Requirements For Registration Of Pharmaceuticals For Human Use Ich Harmonised Tripartite Guideline Validation Of Analytical Procedures: Text And Methodology Q2(R1).
- Ikhtiarini, A. N., Setyaningsih, W., Rafi, M., Aminah, N. S., Insanu, M., Imawati, I., & Rohman, A. (2021). Optimization of ultrasound-assisted extraction and the antioxidant activities of Sidaguri (*Sida rhombifolia*). *Journal of Applied Pharmaceutical Science*, 11(8), 70–76. <https://doi.org/10.7324/JAPS.2021.110810>
- Kapasakalidis, P. G., Rastall, R. A., & Gordon, M. H. (2006). Extraction of polyphenols from processed black currant (*Ribes nigrum* L.) residues. *Journal of Agricultural and Food Chemistry*, 54(11), 4016–4021. <https://doi.org/10.1021/jf052999l>
- Khoo, H. E., Azlan, A., Tang, S. T., & Lim, S. M. (2017). Anthocyanidins and anthocyanins: Colored pigments as food, pharmaceutical ingredients, and the potential health benefits. In *Food and Nutrition Research* (Vol. 61). Swedish Nutrition Foundation. <https://doi.org/10.1080/16546628.2017.1361779>

- Kumar, K., Srivastav, S., & Sharanagat, V. S. (2021). Ultrasound assisted extraction (UAE) of bioactive compounds from fruit and vegetable processing by-products: A review. In *Ultrasonics Sonochemistry* (Vol. 70). Elsevier B.V. <https://doi.org/10.1016/j.ultsonch.2020.105325>
- Lieberman, S. (2007). The antioxidant power of purple corn: A research review. In *Alternative and Complementary Therapies* (Vol. 13, Issue 2, pp. 107–110). Mary Ann Liebert Inc. <https://doi.org/10.1089/act.2007.13210>
- López, C. J., Caleja, C., Prieto, M. A., Barreiro, M. F., Barros, L., & Ferreira, I. C. F. R. (2018). Optimization and comparison of heat and ultrasound assisted extraction techniques to obtain anthocyanin compounds from *Arbutus unedo* L. Fruits. *Food Chemistry*, 264, 81–91. <https://doi.org/10.1016/j.foodchem.2018.04.103>
- Moreno, Y. S., Sanchez, G. S., Hernandez, D. R., & Lobato, N. R. (2005). Characterization of Anthocyanin Extracts from Maize Kernels. *Journal of Chromatographic Science*, 43, 483–487.
- Muangrat, R., Pongsirikul, I., & Blanco, P. H. (2018). Ultrasound assisted extraction of anthocyanins and total phenolic compounds from dried cob of purple waxy corn using response surface methodology. *Journal of Food Processing and Preservation*, 42(2). <https://doi.org/10.1111/jfpp.13447>
- Oladzadabbasabadi, N., Mohammadi Nafchi, A., Ghasemlou, M., Ariffin, F., Singh, Z., & Al-Hassan, A. A. (2022). Natural anthocyanins: Sources, extraction, characterization, and suitability for smart packaging. In *Food Packaging and Shelf Life* (Vol. 33). Elsevier Ltd. <https://doi.org/10.1016/j.fpsl.2022.100872>
- Pérez-Gregorio, R. M., García-Falcón, M. S., Simal-Gándara, J., Rodrigues, A. S., & Almeida, D. P. F. (2010). Identification and quantification of flavonoids in traditional cultivars of red and white onions at harvest. *Journal of Food Composition and Analysis*, 23(6), 592–598. <https://doi.org/10.1016/j.jfca.2009.08.013>
- Salacheep, S., Kasemsiri, P., Pongsa, U., Okhawilai, M., Chindaprasirt, P., & Hiziroglu, S. (2020). Optimization of ultrasound-assisted extraction of anthocyanins and bioactive compounds from butterfly pea petals using Taguchi method and Grey relational analysis. *Journal of Food Science and Technology*, 57(10), 3720–3730. <https://doi.org/10.1007/s13197-020-04404-7>

- Seidi, S., & Yamini, Y. (2012). Analytical sonochemistry; Developments, applications, and hyphenations of ultrasound in sample preparation and analytical techniques. In *Central European Journal of Chemistry* (Vol. 10, Issue 4, pp. 938–976). <https://doi.org/10.2478/s11532-011-0160-1>
- Setyaningsih, W., Saputro, I. E., Carrera, C. A., & Palma, M. (2019). Optimisation of an ultrasound-assisted extraction method for the simultaneous determination of phenolics in rice grains. *Food Chemistry*, 288, 221–227. <https://doi.org/10.1016/j.foodchem.2019.02.107>
- Sipahli, S., Mohanlall, V., & Mellem, J. J. (2017). Stability and degradation kinetics of crude anthocyanin extracts from H. sabdariffa. *Food Science and Technology (Brazil)*, 37(2), 209–215. <https://doi.org/10.1590/1678-457X.14216>
- Suriano, S., Balconi, C., Valoti, P., & Redaelli, R. (2021). Comparison of total polyphenols, profile anthocyanins, color analysis, carotenoids and tocopherols in pigmented maize. *LWT*, 144. <https://doi.org/10.1016/j.lwt.2021.111257>
- Tan, J., Han, Y., Han, B., Qi, X., Cai, X., Ge, S., & Xue, H. (2022). Extraction and purification of anthocyanins: A review. *Journal of Agriculture and Food Research*, 8. <https://doi.org/10.1016/j.jafr.2022.100306>
- Teng, Z., Jiang, X., He, F., & Bai, W. (2020). Qualitative and Quantitative Methods to Evaluate Anthocyanins. *EFood*, 1(5), 339–346. <https://doi.org/10.2991/efood.k.200909.001>
- Thuy, N. M., Han, D. H. N., Minh, V. Q., & Tai, N. Van. (2022). Effect of extraction methods and temperature preservation on total anthocyanins compounds of *Peristrophe bivalvis* L. Merr leaf. *Journal of Applied Biology and Biotechnology*, 10(2), 146–153. <https://doi.org/10.7324/JABB.2022.100218>
- Türker, D. A., & Doğan, M. (2022). Ultrasound-assisted natural deep eutectic solvent extraction of anthocyanin from black carrots: Optimization, cytotoxicity, in-vitro bioavailability and stability. *Food and Bioprocess Processing*, 132, 99–113. <https://doi.org/10.1016/j.fbp.2022.01.002>
- Vázquez-Espinosa, M., González de Peredo, A. V., Ferreira-González, M., Carrera, C., Palma, M., Barbero, G. F., & Espada-Bellido, E. (2019). Assessment of ultrasound assisted extraction as an alternative method for the extraction of anthocyanins and total phenolic compounds from maqui berries (*Aristotelia chilensis* (Mol.) Stuntz). *Agronomy*, 9(3). <https://doi.org/10.3390/agronomy9030148>

- Vázquez-Espinosa, M., González-De-Peredo, A. V., Espada-Bellido, E., Ferreira-González, M., Toledo-Domínguez, J. J., Carrera, C., Palma, M., & Barbero, G. F. (2019). Ultrasound-assisted extraction of two types of antioxidant compounds (TPC and TA) from black chokeberry (*Aronia melanocarpa* L.): Optimization of the individual and simultaneous extraction methods. *Agronomy*, 9(8). <https://doi.org/10.3390/agronomy9080456>
- Wang, Y., Zhao, L., Zhang, R., Yang, X., Sun, Y., Shi, L., & Xue, P. (2020). Optimization of ultrasound-assisted extraction by response surface methodology, antioxidant capacity, and tyrosinase inhibitory activity of anthocyanins from red rice bran. *Food Science and Nutrition*, 8(2), 921–932. <https://doi.org/10.1002/fsn3.1371>
- Xu, D. P., Li, Y., Meng, X., Zhou, T., Zhou, Y., Zheng, J., Zhang, J. J., & Li, H. Bin. (2017). Natural antioxidants in foods and medicinal plants: Extraction, assessment and resources. In *International Journal of Molecular Sciences* (Vol. 18, Issue 1). MDPI AG. <https://doi.org/10.3390/ijms18010096>
- Xue, H., Tan, J., Li, Q., Tang, J., & Cai, X. (2020). Optimization ultrasound-assisted deep eutectic solvent extraction of anthocyanins from raspberry using response surface methodology coupled with genetic algorithm. *Foods*, 9(10). <https://doi.org/10.3390/foods9101409>
- Yang, Z., Fan, G., Gu, Z., Han, Y., & Chen, Z. (2008). Optimization extraction of anthocyanins from purple corn (*Zea mays* L.) cob using tristimulus colorimetry. *European Food Research and Technology*, 227(2), 409–415. <https://doi.org/10.1007/s00217-007-0735-4>
- Yang, Z., & Zhai, W. (2010). Optimization of microwave-assisted extraction of anthocyanins from purple corn (*Zea mays* L.) cob and identification with HPLC-MS. *Innovative Food Science and Emerging Technologies*, 11(3), 470–476. <https://doi.org/10.1016/j.ifset.2010.03.003>
- Zhu, F. (2018). Anthocyanins in cereals: Composition and health effects. In *Food Research International* (Vol. 109, pp. 232–249). Elsevier Ltd. <https://doi.org/10.1016/j.foodres.2018.04.015>