

DAFTAR PUSTAKA

- Abdel-Azeem, A.M., Salem, F.M., Abdel-Azeem, M.A., Nafady, N.A., Mohesien, M.T. and Soliman, E.A. (2016). Biodiversity of the genus aspergillus in different habitats. In: *New and Future Developments in Microbial Biotechnology and Bioengineering: Aspergillus System Properties and Applications*, 3–28. <https://doi.org/10.1016/B978-0-444-63505-1.00001-4>.
- Abdulaal, W.H. (2018). Purification and characterization of α -amylase from *Trichoderma pseudokoningii*. *BMC Biochemistry*, 19(1): 4–9. <https://doi.org/10.1186/s12858-018-0094-8>.
- Adnan, M., Zheng, W., Islam, W., Arif, M., Abubakar, Y.S., Wang, Z. and Lu, G. (2018). Carbon catabolite repression in filamentous fungi, 19(48): 1–23. <https://doi.org/10.3390/ijms19010048>.
- Aggarwal, R., Dutta, T. and Sheikh, J. (2019). Extraction of amylase from the microorganism isolated from textile mill effluent vis a vis desizing of cotton. *Sustainable Chemistry and Pharmacy* 14: 100178. <https://doi.org/10.1016/j.scp.2019.100178>.
- Aiyer, P. (2005). Amylases and their applications. *African Journal of Biotechnology*, 4(13): 1525–1529. <https://doi.org/10.5772/65363>.
- Al-daamy, A.A.H.K. (2021). Production of α – amylase by fungi isolated from oil-contaminated soil in karbala, Iraq, 12(1): 878–883. <https://doi:10.31838/srp.2021.1.123>.
- Aliyah, A., Alamsyah, G., Ramadhani, R. and Hermansyah, H. (2017). Production of α -Amylase and β -Glucosidase from *Aspergillus niger* by solid state fermentation method on biomass waste substrates from rice husk, bagasse and corn cob. *Energy Procedia*, 136: 418–423. <https://doi.org/10.1016/j.egypro.2017.10.269>.
- Altermann, E. and Klaenhammer, T.R. (2005). PathwayVoyager: Pathway mapping using the Kyoto Encyclopedia of Genes and Genomes (KEGG) database. *BMC Genomics*, 6: 1–7. <https://doi.org/10.1186/1471-2164-6-60>.
- Atmaja, D.S. (2013). Isolasi, purifikasi dan karakterisasi α -amilase dari *Trichoderma viride* FNCC 6013. *Chemical Information*, 1(1): 85–93. <https://ejournal3.undip.ac.id/index.php/kimia/article/view/1857/1856>.
- Avwioroko, O.J., Anigboro, A.A., Unachukwu, N.N. and Tonukari, N.J. (2018). Isolation, identification and in silico analysis of alpha-amylase gene of *Aspergillus niger* strain CSA35 obtained from cassava undergoing spoilage.

Biochemistry and Biophysics Reports 14: 35–42.
<https://doi.org/10.1016/j.bbrep.2018.03.006>.

Bagheri, H., Dyer, R., Severin, A. and Rajan, H. (2020). Comprehensive analysis of non-redundant protein database. Research Square, 1–9.
<https://doi.org/10.21203/rs.3.rs-54568/v1>.

Baks, T., Janssen, A. E. M., and Boom, R. M. (2006). The effect of carbohydrates on α -amylase activity measurements. Enzyme and Microbial Technology, 39(1), 114–119. <https://doi.org/10.1016/j.enzmictec.2005.10.005>

Baroncelli, R., Piaggieschi, G., Fiorini, L., Bertolini, E., Zapparata, A., Pè, M.E., Sarrocco, S. and Vannacci, G. (2015). Draft whole-genome sequence of the biocontrol agent *Trichoderma harzianum* T6776. Genome Announcements, 3(3): 9–10. <https://doi.org/10.1128/genomeA.00647-15>.

Boulesteix, M., Weiss, M. and Biémont, C. (2006). Differences in genome size between closely related species: The *Drosophila melanogaster* species subgroup. Molecular Biology and Evolution, 23(1): 162–167.
<https://doi.org/10.1093/molbev/msj012>.

Cha, H.J. and Yoo, Y.J. (1996). Determination of optimal glucose concentrations for glucoamylase production from plasmid-harboring and chromosome-integrated recombinant yeasts using a SUC2 promoter. Process Biochemistry, 31(5): 499–506. [https://doi.org/10.1016/0032-9592\(95\)00093-3](https://doi.org/10.1016/0032-9592(95)00093-3).

Chand, P., Aruna, A., Maqsood, A.M. and Rao, L.V. (2005). Novel mutation method for increased cellulase production. Journal of Applied Microbiology, 98(2): 318–323. <https://doi.org/10.1111/j.1365-2672.2004.02453.x>.

Copetti, M.V. (2019). ScienceDirect Fungi as industrial producers of food ingredients. Current Opinion in Food Science, 25: 52–56.
<https://doi.org/10.1016/j.cofs.2019.02.006>.

Crozet, P., Margalha, L., Confraria, A., Rodrigues, A., Martinho, C., Adamo, M., Elias, C.A. and Baena-González, E. (2014). Mechanisms of regulation of SNF1/AMPK/SnRK1 protein kinases. Frontiers in Plant Science 5: 1–17.
<https://doi.org/10.3389/fpls.2014.00190>.

Cuomo, C.A., and Birren, B.W. (2010). The Fungal Genome Initiative and Lessons Learned from Genome Sequencing. In Guide to Yeast Genetics: Functional Genomics, Proteomics, and Other Systems Analysis (2nd ed., Vol. 470, Issue 10). Elsevier Inc. [https://doi.org/10.1016/S0076-6879\(10\)70034-3](https://doi.org/10.1016/S0076-6879(10)70034-3).

- Davolos, D., Russo, F., Canfora, L., Malus, E., Tartanus, M., Furmanczyk, E. M., Ceci, A., Maggi, O., & Persiani, A. M. (2021). A genomic and transcriptomic study on the DDT-resistant *Trichoderma hamatum* FBL 587: first genetic data into mycoremediation strategies for DDT-Polluted Sites. *Microorganisms*, 9(1680): 1-21. <https://doi.org/10.3390/microorganisms9081680>
- de Almeida, P.Z., Pereira, M.G., de Carvalho, C.C., Heinen, P.R., Ziotti, L.S., Messias, J.M., Jorge, J.A, Polizeli, M. and Maria, L. (2017). Bioprospection and characterization of the amylolytic activity by filamentous fungi from Brazilian atlantic forest. *Biota Neotropica*, 17(3): 1–9. <https://doi.org/10.1590/1676-0611-bn-2017-0337>.
- de Assis, L.J., Silva, L.P., Bayram, O., Dowling, P., Kniemeyer, O., Krüger, T., Brakhage, A.A., Chen, Y., Dong, L., Tan, K., Wong, K.H., Ries, L.N.A. and Goldman, G.H. (2021). Carbon catabolite repression in filamentous fungi is regulated by phosphorylation of the transcription factor creA. *MBio*, 12(1): 1–21. <https://doi.org/10.1128/mBio.03146-20>.
- de Souza, P.M. and Magalhães. P.O. (2010). Application of microbial α -amylase in industry - a review. *Brazilian Journal of Microbiology*, 41(4): 850–861. <https://doi.org/10.1590/s1517-83822010000400004>.
- Dey, T.B., Kumar, A., Banerjee, R., Chandna, P. and Kuhad, R.C. (2016). Improvement of microbial α -amylase stability: Strategic approaches. *Process Biochemistry*, 51(10): 1380–1390. <https://doi.org/10.1016/j.procbio.2016.06.021>.
- Dolfini, D., Zambelli, F., Pavesi, G. and Mantovani, R. (2009). A perspective of promoter architecture from the CCAAT box, *Cell Cycle* 8(24): 4127-4137. <https://doi.org/10.4161/cc.8.24.10240>.
- Dorn, A., Bollekens, J., Staub, A., Mathis, D. and Mol, L.D.G. (1987). A multiplicity of CCAAT Box-Binding Proteins, *Cell* 50: 663–672. [https://doi.org/10.1016/0092-8674\(87\)90513-7](https://doi.org/10.1016/0092-8674(87)90513-7).
- Dung, N. T. P., Rombouts, F. M., & Nout, M. J. R. (2006). Functionality of selected strains of moulds and yeasts from Vietnamese rice wine starters. *Food Microbiology*, 23(4), 331–340. <https://doi.org/10.1016/j.fm.2005.05.002>
- Egbuta, A.M., Mwanza, M. and Babalola, O.O. (2016). A Review of the Ubiquity of Ascomycetes Filamentous Fungi in Relation to Their Economic and Medical Importance. *Advances in Microbiology*, 6(14): 1140–1158. <https://doi.org/10.4236/aim.2016.614103>.

- Elizabeth, A., Widjaja, Rahayuningsih, Y., Rahajoe, J.S., Ubaidillah, R., Maryanto, I., Walujo, E.B. and Semiadi, G. (2014). Kekinian Keanekaragaman Hayati Indonesia. Igarss (Issue 1). <https://doi.org/10.1007/s13398-014-0173-7.2>.
- Fadel, M., Abdel-Halim, S., Sharada, H. and Yehia, A. (2020). Production of glucoamylase, α -amylase and cellulase by aspergillus oryzae F-923 cultivated on wheat bran under solid state fermentation. 23(4): 8–22. <https://doi.org/10.9734/JABB/2020/v23i430149>.
- Far, B.E., Ahmadi, Y., Khosroushahi, A.Y. and Dilmaghani, A. (2020). Microbial alpha-amylase production: progress, challenges, and perspectives. Advanced Pharmaceutical Bulletin, 10(3): 350–358. <https://doi.org/10.34172/apb.2020.043>.
- Farooq, M.A., Ali, S., Hassan, A., Tahir, H.M., Mumtaz, S. and Mumtaz S. (2021). Biosynthesis and industrial applications of α -amylase: a review. Archives of Microbiology, 203(4): 1281–1292. <https://doi.org/10.1007/s00203-020-02128-y>.
- Fast, N.M., Logsdon, J.M.Jr. and Doolittle W F. (1999). Letter to the editor phylogenetic analysis of the tata box binding protein (TBP) gene from *Nosema locustae*: evidence for a microsporidia – fungi relationship and spliceosomal intron loss. Molecular Biology and Evolution, 16(10): 1415–1419. <https://doi.org/10.1093/oxfordjournals.molbev.a026052>.
- Ferreira, E.O., Montijo, A.N., Martins, da.S.E. and Mutton, J.R.M. (2015). Production of -amylase by solid state fermentation by *Rhizopus oryzae*. African Journal of Biotechnology, 14(7): 622–628. <https://doi.org/10.5897/AJB2014.14296>.
- Filho, J.A.F., Horta, M.A.C, Beloti, L.L, dos Santos, C.A. and de Souza, A.P. (2017). Carbohydrate-active enzymes in *Trichoderma harzianum*: A bioinformatic analysis bioprospecting for key enzymes for the biofuels industry. BMC Genomics, 18(1): 1–12. <https://doi.org/10.1186/s12864-017-4181-9>.
- Fifendy, M, Periadnadi, Indriati, G., Osnita, S. and Annisa, L. (2020). Isolation and activity of amylase enzyme in isolates of fungi from black rice lemang (*Oryza sativa* siarang). Proceedings of the International Conference on Biology, Sciences and Education 10: 46–50. <https://doi.org/10.2991/absr.k.200807.011>.
- Forment, J.V., Flipphi, M., Ventura, L., González, R., Ramón, D. and MacCabe, A.P. (2014). High-affinity glucose transport in *Aspergillus nidulans* is mediated by the products of two related but differentially expressed genes. Plos one, 9(4): e94662 <https://doi.org/10.1371/journal.pone.0094662>.

- Gancedo, J.M. (1998). Yeast Carbon Catabolite Repression. *Microbiology and Molecular Biology Reviews*, 62(2): 334–361. <https://doi.org/10.1128/membr.62.2.334-361.1998>.
- Ghadi, A., Mahjoub, S. and Mehravar, R. (2011). Management of glucose production process from rice husk by solid state fermentation method. *International Conference on Biotechnology and Environment Management*, 18: 93–97.
- Ghazala, I., Haddar, A., Romdhane, M.B. and Ellouz-Chaanouni, S. (2016). Screening and molecular identification of new microbial strains for production of enzymes of biotechnological interest. *Brazilian Archives of Biology and Technology*, 59: 1–12. <https://doi.org/10.1590/1678-4324-2016150152>.
- Godbey, W.T. (2014). An Introduction to Biotechnology The Science, Technology and Medical Applications Chapter 4. In *Introduction to Biotechnology “genes.”* Elsevier Ltd. <https://doi.org/10.1016/B978-1-907568-28-2.00004-6>.
- Gomi, K. (2019). Regulatory mechanisms for amylolytic gene expression in the koji mold *Aspergillus oryzae*. *Bioscience, Biotechnology and Biochemistry*, 83(8): 1385–1401. <https://doi.org/10.1080/09168451.2019.1625265>.
- Gomi, K., Akeno, T., Minetoki, T., Ozeki, K., Kumagai, C., Okazaki, N. and Imura, Y. (2000). Molecular cloning and characterization of a transcriptional activator gene, amyR, involved in the amylolytic gene exoression in *Aspergillus oryzae*. *Bioscience, Biotechnology and Biochemistry*, 64(4): 816–827.
- Gopinath, S.C.B., Anbu, P., Arshad, M.K.M., LakshmiPriya, T., Voon, C.H., Hashim, U. and Chinni, S.V. (2017). Biotechnological processes in microbial amylase production. *BioMed Research International*, Feb 9: 1-9. <https://doi.org/10.1155/2017/1272193>.
- Guo, L., Han, L., Yang, L., Zeng, H., Fan, D., Zhu, Y. and Feng, Y. (2014). Genome and transcriptome analysis of the fungal pathogen fusarium *Oxysporum* f. Sp . *Cubense* causing banana vascular wilt disease. *Plos one*, 10(1): e0117621. <https://doi.org/10.1371/journal.pone.0095543>.
- Gupta, A., Gupta, V.K., Modi, D.R. and Yadava, L.P. (2008). Production and characterization of α -amylase from *Aspergillus niger*. *Biotechnology*, 7(3): 551–556.
- Haq, I., Abdullah, R., Ashraf, H. and Shah, A.H. (2002). Isolation and screening of fungi for the biosynthesis of alpha amylase. *Biotechnology*, 1(2): 61–66. <https://doi.org/10.3923/biotech.2002.61.66>.

- Hasegawa, K., Kubota, M. and Matsuura, Y. (1999). Roles of catalytic residues in α -amylases as evidenced by the structures of the product-complexed mutants of a maltotetraose-forming amylase. *Protein Engineering*, 12(10): 819–824. <https://doi.org/10.1093/protein/12.10.819>.
- Haseltine, C., Rolfsmeier, M. and Blum, P. (1996). The glucose effect and regulation of α -amylase synthesis in the hyperthermophilic archaeon *Sulfolobus solfataricus*. *Journal of Bacteriology*, 178(4): 945–950. <https://doi.org/10.1128/jb.178.4.945-950.1996>.
- Hidayati F.L.N., Suroto D.A., Sardjono, and Cahyanto M.N. (2022). Effect of glucose on endoglucanase and β -glucosidase production by some indigenous Indonesian fungi. *International Journal of Agricultural Technology* 18(4): 1505-1520. <https://doi.org/10.22207/JPAM.16.1.12>.
- Hiramoto, T., Tanaka, M., Ichikawa, T., Matsuura, Y., Hasegawa-Shiro, S., Shintani, T. and Gomi K. (2015). Endocytosis of a maltose permease is induced when amylolytic enzyme production is repressed in *Aspergillus oryzae*. *Fungal Genetics and Biology*, 82: 136–144. <https://doi.org/10.1016/j.fgb.2015.05.015>.
- Hortschansky, P., Haas, H., Huber, E.M., Groll, M. and Brakhage, A.A. (2017). The CCAAT-binding complex (CBC) in *Aspergillus* species. *Biochimica et Biophysica Acta- Gene Regulatory Mechanisms*, 1860(5): 560–570. <https://doi.org/10.1016/j.bbagr.2016.11.008>.
- Huang, M., Bai, Y., Sjostrom, S. L., Hallström, B. M., Liu, Z., Petranovic, D., Uhlén, M., Joensson, H. N., Andersson-Svahn, H., & Nielsen, J. (2015). Microfluidic screening and whole-genome sequencing identifies mutations associated with improved protein secretion by yeast. *Proceedings of the National Academy of Sciences of the United States of America*, 112(34), E4689–E4696. <https://doi.org/10.1073/pnas.1506460112>
- Hunter, A.J., Morris, T.A., Jin, B., Saint, C.P. and Kelly, J.M. (2013). Deletion of CreB in *Aspergillus oryzae* increases secreted hydrolytic enzyme activity. *Applied and Environmental Microbiology*, 79(18): 5480–5487. <https://doi.org/10.1128/AEM.01406-13>.
- Ichinose, S., Tanaka, M., Shintani, T. and Gomi, K. (2014). Improved α -amylase production by *Aspergillus oryzae* after a double deletion of genes involved in carbon catabolite repression. *Applied Microbiology and Biotechnology*, 98(1): 335–343. <https://doi.org/10.1007/s00253-013-5353-4>.
- Ichinose, S., Tanaka, M., Shintani, T. and Gomi, K. (2018). Increased production of biomass-degrading enzymes by double deletion of creA and creB genes involved in carbon catabolite repression in *Aspergillus oryzae*. *Journal of*

- Bioscience and Bioengineering, 125(2): 141–147.
<https://doi.org/10.1016/j.jbiosc.2017.08.019>.
- Imran, M., Asad, M.J. and Hadri, S.H. (2012). Production and industrial applications of laccase enzyme. *Journal of Cell and Molecular Biology*, 10(1): 1-11.
- Ivanova, C., Ramoni, J., Aouam, T., Frischmann, A., Seiboth, B., Baker, S.E., Crom, S.L., Lemoine, S., Margeot, A., and Bidard, F. (2017). Genome sequencing and transcriptome analysis of *Trichoderma reesei* QM9978 strain reveals a distal chromosome translocation to be responsible for loss of *vib1* expression and loss of cellulase induction. *Biotechnology for Biofuels*, 10(209): 1–15. <https://doi.org/10.1186/s13068-017-0897-7>
- Jablonski, D., Kaustuv, R. and Valentine, J.W. (2006). Out of the Tropics: Evolutionary Diversity Gradient. *Science*, 314: 102–106. <https://doi.org/10.1126/science.1130880>.
- Ju, F., Zhu, J., Shao, B., Kong, L., Liu, T.Y., Zheng, W.M., and Bu, D. (2021). CopulaNet: Learning residue co-evolution directly from multiple sequence alignment for protein structure prediction. *Nature Communications* 12(1): 1-9. <https://doi.org/10.1038/s41467-021-22869-8>
- Kanti, A. (2017). Potensi dari kapang aspergillus niger, rhizophus oryzae dan neurospora sitophila sebagai penghasil ezim fitase dan amilase pada substrate ampas tahu. *Buletin Peternakan*, 41(1): 26. <https://doi.org/10.21059/buletinpeternak.v41i1.13337>.
- Karim, K.M.R., Husaini, A., Sing, N.N., Sinang, F.M., Roslan, H.A. and Hussain, H. (2018). Purification of an alpha amylase from *Aspergillus flavus* NSH9 and molecular characterization of its nucleotide gene sequence. *Biotech*, 8(4): 1–14. <https://doi.org/10.1007/s13205-018-1225-z>.
- Kato, M. (2005). An overview of the CCAAT-box binding factor in filamentous fungi: assembly, nuclear translocation, and transcriptional enhancement. *Bioscience, biotechnology, and biochemistry*, 69(4): 663–672. <https://doi.org/10.1271/bbb.69.663>.
- Kato, M., Sekine, K. and Tsukagoshi, N. (1996). Sequence-specific binding sites in the Taka- amylase A G2 promoter for the CreA repressor mediating carbon catabolite repression. *Bioscience, Biotechnology, and Biochemistry*, 60(11): 1776-1779. <https://doi.org/10.1271/bbb.60.1776>
- Kelly, R.M., Dijkhuizen, L. and Leemhuis, H. (2009). Starch and a-glucan acting enzyme, modulating their properties by directed evolution. *Journal of Biotechnology*, 140: 184–193. <https://doi.org/10.1016/j.jbiotec.2009.01.020>.

- Khokhar, I., Mukhtar, I. and Mushtaq, S. (2011). Isolation and Screening of Amyolytic Filamentous Fungi. *Journal of Applied Sciences and Environmental Management*, 15(1): 203-206. <https://doi.org/10.4314/jasem.v15i1.68442>.
- Khosravi, C. (2017). Sugar catabolism during growth on plant biomass in *Aspergillus*. [PhD thesis, Utrecht University].
- Kobayashi, A., Sano, M., Oda, K., Hisada, H., Hata, Y. and Ohashi, S. (2007). The glucoamylase-encoding gene (*gluB*) is expressed in solid-state culture with a low water content. *Bioscience, Biotechnology and Biochemistry* 71(7): 1797–1799. <https://doi.org/10.1271/bbb.70132>
- Koboldt, D.C., Steinberg, K.M., Larson, D.E., Wilson, R.K. and Mardis, E.R. (2013). Review the next-generation sequencing revolution and its impact on genomics. *Cell*, 15(2013): 27–38. <https://doi.org/10.1016/j.cell.2013.09.006>
- Krishnan, A., Convey, P., Gonzalez-Rocha, G. and Alias, S.A. (2016). Production of extracellular hydrolase enzymes by fungi from King George Island. *Polar Biology*, 39: 65–76. <https://doi.org/10.1007/s00300-014-1606-7>.
- Krisnawati, R., Sardjono, Widada, J., Suroto, D.A. and Cahyanto, M.N. (2022). Effect of Glucose on Endo-xylanase and β -xylosidase Production by Fungi Isolated in Indonesia. *Journal of Pure and Applied Microbiology*, 16(1): 226–234. <https://doi.org/10.22207/JPAM.16.1.12>.
- Kubicek, C.P., Steindorff, A.S., Chenthamara, K., Manganiello, G., Henrissat, B., Zhang, J., Cai, F, Kopchinskiy, A.G, Kubicek, E.M, Kuo, A., Baroncelli, R., Sarrocco, S., Noronha, E.F, Vannacci, G. and Shen, Q. (2019). Evolution and comparative genomics of the most common *Trichoderma* species. *BMC Genomics*, 20:1–24. <https://doi.org/10.1186/s12864-019-5680-7>.
- Kumar, B.L. and Gopal, D.V.R.S. (2015). Effective role of indigenous microorganisms for sustainable environment. *3 Biotech* 5(6): 867–876. <https://doi.org/10.1007/s13205-015-0293-6>
- Kumar, P. and Satyanarayana, T. (2009). Microbial glucoamylases: characteristics and applications. *Critical Reviews in Biotechnology*, 29(3): 225–255. <https://doi.org/10.1080/07388550903136076>.
- Kunamneni, A., Permaul, K. and Singh, S. (2005). Amylase production in solid state fermentation by the thermophilic fungus *Thermomyces lanuginosus*. *Journal of Bioscience and Bioengineering*, 100(2): 168–171. <https://doi.org/10.1263/jbb.100.168>.

- Kusnadi, Saefudin, and Astri, E. (2009). Keanekaragaman jamur selulolitik dan amilolitik pengurai sampah organik dari berbagai substrat. *Jurnal Pendidikan Biologi*, 229: 1–10.
- Lee, J. and Paetzel, M. (2011). Structure of the catalytic domain of glucoamylase from *Aspergillus niger*. *Acta Crystallographica Section F: Structural Biology and Crystallization Communications*, 67(2): 188–192. <https://doi.org/10.1107/S1744309110049390>.
- Li, C., Shu, W., Wang, S., Liu, P., Zhuang, Y., Zhang, S. and Xia J. (2018). Dynamic metabolic response of *Aspergillus niger* to glucose perturbation: evidence of regulatory mechanism for reduced glucoamylase production. *Journal of Biotechnology*, 287: 28–40. <https://doi.org/10.1016/j.jbiotec.2018.08.005>.
- Li, L., Stoekert, C.J.J. and Roos, D.S. (2003). OrthoMCL: Identification of Ortholog Groups for Eukaryotic Genomes. *Genome Research*, 13(9): 2178–2189. <https://doi.org/10.1101/gr.1224503.candidates>.
- Liao, G.Y., Zhao, S., Zhang, T., Li, C.X., Liao, L.S., Zhang, F.F., Luo, X.M. and Feng, J.X. (2018). The transcription factor TpRfx1 is an essential regulator of amylase and cellulase gene expression in *Talaromyces pinophilus*. *Biotechnology for Biofuels*, 11(1): 1–17. <https://doi.org/10.1186/s13068-018-1276-8>.
- Londoño-Hernández, L., Ramírez-Toro, C., Ruiz, H.A., Ascacio-Valdés, J.A., Aguilar-Gonzalez, M.A., Rodríguez-Herrera, R. and Aguilar, C.N. (2017). *Rhizopus oryzae* – Ancient microbial resource with importance in modern food industry. *International Journal of Food Microbiology*, 257: 110–127. <https://doi.org/10.1016/j.ijfoodmicro.2017.06.012>.
- Lubertozzi, D. and Keasling, J.D. (2009). Developing *Aspergillus* as a host for heterologous expression. *Biotechnology Advances*, 27(1): 53–75. <https://doi.org/10.1016/j.biotechadv.2008.09.001>.
- Lv, C., Wang, P., Ma, L., Zheng, M., Liu, Y. and Xing F. (2018). Large-scale comparative analysis of eugenol-induced / repressed genes expression in *Aspergillus flavus* using RNA-seq. *Frontiers in Microbiology*, 9: 1116. <https://doi.org/10.3389/fmicb.2018.01116>.
- MacPherson, S., Larochelle, M. and Turcotte, B. (2006). A fungal family of transcriptional regulators: the zinc cluster proteins. *Microbiology and Molecular Biology Reviews*, 70(3): 583–604. <https://doi.org/10.1128/mubr.00015-06>
- Mahmood, S. and Rahman, S.R. (1970). Production and partial characterization of extracellular & alpha-amylase by *Trichoderma viride*. *Bangladesh Journal of Microbiology*, 25(2): 99–103. <https://doi.org/10.3329/bjm.v25i2.4870>.

- Mandels, M., Hontz, L. and Nystrom, J. (1974). Enzymatic hydrolysis of waste cellulose. *Biotechnology and Bioengineering*, 16(11): 1471–1493. <https://doi.org/10.1002/bit.260161105>.
- Marín-Navarro, J. and Polaina, J. (2011). Glucoamylases: structural and biotechnological aspects. *Applied microbiology and biotechnology*, 89: 1267–1273. <https://doi.org/10.1007/s00253-010-3034-0>.
- McCleary, B. V., Bouhet, F., and Driguez, H. (1991). Measurement of amyloglucosidase using P-nitrophenyl β -maltoside as substrate. *Biotechnology Techniques*, 5(4), 255–258. <https://doi.org/10.1007/BF02438658>
- Megazyme International Ireland. (2014). Assay of amyloglucosidase using p-nitrophenyl plus thermostable β -glucosidase. www.megazyme.com diakses pada 20 Agustus 2018
- Megazyme International Ireland. (2012). Alpha-amylase, assay procedure: For the measurement of cereal and microbial alpha-amylase. <https://secure.megazyme.com> diakses pada 20 Agustus 2018
- Mertens, J.A. and Skory, C.D. (2007). Isolation and characterization of a second glucoamylase gene without a starch binding domain from *Rhizopus oryzae*. *Enzyme and Microbial Technology*, 40(4): 874–880. <https://doi.org/10.1016/j.enzmictec.2006.07.003>.
- Mohamed, S.A.H, Azhar, E.I., Ba-Akdah, M.M., Tashkandy, N. and Kumosani T. (2011). Production, purification and characterization of α -amylase from *Trichoderma harzianum* grown on mandarin peel. *African Journal of Microbiology Research*, 5(9): 1018-1028. <https://doi.org/10.5897/ajmr10.890>.
- Mohanta, T.K. and Bae, H. (2015). The diversity of fungal genome. *Biological Procedures Online*, 17(1): 1–9. <https://doi.org/10.1186/s12575-015-0020-z>.
- Mojsov, K. (2014). Microbial alpha-amylases and their industrial applications: A review. *International Journal of Management, IT, and Engineering*, 2(10): 583–609.
- Mojzita, D., Rantasalo, A. and Jantti, J. (2019). Gene expression engineering in fungi. *Current Opinion in Biotechnology* 59: 141–149. <https://doi.org/10.1016/j.copbio.2019.04.007>
- Morozova, O., Hirst, M. and Marra, M.A. (2009). Applications of new sequencing technologies for transcriptome analysis. *Annual Review of Genomics and Human Genetics*, 10:135-151. <https://doi.org/10.1146/annurev-genom-082908-145957>.

- Mukherjee, P.K., Horwitz, B.A., Herrera-estrella, A., Schmoll, M. and Kenerley, C.M. (2013). Trichoderma research in the genome era. Annual Review of Phytopathology, 5: 105–29. <https://doi.org/10.1146/annurev-phyto-082712-102353>.
- Murakoshi, Y., Makita, T. and Kato, M. (2012). Comparison and characterization of α -amylase inducers in *Aspergillus nidulans* based on nuclear localization of AmyR. Applied Microbiology and Biotechnology, 94: 1629–1635. <https://doi.org/10.1007/s00253-012-3874-x>.
- Mutz, K., Heilkenbrinker, A., Lo, M. and Stahl, F. (2013). Transcriptome analysis using next-generation sequencing. Current Opinion in Biotechnology, 24(1): 22–30. <https://doi.org/10.1016/j.copbio.2012.09.004>.
- Nagalakshmi, U., Waern, K. and Snyder, M. (2010). RNA-seq: a method for comprehensive transcriptome analysis. Current Protocols in Molecular Biology, January: 1–13. <https://doi.org/10.1002/0471142727.mb0411s89>.
- Naili, B., Sahnoun, M., Bejar, S. and Kammoun R. (2016). Optimization of submerged *Aspergillus oryzae* S2 α -amylase production. Food Science and Biotechnology, 25(1): 185–192. <https://doi.org/10.1007/s10068-016-0028-4>.
- Nangin, D. and Sutrisno, A. (2015). Potensi kapang *Aspergillus niger*, *Rhizopus oryzae* dan *Neurospora sitophila* sebagai penghasil enzim fitase dan amilase pada substrat ampas tahu. Jurnal Pangan Dan Agroindustri, 3(3): 1032–1039.
- Napitupulu, T.P., Silaban, N.R., Kanti, A. and Sudiana, I.M. (2019). The effect of substrate composition on the activity of amylase and cellulase by *Trichoderma harzianum* strains under solid state fermentation. Journal of Microbial Systematics and Biotechnology, 1(2): 41–48. <https://doi.org/10.37604/jmsb.v1i2.26>.
- Nardone, V., Chaves-sanjuan, A. and Nardini, M. (2017). Structural determinants for NF-Y / DNA interaction at the CCAAT box. Biochimica et Biophysica Acta - Gene Regulatory Mechanisms, 1860(5): 571–580. <https://doi.org/10.1016/j.bbagr.2016.09.006>.
- Ng, P.C. and Kirkness, E.F. (2010). Whole Genome Sequencing. In: Barnes, M., Breen, G. (eds) Genetic Variation. Methods in Molecular Biology, vol 628. Humana Press, Totowa, NJ. <https://doi.org/10.1007/978-1-60327-367-1>.
- Nitta, M., Furukawa, T., Shida, Y., Mori, K., Kuhara, S. and Morikawa, Y. (2012). A new Zn (II) 2 Cys 6 -type transcription factor BglR regulates β -glucosidase expression in *Trichoderma reesei*. Fungal Genetics and Biology, 49(5): 388–397. <https://doi.org/10.1016/j.fgb.2012.02.009>.

- Norouzian, D., Norouzian, D., Akbarzadeh, A., Akbarzadeh, A., Scharer, J.M., Scharer, J.M. and Moo, Y.M. (2006). Fungal glucoamylases. *Biotechnology Advances*, 24 (1): 80–85. <https://doi.org/10.1016/j.biotechadv.2005.06.003>.
- Obafemi, Y.D., Ajayi, A.A., Olasehinde, G.I., Atolagbe, O.M. and Onibokun, E.A. (2017). Screening and partial purification of amylase from *Aspergillus Niger* isolated from deteriorated tomato (*Lycopersicon Esculentum* mill.) fruits. *African Journal of Clinical and Experimental Microbiology*, 19(1): 47-57. <https://doi.org/10.4314/ajcem.v19i1.7>.
- Ogbonna, C.N., Okpokwu, N.M., Okafor, C.U. and Onyia, C.E. (2014). Isolation and screening of amylase producing fungi obtained from garri processing site. *International Journal of Biotechnology and Food Science*, 2(5): 88–93.
- Oktavia, Y., Lestari, S.D., Lestari, S., Herpandi and Jannah, M. (2018). Optimasi waktu inkubasi produksi protease dan amilase isolat bakteri asal terasi ikan teri *Stolephorus* sp. *Jurnal Ilmu dan Teknologi Kelautan Tropis* 10(3): 719–725. <https://doi.org/10.29244/jitkt.v10i3.18840>
- Pachauri, S., Sherkhane, P.D., Kumar, V. and Mukherjee, P.K. (2020). Whole genome sequencing reveals major deletions in the genome of m7, a gamma ray-induced mutant of *Trichoderma virens* that is repressed in conidiation, secondary metabolism, and mycoparasitism. *Frontiers in Microbiology*, 11: 1–12. <https://doi.org/10.3389/fmicb.2020.01030>.
- Parra-marín, O., López-pacheco, K., Hernández, R. and López-villaseñor, I. (2020). Molecular & biochemical parasitology the highly diverse TATA box-binding proteins among protists: a review. *Molecular and Biochemical Parasitology* 239: 111312. <https://doi.org/10.1016/j.molbiopara.2020.111312>.
- Pasin, T.M., Benassi, V.M., Heinen, P.R., Damasio, A.R. de, L., Cereia, M., Jorge, J.A. and Polizeli, M. (2017). Purification and functional properties of a novel glucoamylase activated by manganese and lead produced by *Aspergillus japonicus*. *International Journal of Biological Macromolecules*, 102: 779–788. <https://doi.org/10.1016/j.ijbiomac.2017.04.016>.
- Paul, D. and Bengal, W. (2016). Microorganisms and a-amylase: A Concise Review. *Innovare Journal of Science*, 4(4): 1–5.
- Pavezzi, F. C., Gomes, E and Da Silva, R. (2008). Production and characterization of glucoamylase from fungus *Aspergillus awamori* expressed in yeast *Saccharomyces cerevisiae* using different carbon sources. *Brazilian Journal of Microbiology*, 39(1), 108–114. <https://doi.org/10.1590/S1517-83822008000100024>

- Pel, H.J., Winde, J.H.De, Archer, D.B., Dyer, P.S., Hofmann, G., Schaap, P.J., Turner, G., Vries, R.P. De, Albang, R., Albermann, K., Andersen, M.R., Bendtsen, J.D., Benen, J.A.E., Berg, M., Van Den, Breestraat, S., Caddick, M.X., Contreras, R., Cornell, M., Coutinho, P.M. and Pal K. (2007). Genome sequencing and analysis of the versatile cell. *Nature Biotechnology*, 25(2): 221–231. <https://doi.org/10.1038/nbt1282>
- Petersen, K.L., Lehmbeck, J. and Christensen, T. (1999). A new transcriptional activator for amylase genes in *Aspergillus*. *Molecular and General Genetics*, 262(4–5): 668–676. <https://doi.org/10.1007/s004380051129>.
- Pitt, J.I. and Hocking, A.D. (2009). *Fungi and Food Spoilage* (Third, Issue April). Springer.
- Portnoy, T., Margeot, A., Linke, R., Atanasova, L., Fekete, E., Sándor, E., Hartl, L., Karaffa, L., Druzhinina, I.S., Seiboth, B., Crom, S.L. and Kubicek, C.P. (2011). The CRE1 carbon catabolite repressor of the fungus *Trichoderma reesei*: a master regulator of carbon assimilation. *BMC Genomics*, 12(1): 1–12. <https://doi.org/10.1186/1471-2164-12-269>.
- Rajoka, M.I. and Yasmeen, A. (2005). Induction, and production studies of a novel glucoamylase of *Aspergillus niger*. *World Journal of Microbiology and Biotechnology*, 21(2): 179–187. <https://doi.org/10.1007/s11274-004-1766-7>.
- Reilly, M.C., Magnuson, J.K. and Baker, S.E. (2016). Approaches to understanding protein hypersecretion in fungi. *Fungal Biology Reviews*, 30(4): 145–151. <https://doi.org/10.1016/j.fbr.2016.06.002>.
- Rezaei-Matehkolaei, A., Mirhendi, H., Makimura, K., Sybren De Hoog, G., Satoh, K., Najafzadeh, M.J. and Shidfar, M.R. (2014). Nucleotide sequence analysis of beta tubulin gene in a wide range of dermatophytes. *Medical Mycology*, 52(7): 674–688. <https://doi.org/10.1093/mmy/myu033>.
- Ronne, H. (1995). Glucose repression in fungi. *Trends in Genetics*, 11(1): 12–17. [https://doi.org/10.1016/S0168-9525\(00\)88980-5](https://doi.org/10.1016/S0168-9525(00)88980-5).
- Sadhasivam, B., Thanaraj, P., Veerichetty, V., Nachimuthu, S., Ponnusamy, R., Peraman, M., Jabamalai, H. and Ramachandran, S.D. (2018). Production and application of α -amylase from indigenous fungal strain *Aspergillus luchuensis* bs1. *Malaysian Journal of Microbiology* 14(4): 215–228. <https://doi.org/10.21161/mjm.107117>.
- Sahnoun, M., Kriaa, M., Elgharbi, F., Ayadi, D.Z., Bejar, S. and Kammoun, R. (2015). *Aspergillus oryzae* S2 alpha-amylase production under solid state fermentation: Optimization of culture conditions. *International Journal of*

Biological Macromolecules, 75: 73–80.
<https://doi.org/10.1016/j.ijbiomac.2015.01.026>.

- Saini, R., Singh Saini, H., Dahiya, A. and Harnek, S. S. C. (2017). Amylases: Characteristics and industrial applications. *Journal of Pharmacognosy and Phytochemistry*, 6(4): 1865–1871.
- Saleem, A. and Ebrahim, M.K.H. (2014). Production of amylase by fungi isolated from legume seeds collected in Almadinah Almunawwarah, Saudi Arabia. *Journal of Taibah University for Science*, 8(2): 90–97. <https://doi.org/10.1016/j.jtusci.2013.09.002>.
- Samanta, S., Jana, S., Kar, S., Mohapatra, P.K.D., Pati, B.R. and Mondal, K.C. (2017). Induction and catabolite repression of alpha-amylase synthesis in *Bacillus licheniformis* SKB4. *Journal of Microbiology and Biotechnology Research*, 7(6): 8. <https://doi.org/10.24896/jmbr.2017762>.
- Santos, E.D.O. and Martins, M.L.L. (2003). Effect of the medium composition on formation of amylase by *Bacillus* sp. *Brazilian Archives of Biology and Technology*, 46(1): 129–134. <https://doi.org/10.1590/S1516-89132003000100018>.
- Sanz, P., Alms, G.R., Haystead, T.A.J. and Carlson, M. (2000). Regulatory interactions between the reg1-glc7 protein phosphatase and the snf1 protein kinase. *Molecular and Cellular Biology*, 20(4): 1321–1328. <https://doi.org/10.1128/mcb.20.4.1321-1328.2000>.
- Saranraj, P. and Stella, D. (2013). Fungal Amylase - A Review. *International Journal of Microbiological Research*, 4(2): 203–211. <https://doi.org/10.5829/idosi.ijmr.2013.4.2.75170>.
- Sharma, A. and Satyanarayana, T. (2013). Microbial acid-stable α -amylases: Characteristics, genetic engineering and applications. *Process Biochemistry*, 48(2): 201–211. <https://doi.org/10.1016/j.procbio.2012.12.018>.
- Shi, L. L., Mortimer, P.E., Ferry, S.J.W, Zou, X.M., Xu, J., Feng, W.T. and Qiao, L. (2014). Variation in forest soil fungal diversity along a latitudinal gradient. *Fungal Diversity*, 64(1): 305–315. <https://doi.org/10.1007/s13225-013-0270-5>.
- Shroff, R.A., Lockington, R.A. and Kelly, J.M. (1996). Analysis of mutations in the creA gene involved in carbon catabolite depression in *Aspergillus nidulans*. *Canadian Journal of Microbiology*, 42(9): 950–959. <https://doi.org/10.1139/m96-122>.

- Sibthorp, C., Wu, H., Cowley, G., Wong, P.W.H., Palaima, P., Morozov, I.Y., Weedall, G.D. and Caddick, M.X. (2013). Transcriptome analysis of the filamentous fungus *Aspergillus nidulans* directed to the global identification of promoters. *BMC Genomics*, 14(1): 1-40. <https://doi.org/10.1186/1471-2164-14-847>.
- Singh, K.D., Schmalisch, M.H., Stülke, J. and Görke, B. (2008). Carbon catabolite repression in *Bacillus subtilis*: Quantitative analysis of repression exerted by different carbon sources. *Journal of Bacteriology*, 190(21): 7275–7284. <https://doi.org/10.1128/JB.00848-08>.
- Singh, S., Singh, S., Bali, V., Sharma, L. and Mangla, J. (2014). Production of fungal amylases using cheap, readily available agriresidues, for potential application in textile industry. *BioMed Research International*, January:215748. <https://doi.org/10.1155/2014/215748>.
- Sivaramakrishnan, S., Gangadharan, D., Nampoothiri, K.M., Soccol, C.R. and Pandey, A. (2007). Alpha amylase production by *Aspergillus oryzae* employing solid-state fermentation. *Journal of Scientific and Industrial Research*, 66(8): 621–626.
- Slivinski, C.T., Machado, A.V.L., Iulek, J., Ayub, R.A. and de Almeida, M.M. (2011). Biochemical characterisation of a glucoamylase from *Aspergillus niger* produced by solid-state fermentation. *Brazilian Archives of Biology and Technology*, 54(3): 559–568. <https://doi.org/10.1590/S1516-89132011000300018>.
- Soni, S.K., Kaur, A. and Gupta, J.K. (2003). A solid state fermentation based bacterial α -amylase and fungal glucoamylase system and its suitability for the hydrolysis of wheat starch. *Process Biochemistry*, 39(2): 185–192. [https://doi.org/10.1016/S0032-9592\(03\)00058-X](https://doi.org/10.1016/S0032-9592(03)00058-X).
- Srivastava, M. and Shahid, M. (2014). Trichoderma genome to genomics: a review. *Journal of Data Mining in Genomics & Proteomics*, 5(3): 3–6. <https://doi.org/10.4172/2153-0602.1000162>.
- Steidl, S., Tuncher, A., Goda, H., Guder, C., Papadopoulou, N., Kobayashi, T., Tsukagoshi, N., Kato, M. and Brakhage, A.A. (2004). A single subunit of a heterotrimeric CCAAT-binding complex carries a nuclear localization signal: piggy back transport of the pre-assembled complex to the nucleus. *Journal of Molecular Biology*, 342(2): 515–524. <https://doi.org/10.1016/j.jmb.2004.07.011>.
- Sterflinger, K., Tesei, D. and Zakharova, K. (2012). Fungi in hot and cold deserts with particular reference to microcolonial fungi. *Fungal Ecology*, 5(4): 453–462. <https://doi.org/10.1016/j.funeco.2011.12.007>.

- Suleimenova, Zh., Rakhmetova, Zh. and Zhakipbekova, A. (2017). Screening of *Aspergillus oryzae* M/4 for extra cellular alphaamylase production. *International Journal of Engineering Research and Applications*, 07(06), 55–57. <https://doi.org/10.9790/9622-0706015557>
- Sun, T., Zhang, Y., Jiang, H., Yang, K., Wang, S., Wang, R., Li, S., Lei, P., Xu, H. and Qiu Y. (2022). Whole genome sequencing and annotation of *Naematelia aurantialba* (Basidiomycota, edible-medicinal fungi). *Journal of Fungi*, 8(1): 6–14. <https://doi.org/10.3390/jof8010006>.
- Sundarram, A. and Murthy, T.P.K. (2014). α -Amylase Production and Applications: A Review. *Journal of Applied & Environmental Microbiology*, 2(4): 166–175. <https://doi.org/10.12691/jaem-2-4-10>.
- Suzuki, K., Tanaka, M., Konno, Y., Ichikawa, T., Ichinose, S., Hasegawa-Shiro, S., Shintani, T. and Gomi, K. (2015). Distinct mechanism of activation of two transcription factors, AmyR and MalR, involved in amylolytic enzyme production in *Aspergillus oryzae*. *Applied Microbiology and Biotechnology*, 99(4): 1805–1815. <https://doi.org/10.1007/s00253-014-6264-6268>.
- Tanaka, M., Hiramoto, T., Tada, H., Shintani, T. and Gomi, K. (2017). Improved α -amylase production by dephosphorylation mutation of CreD, an arrestin-like protein required for glucose-induced endocytosis of maltose permease and carbon catabolite derepression in *Aspergillus oryzae*. *Applied and Environmental Microbiology*, 83(13): 1–15. <https://doi.org/10.1128/AEM.00592-17>.
- Taniguchi, H. and Honnda, Y. (2009). Amylases. *Applied Microbiology: Industrial*, 163–180. <https://doi.org/10.1016/B978-0-12-805419-2.00008-3>.
- Tarawneh, K.A., Khleifat, K.M., Al-Mustafa, A., Alioui, N. and Wedyan, M.A. (2008). Temporal expression of *Neurospora crassa* tyrosinase gene under the control of glucose -repressible gene-1(*grg-1*) promoter. *Australian Journal of Basic and Applied Sciences*, 2(4): 805–814.
- Tatusov, R.L., Fedorova, N.D., Jackson, J.D., Jacobs, A.R., Kiryutin, B., Koonin, E.V., Krylov, D.M., Mazumder, R., Smirnov, S., Nikolskaya, A.N., Rao, B.S., Mekhedov, S.L., Sverlov, A.V., Vasudevan, S., Wolf, Y.I., Yin, J.J. and Natale, D.A. (2003). The COG database: An updated vesion includes eukaryotes. *BMC Bioinformatics*, 4: 1–14. <https://doi.org/10.1186/1471-2105-4-41>.

- Terrapon, N., Lombard, V., Drula, E. and Coutinho, P.M. (2017). A practical guide to using glycomics databases. Springer: Tokyo. <https://doi.org/10.1007/978-4-431-56454-6>
- Thalmann, M., Coiro, M., Meier, T., Wicker, T., Zeeman, S. C. and Santelia, D. (2019). The evolution of functional complexity within the α -amylase gene family in land plants. *BMC Evolutionary Biology* 19(1): 1–18. <https://doi.org/10.1186/s12862-019-1395-2>
- Thammahong, A., Puttikamonkul, S., Perfect, J.R., Brennan, R.G. and Cramer, R.A. (2017). Central role of the trehalose biosynthesis pathway in the pathogenesis of human fungal infections: opportunities and challenges for therapeutic development. *Microbiology and Molecular Biology Reviews*, 81(2): e00053-16. <https://doi.org/10.1128/membr.00053-16>
- Tiwari, S., Srivastava, R., Singh, C., Shukla, K., Singh, P., Singh, R., Singh, N. and Sharma, R. (2015). Amylases: an overview with special reference to alpha amylase. *Journal of Global Biosciences*, 4(1): 1886–1901.
- Todd, R.B., Lockington, R.A. and Kelly, J.M. (2000). The *Aspergillus nidulans* CreC gene involved in carbon catabolite repression encodes a WD40 repeat protein. *Molecular and General Genetics*, 263(4): 561–570. <https://doi.org/10.1007/s004380051202>.
- Tolic-nørrelykke, S.F., Rasmussen, M.B., Pavone, F.S., Berg-sørensen, K. and Oddershede, L.B. (2006). Stepwise bending of DNA by a single tata-box binding protein. *Biophysical Journal*, 90: 3694–3703. <https://doi.org/10.1529/biophysj.105.074856>.
- Tora, L. and Timmers, H.T.M. (2010). The TATA box regulates TATA-binding protein (TBP) dynamics in vivo. *Trends in Biochemical Sciences*, 35(6): 309–314. <https://doi.org/10.1016/j.tibs.2010.01.007>.
- van Huijsduijnen, R.A.M.H., Bollekens, J., Dorn, A., Benoist, C. and Mathis, D. (1987). Nucleic acids research. *Nucleic Acids Research* 15(18): 7265–7282
- Varalakshmi, K.N., Kumudini, B.S., Nandini, B.N., Solomon, J., Suhas, R., Mahesh, B. and Kavitha, A.P. (2009). Production and characterization of alpha amylase from *Aspergillus niger* JGI 24 isolated in Bangalore. *Polish Journal of Microbiology*, 58(1): 29–36.
- Veerapagu, M., Jeya, K.R. and Sankaranarayanan, A. (2016). Screening and production of fungal amylase from *Aspergillus* sp. by SSF. *Journal of Global Biosciences*, 5(8): 4443–4450.
- Verdoes, J.C., Punt, P.J., Schrickx, J.M., van Verseveld, H.W., Stouthamer, A.H. and van den Hondel, C.A.M.J.J. (1993). Glucoamylase overexpression in

- Aspergillus niger*: Molecular genetic analysis of strains containing multiple copies of the *glaA* gene. *Transgenic Research* 2(2): 84–92. <https://doi.org/10.1007/BF01969381>
- Verdoes, J.C., Punt, P.J., Stouthamer, A.H. and van den Hondel, C.A.M.J.J. (1994). The effect of multiple copies of the upstream region on expression of the *Aspergillus niger* glucoamylase-encoding gene. *Gene* 145(2): 179–187. [https://doi.org/10.1016/0378-1119\(94\)90003-5](https://doi.org/10.1016/0378-1119(94)90003-5)
- Verma, N., Hung, K. and Kang, J.J. (2013). Differential utilization of TATA box-binding protein (TBP) and TBP-related factor 1 (*trf1*) at different classes of RNA polymerase iii promoters. *Journal of Biological Chemistry*, 288(38): 27564–27570. <https://doi.org/10.1074/jbc.C113.503094>.
- Verma, S., Gazara, R.K., Nizam, S., Parween, S., Chattopadhyay, D. and Verma, P.K. (2016). Draft genome sequencing and secretome analysis of fungal phytopathogen *Ascochyta rabiei* provides insight into the necrotrophic effector repertoire. *Scientific Reports*, 6: 1–14. <https://doi.org/10.1038/srep24638>.
- Wang, B., Li, J., Gao, J., Cai, P., Han, X. and Tian, C. (2017). Identification and characterization of the glucose dual-affinity transport system in *Neurospora crassa*: Pleiotropic roles in nutrient transport, signaling, and carbon catabolite repression. *Biotechnology for Biofuels*, 10(1): 1–22. <https://doi.org/10.1186/s13068-017-0705-4>.
- Wang, Z., Deak, M. and Free, S.J. (1994). A cis-acting region required for the regulated expression of *grg-1*, a neurospora glucose-repressible gene: Two regulatory sites (CRE and NRS) are required to repress *grg-1* expression. *Journal of Molecular Biology*, 237(1): 65–74. <https://doi.org/10.1006/jmbi.1994.1209>.
- Watanabe, T. (2002). Pictorial atlas of soil and seed fungi: morphologies of cultured fungi and key to species. In CRC Press (Second edition). CRC Press. <https://doi.org/10.5860/choice.40-3398>.
- Wirsel, S., Lachmund, A., Wildhardt, G. and Ruttkowski, E. (1989). Three α -amylase genes of *Aspergillus oryzae* exhibit identical intron-exon organization. *Molecular Microbiology*, 3(1): 3–14. <https://doi.org/10.1111/j.1365-2958.1989.tb00097.x>.
- Xu, Q., Cao, Y., Li, X., Liu, L., Qin, S., Wang, Y., Cao, Y., Xu, H. and Qiao, D. (2018). Purification and characterization of a novel intracellular α -amylase with a wide variety of substrates hydrolysis and transglycosylation activity from *Paenibacillus* sp. SSG-1. *Protein Expression and Purification*, 144: 62–70. <https://doi.org/10.1016/j.pep.2016.04.007>.

- Yamashita, M., Tsujikami, M., Murata, S., Kobayashi, T., Shimizu, M. and Kato, M. (2021). Artificial AmyR::XlnR transcription factor induces α -amylase production in response to non-edible xylan-containing hemicellulosic biomass. *Enzyme and Microbial Technology*, 145: 109762. <https://doi.org/10.1016/j.enzmictec.2021.109762>.
- Yao, T.T., Wang, Y.M., Gu, J.L. and Wang, Z.X. (2006). Overproduction of glucoamylase by recombinant *Aspergillus niger* harboring multiple copies of glaA. *Chinese Journal of Biotechnology* 22(4): 567–571. [https://doi.org/10.1016/S1872-2075\(06\)60045-9](https://doi.org/10.1016/S1872-2075(06)60045-9)
- Yu, R. and Hang, Y.D. (1991). Purification and characterization of a glucoamylase from *Rhizopus oryzae*. *Food Chemistry*, 40(3): 301–308. [https://doi.org/10.1016/0308-8146\(91\)90114-4](https://doi.org/10.1016/0308-8146(91)90114-4).
- Zangirolami, T.C., Carlsen, M., Nielsen, J. and Jorgensen, S.B. (2002). Growth and enzyme production during continuous cultures of a high amylase-producing variant of *Aspergillus oryzae*. *Brazilian Journal of Chemical Engineering*, 19(1): 1–32. <https://doi.org/10.1590/S0104-66322002000100005>.
- Zhang, Q., Han, Y. and Xiao, H. (2017). Microbial α -amylase: A biomolecular overview. *Process Biochemistry*, 53: 88–101. <https://doi.org/10.1016/j.procbio.2016.11.012>.
- Zhao, Z., Liu, H., Luo, Y., Zhou, S., An, L., Wong, C., Jin, Q., Zhou, M. and Xu J. (2014). Molecular evolution and functional fungal tree of life. *Scientific Reports*, 4: 1–13. <https://doi.org/10.1038/srep06746>.
- Zoll, J., Snelders, E., Verweij, P.E. and Melchers, W.J.G. (2016). Next-generation sequencing in the mycology lab. *Current Fungal Infection Reports*, 10: 37–42. <https://doi.org/10.1007/s12281-016-0253-6>.