

DAFTAR PUSTAKA

- Abdelaziz, M. N., Xuan, T. D., Mekawy, A. M. M., Wang, H., & Khanh, T. D. (2018). Relationship of salinity tolerance to Na⁺ exclusion, proline accumulation, and antioxidant enzyme activity in rice seedlings. *Agriculture*, 8(11), 166.
- Ahmadizadeh, M., Vispo, N. A., Calapit-Palao, C. D. O., Pangaan, I. D., Viña, C. D., & Singh, R. K. (2016). Reproductive stage salinity tolerance in rice: a complex trait to phenotype. *Indian Journal of Plant Physiology*, 21(4), 528–536.
- Assaha, D. V., Ueda, A., Saneoka, H., Al-Yahyai, R., & Yaish, M. W. (2017). The role of Na⁺ and K⁺ transporters in salt stress adaptation in glycophytes. *Frontiers in physiology*, 8, 509.
- Bahcesular, B., Yildirim, E. D., Karaçocuk, M., Kulak, M., & Karaman, S. (2020). Seed *priming* with melatonin effects on growth, essential oil compounds and antioxidant activity of basil (*Ocimum basilicum* L.) under salinity stress. *Industrial Crops and Products*, 146, 112165.
- Bates, L. S., Waldren, R. P., & Teare, I. D. (1973). Rapid determination of free proline for water-stress studies. *Plant and soil*, 39(1), 205–207.
- Bdr, M. F., Musa, Y., Nasaruddin, N., Ridwan, I., & Yassi, A. H. (2020). Screening of Indonesian rice (*Oryza sativa* L.) genotypes against salinity stress using NaCl under hydroponic condition. *IOP Conference Series: Earth and Environmental Science*, 575, 012111.
- Demiral, T., & Türkan, I. (2005). Comparative lipid peroxidation, antioxidant defense systems and prolina content in roots of two rice cultivars differing in salt tolerance. *Environmental and experimental botany*, 53(3), 247–257.
- Fageria, N. K. (2007). Yield physiology of rice. *Journal of Plant Nutrition*, 30(6), 843–879.
- Fahad, S., Hussain, S., Matloob, A., Khan, F. A., Khaliq, A., Saud, S., Hassan, S., Shan, D., Khan, F., Ullah, N., Faiq, M., Khan, M. R., Tareen, A. K., Khan, A., Ullah, A., Ullah, N., & Huang, J. (2014). Phytohormones and plant responses to salinity stress: a review. *Plant Growth Regulation*, 75(2), 391–404.
- Ghosh, B., Ali, M. N., & Saikat, G. (2016). Response of rice under salinity stress: a review update. *J Res Rice*, 4(2), 167–174.
- Hairmansis, A., Nafisah, N., & Jamil, A. (2017). Towards developing salinity tolerant rice adaptable for coastal regions in Indonesia. *KnE Life Sciences*, 2(6), 72.
- Hidayah, A., Nisak, R. R., Susanto, F. A., Nuringtyas, T. R., Yamaguchi, N., & Purwestri, Y. A. (2022). Seed *Halopriming* Improves Salinity Tolerance of Some Rice Cultivars During Seedling Stage. *Botanical Studies*, 63(1), 1–12.
- Hoang, T. M. L., Tran, T. N., Nguyen, T. K. T., Williams, B., Wurm, P., Bellairs, S., & Mundree, S. (2016). Improvement of salinity stress tolerance in rice: challenges and opportunities. *Agronomy*, 6(4), 54.

- Hussain, S., Zhang, J., Zhong, C., Zhu, L., Cao, X., Yu, S., Bohr, J. A., JiJie, H., & Jin, Q. (2017). Effects of salt stress on rice growth, development characteristics, and the regulating ways: A review. *Journal of Integrative Agriculture*, 16(11), 2357–2374.
- Iedan, S. F., & Alag, M. K. (2021). The role of *priming* seeds with salicylic acid in improving some vegetative growth characteristics of sweet corn under water stress conditions. *Plant Archives*, 21(Suppliment-1), 1548–1554.
- IRRI. (2002). The standard evaluation system for rice. International Rice Research International. <http://www.knowledgebank.irri.org/images/docs/rice-standard-evaluation-system.pdf>
- Isayenkov, S. V., & Maathuis, F. J. M. (2019). Plant salinity stress: many unanswered questions remain. *Frontiers in plant science*, 10, 80.
- ITIS - Report: *Oryza sativa*. (n.d.).
https://www.itis.gov/servlet/SingleRpt/SingleRpt?search_topic=TSN&search_value=41976#null
- Ito, V. C., & Lacerda, L. G. (2019). Black rice (*Oryza sativa* L.): A review of its historical aspects, chemical composition, nutritional and functional properties, and applications and processing technologies. *Food chemistry*, 301, 125304.
- Jafar, M., Farooq, M., Cheema, M., Afzal, I., Basra, S. M. A., Wahid, M. A., Aziz, T., & Shahid, M. (2011). Improving the performance of wheat by seed *priming* under saline conditions. *Journal of Agronomy and Crop Science*, 198(1), 38–45.
- Jisha, K. C., & Puthur, J. T. (2014). Seed *halopriming* outdo *hydropriming* in enhancing seedling vigor and osmotic stress tolerance potential of rice varieties. *Journal of Crop Science and Biotechnology*, 17, 209–219
- Johnson, R., & Puthur, J. T. (2021). Seed *priming* as a cost effective technique for developing plants with cross tolerance to salinity stress. *Plant physiology and biochemistry*, 162, 247–257.
- Kementerian Pertanian RI. (2023).
https://satudata.pertanian.go.id/assets/docs/publikasi/Buku_Statistik_Konsumsi_2022.pdf
- Khan, M. O., Irfan, M., Muhammad, A., Ullah, I., Nawaz, S., Khalil, M. K., & Ahmad, M. (2022). A practical and economical strategy to mitigate salinity stress through seed *priming*. *Frontiers in Environmental Science*, 1613.
- Kibria, M. G., Hossain, M., Murata, Y., & Hoque, M. A. (2017). Antioxidant defense mechanisms of salinity tolerance in rice genotypes. *Rice Science*, 24(3), 155–162.
- Kumar, K., Kumar, M., Kim, S. R., Ryu, H., & Cho, Y. G. (2013). Insights into genomics of salt stress response in rice. *Rice*, 6(1), 1–15.
- Kumari, S., Panjabi nee Sabharwal, V., Kushwaha, H. R., Sopory, S. K., Singla-Pareek, S. L., & Pareek, A. (2009). Transcriptome map for seedling stage

specific salinity stress response indicates a specific set of genes as candidate for saline tolerance in *Oryza sativa* L. *Functional & Integrative Genomics*, 9, 109–123.

- Lichtenthaler, H. K., & Wellburn, A. R. (1983). Determinations of total carotenoids and chlorophylls a and b of leaf extracts in different solvents. *Biochemical Society Transactions*, 11(5), 591–592.
- Liu, C., Mao, B., Yuan, D., Chu, C., & Duan, M. (2022). Salt tolerance in rice: Physiological responses and molecular mechanisms. *The Crop Journal*, 10(1), 13–25.
- Livak, K. J., & Schmittgen, T. D. (2001). Analysis of relative gene expression data using real-time quantitative PCR and the 2- $\Delta\Delta$ CT method. *Methods*, 25(4), 402–408.
- Ludwiczak, A., Osiak, M., Cárdenas-Pérez, S., Lubińska-Mielińska, S., & Piernik, A. (2021). Osmotic stress or ionic composition: which affects the early growth of crop species more?. *Agronomy*, 11(3), 435.
- Manimaran, P., Reddy, S. V., Moin, M., Reddy, M. R., Yugandhar, P., Mohanraj, S. S., Balachandran, S. M., & Kirti, P. B. (2017). Activation-tagging in indica rice identifies a novel transcription factor subunit, NF-YC13 associated with salt tolerance. *Scientific Reports*, 7(1).
- Mansour, M. M. F., & Ali, E. F. (2017). Evaluation of proline functions in saline conditions. *Phytochemistry*, 140, 52–68.
- Maoka, T. (2020). Carotenoids as natural functional pigments. *Journal of natural medicines*, 74(1), 1–16.
- Marklund, S., & Marklund, G. (1974). Involvement of the superoxide anion radical in the autoxidation of pyrogallol and a convenient assay for superoxide dismutase. *European journal of biochemistry*, 47(3), 469–474.
- Mishra, P., Bhoomika, K., & Dubey, R. S. (2013). Differential responses of antioxidative defense system to prolonged salinity stress in salt-tolerant and salt-sensitive Indica rice (*Oryza sativa* L.) seedlings. *Protoplasma*, 250, 3–19.
- Nakano, Y., & Asada, K. (1981). Hydrogen peroxide is scavenged by ascorbate-specific peroxidase in spinach chloroplasts. *Plant and cell physiology*, 22(5), 867–880.
- Nakhoda, B., Leung, H., Mendioro, M. S., Mohammadi-nejad, G., & Ismail, A. M. (2012). Isolation, characterization, and field evaluation of rice (*Oryza sativa* L., Var. IR-64) mutants with altered responses to salt stress. *Field Crops Research*, 127, 191–202.
- Negrão, S., Schmöckel, S. M., & Tester, M. J. A. O. B. (2017). Evaluating physiological responses of plants to salinity stress. *Annals of botany*, 119(1), 1–11.

- Noorzuraini, A. R. S., Ramdzan, O. M., Idayu, A., & Hafiz, M. (2021). Evaluating the rice germplasm for salinity tolerance based on phenotypic traits. *IOP Conference Series*, 736(1), 012067.
- Panga, N. J., Sjahril, R., & Kaimuddin. (2019). Seed *priming* effect on Banyuasin and IR-64 rice varieties growth at the seedling stage under salinity stress. *IOP Conference Series: Earth and Environmental Science*, 343(1), 012171.
- Paparella, S., Araújo, S. S., Rossi, G., Wijayasinghe, M., Carbonera, D., & Balestrazzi, A. (2015). Seed *priming*: state of the art and new perspectives. *Plant cell reports*, 34(8), 1281–1293.
- Parihar, P., Singh, S., Singh, R., Singh, V. P., & Prasad, S. M. (2015). Effect of salinity stress on plants and its tolerance strategies: a review. *Environmental science and pollution research*, 22(6), 4056–4075.
- Rafii, M. Y., Rafii, M. Y., Yusoff, M. M., Ali, N. S., Oladosu, Y., Datta, D. R., Anisuzzaman, M., & Ikbal, M. F. (2021). Advanced breeding strategies and future perspectives of salinity tolerance in rice. *Agronomy*, 11(8), 1631.
- Reddy, I. N. B. L., Kim, B. K., Yoon, I. S., Kim, K. H., & Kwon, T. R. (2017). Salt tolerance in rice: focus on mechanisms and approaches. *Rice Science*, 24(3), 123–144.
- Rhoades, J. D., Kandiah, A., & Mashali, A. M. (1992). The use of saline waters for crop production-FAO irrigation and drainage paper 48. *FAO, Rome*, 133.
- Sairam, R. K., Deshmukh, P. S., & Shukla, D. S. (1997). Tolerance of drought and temperature stress in relation to increased antioxidant enzyme activity in wheat. *Journal of Agronomy and Crop Science*, 178(3), 171–178.
- Scardaci, S., Shannon, M., Grattan, S., Eke, A., Roberts, S., Goldman-Smith, S., & Hill, J. (2002). Water management practices can affect salinity in rice fields. *California Agriculture*, 56(6), 184–188.
- Shakri, T., Che-Othman, M. H., Md Isa, N., Sukiran, N. L., & Zainal, Z. (2022). Morpho-physiological and stress-related gene expression of rice varieties in response to salinity stress at early vegetative stage. *Agriculture*, 12(5), 638.
- Silva, F., Stevens, C. J., Weisskopf, A., Castillo, C., Qin, L., Bevan, A., & Fuller, D. Q. (2015). Modelling the geographical origin of rice cultivation in Asia using the rice archaeological database. *PloS one*, 10(9), e0137024.
- Sriskantharajah, K., Osumi, S., Chuamnakhong, S., Nampei, M., Amas, J. C., Gregorio, G. B., & Ueda, A. (2020). Contribution of two different Na⁺ transport systems to acquired salinity tolerance in rice. *Plant Science*, 297, 110517.
- Subekti, N. A., Sembiring, H., Nugraha, D., & Priatmojo, B. (2020). Yield of different rice cultivars at two levels of soil salinity under seawater intrusion in West Java, Indonesia. *Biodiversitas Journal of Biological Diversity*, 21(1).

- Subramanyam, K., Du Laing, G., & Van Damme, E. J. (2019). Sodium selenate treatment using a combination of seed *priming* and foliar spray alleviates salinity stress in rice. *Frontiers in plant science*, 10, 116.
- Talubaghi, M. J., Daliri, M. S., Mazloun, P., Rameeh, V., & Mousavi, A. (2023). Effect of salt stress on growth, physiological and biochemical parameters and activities of antioxidative enzymes of rice cultivars. *Cereal Research Communications*, 51(2), 403–411.
- Tamimi, S. M. (2016). Effect of seed *priming* on growth and physiological traits of five Jordanian wheat (*Triticum aestivum* L.) landraces under salt stress. *J Biosci Agric Res*, 11(1), 906–922.
- Theerakulpisut, P., Kanawapee, N., & Panwong, B. (2017). Seed *priming* alleviated salt stress effects on rice seedlings by improving Na⁺/K⁺ and maintaining membrane integrity. *International Journal of Plant Biology*, 7(1), 6402.
- Vighi, I. L., Benitez, L. C., Amaral, M. N. D., Moraes, G. P., Auler, P. A., Rodrigues, G., Deuner, S., Da Maia, L. C., & Braga, E. J. B. (2017). Functional characterization of the antioxidant enzymes in rice plants exposed to salinity stress. *Biologia Plantarum*, 61(3), 540–550.
- Wang, W., Zhao, X., Li, M., Huang, L., Xu, J., Zhang, F., Cui, Y., Fu, B., & Li, Z. (2015). Complex molecular mechanisms underlying seedling salt tolerance in rice revealed by comparative transcriptome and metabolomic profiling. *Journal of Experimental Botany*, 67(1), 405–419.
- Wangsawang, T., Chuamnakhong, S., Kohnishi, E., Sripichitt, P., Sreewongchai, T., & Ueda, A. (2018). A salinity-tolerant japonica cultivar has Na⁺ exclusion mechanism at leaf sheaths through the function of a Na⁺ transporter OsHKT 1; 4 under salinity stress. *Journal of Agronomy and Crop Science*, 204(3), 274–284.
- Yadav, P., Singh, P., Yadav, B., Pandey, S., Dubey, V., Singh, V., & Dwivedi, D. K. (2018). Evaluation of rice genotypes for salinity tolerance at seedling stage. *Journal of Pharmacognosy and Phytochemistry*, 7(4S), 202–205.
- Yoshida, S. (1981). Fundamentals of rice crop science. Int. Rice Res. Inst..
- Youssef, N. B., Shaher Al-Moaikal, R. M., Saleh Al-Hawas, G. H., & Alrammah, F. M. (2020). Up-modulation of membrane lipid composition and functionality by seed *priming* under salinity in the Hasawi rice variety. *Journal of Applied Botany & Food Quality*, 93.
- Zhu, L., Sun, J., Wu, G., Wang, Y., Wang, L., Qian, H., & Qi, X. (2018). Identification of rice varieties and determination of their geographical origin in China using Raman spectroscopy. *Journal of Cereal Science*, 82, 175–182.