

REFERENCES

- Azevedo, F. C., Lemos, F. G., Freitas-Junior, M. C., Rocha, D. G., & Azevedo, F. C. C. (2018). Puma activity patterns and temporal overlap with prey in a human-modified landscape at Southeastern Brazil. *Journal of Zoology*, 305(4), 246–255. <https://doi.org/10.1111/jzo.12558>
- Ballari, S. A., & Barrios-García, M. N. (2014). A review of wild boar *Sus scrofa* diet and factors affecting food selection in native and introduced ranges. *Mammal Review*, 44(2), 124–134. <https://doi.org/10.1111/MAM.12015/SUPPINFO>
- Barros, A. L., Curveira-Santos, G., Marques, T. A., & Santos-Reis, M. (2020). Accounting for detection unveils the intricacy of wild boar and rabbit co-occurrence patterns in a Mediterranean landscape. *Scientific Reports*, 10(1). <https://doi.org/10.1038/s41598-020-63492-9>
- Brodie, J. F., Helmy, O. E., Mohd-Azlan, J., Granados, A., Bernard, H., Giordano, A. J., & Zipkin, E. (2018). Models for assessing local-scale co-abundance of animal species while accounting for differential detectability and varied responses to the environment. *Biotropica*, 50(1), 5–15. <https://doi.org/10.1111/btp.12500>
- Cahill, S., Llimona, F., & Gràcia, J. (2003). Spacing and nocturnal activity of wild boar *Sus scrofa* in a Mediterranean metropolitan park. *Wildlife Biology*, 9(SUPPL. 1), 3–13. <https://doi.org/10.2981/wlb.2003.058>
- Caruso, N. I., J Valenzuela, A. E., Burdett, C. L., Luengos Vidal, E. M., Birochio, D., & CasanaveID, E. B. (2018). *Summer habitat use and activity patterns of wild boar *Sus scrofa* in rangelands of central Argentina*. <https://doi.org/10.1371/journal.pone.0206513>
- Chatani, K. (2003). Positional behavior of free-ranging Japanese macaques (*Macaca fuscata*). *Primates*, 44(1), 13–23. <https://doi.org/10.1007/s10329-002-0002-z>
- Chetri, M., Odden, M., Sharma, K., Flagstad, Ø., & Wegge, P. (2019). Estimating snow leopard density using fecal DNA in a large landscape in north-central

Nepal. *Global Ecology and Conservation*, 17.
<https://doi.org/10.1016/J.GECCO.2019.E00548>

Colyn, R. B., Radloff, F. G. T., & O'Riain, M. J. (2018). Camera trapping mammals in the scrubland's of the Cape Floristic Kingdom—the importance of effort, spacing and trap placement. *Biodiversity and Conservation*, 27(2), 503–520. <https://doi.org/10.1007/S10531-017-1448-Z/FIGURES/7>

Crombie, A. C. (1947). Interspecific Competition. *The Journal of Animal Ecology*, 16(1), 44. <https://doi.org/10.2307/1506>

Eccard, J. A., & Ylönen, H. (2002). *Direct Interference or Indirect Exploitation? An Experimental Study of Fitness Costs of Interspecific Competition in Voles* (Vol. 99, Issue 3). <https://about.jstor.org/terms>

Franklin, Janet, Miller, & Jennifer A. (2009). *Mapping Species Distributions: Spatial Inference and Prediction*. www.cambridge.org/9780521876353

Garza, S. J., Tabak, M. A., Miller, R. S., Farnsworth, M. L., & Burdett, C. L. (2018). Abiotic and biotic influences on home-range size of wild pigs (*Sus scrofa*). *Journal of Mammalogy*, 99(1), 97–107. <https://doi.org/10.1093/jmammal/gyx154>

Geiger, M. (2021). Pattern and pace of morphological change due to variable human impact: the case of Japanese macaques. *Primates*, 62, 955–970. <https://doi.org/10.1007/s10329-021-00933-7>

Hallett, J. G., O'connell, M. A., & Honeycutt, R. L. (1983). Competition and Habitat Selection: Test of a Theory Using Small Mammals. *Oikos*, 40(2), 175–181.

Hamada, Y., & Yamamoto, A. (2010). Chapter 2: Morphological Characteristics, Growth, and Aging in Japanese Macaques. In *The Japanese Macaques* (pp. 27–33).

Hanya, G. (2004). Diet of a Japanese Macaque Troop in the Coniferous Forest of Yakushima. *International Journal of Primatology*, 12(1).

Hanya, G., Noma, N., & Agetsuma, N. (2003). Altitudinal and seasonal variations in the diet of Japanese macaques in Yakushima. *Primates 2003* 44:1, 44(1), 51–59. <https://doi.org/10.1007/S10329-002-0007-7>



- Hanya, G., Yoshihiro, S., Zamma, K., Matsubara, H., Ohtake, M., Kubo, R., Noma, N., Agetsuma, N., & Takahata, Y. (2004). Environmental determinants of the altitudinal variations in relative group densities of Japanese macaques on Yakushima. *Ecological Research*, 19(5), 485–493.
<https://doi.org/10.1111/J.1440-1703.2004.00662.X>
- Haynes, T. B., Schmutz, J. A., Lindberg, M. S., Wright, K. G., Uher-Koch, B. D., & Rosenberger, A. E. (2014). Occupancy of yellow-billed and Pacific loons: Evidence for interspecific competition and habitat mediated co-occurrence. *Journal of Avian Biology*, 45(3), 296–304. <https://doi.org/10.1111/jav.00394>
- Higashide, D., Kuriyama, T., Takagi, S., Nakashima, Y., Fukasawa, K., Yajima, G., Kasada, M., & Yokoyama, M. (2021). Effectiveness of signs of activity as relative abundance indices for wild boar. *Wildlife Biology*, 2021(4).
<https://doi.org/10.2981/wlb.00869>
- Honda, T. (2009). Environmental factors affecting the distribution of the wild boar, sika deer, Asiatic black bear and Japanese macaque in Central Japan, with implications for human-wildlife conflict. *Mammal Study*, 34(2), 107–116.
<https://doi.org/10.3106/041.034.0206>
- Ikeda, T., Asano, M., & Suzuki, M. (2021). The potential negative impacts of the classical swine fever virus on wild boar population in Gifu prefecture, Japan. *J. Vet. Med. Sci*, 83(5), 846–849. <https://doi.org/10.1292/jvms.20-0675>
- Ikeda, T., Uchida, K., Matsuura, Y., Takahashi, H., Yoshida, T., Kaji, K., & Koizumi, I. (2016). *Seasonal and Diel Activity Patterns of Eight Sympatric Mammals in Northern Japan Revealed by an Intensive Camera-Trap Survey*.
<https://doi.org/10.1371/journal.pone.0163602>
- Izumiyama, S., Mochizuki, T., & Shiraishi, T. (2003). Troop size, home range area and seasonal range use of the Japanese macaque in the Northern Japan Alps. *Ecological Research*, 18(5), 465–474. <https://doi.org/10.1046/J.1440-1703.2003.00570.X>
- Jacquin, A., Goulard, M., Hutchinson, J. M. S., & Hutchinson, S. L. (2015). A Statistical Approach For Predicting Grassland Degradation in Disturbance-driven Landscapes. *2015 8th International Workshop on the Analysis of*



Multitemporal Remote Sensing Images (Multi-Temp), 1–5.

<https://doi.org/10.1109/Multi-Temp.2015.7245759>

- Johann, F., Handschuh, M., Linderoth, P., Dormann, C. F., & Arnold, J. (2020). Adaptation of wild boar (*Sus scrofa*) activity in a human-dominated landscape. *BMC Ecology*, 20(1). <https://doi.org/10.1186/S12898-019-0271-7>
- Kaneko, Y., Kanda, E., Tashima, S., Masuda, R., Newman, C., & MacDonald, D. W. (2014). The socio-spatial dynamics of the Japanese badger (*Meles anakuma*). *Journal of Mammalogy*, 95(2), 290–300. <https://doi.org/10.1644/12-MAMM-A-158>
- Kaneko, Y., Maruyama, N., & Macdonald, D. W. (2006). Food habits and habitat selection of suburban badgers (*Meles meles*) in Japan . *Journal of Zoology*, 270(1), 78–89. <https://doi.org/10.1111/j.1469-7998.2006.00063.x>
- Kaneko, Y., Masuda, R. &, & Abramov, A. v. (2016). *Meles anakuma, Japanese Badger THE IUCN RED LIST OF THREATENED SPECIES™*. <https://doi.org/10.2305/IUCN.UK.2016>
- Kauhala, K. &, & Saeki, M. (2016). *Nyctereutes procyonoides. The IUCN Red List of Threatened Species*. <https://doi.org/10.2305/IUCN.UK.2016>
- Keuling, O., & Leus, K. (2019). *Sus scrofa, Wild Boar THE IUCN RED LIST OF THREATENED SPECIES™*. <https://doi.org/10.2305/IUCN.UK.2019-3.RLTS.T41775A44141833.en>
- Keuling, O., Podgórski, T., Monaco, A., Melletti, M., Merta, D., Albrycht, M., Genov, P. v., Gethöffer, F., Vetter, S. G., Jori, F., Scalera, R., & Gongora, J. (2017). *Ecology, Conservation and Management of Wild Pigs and Peccaries* (M. Melletti & E. Meijaard, Eds.). Cambridge University Press. <https://doi.org/10.1017/9781316941232>
- Kitao, N., Fukui, D., Hashimoto, M., & Osborne, P. G. (2009). Overwintering strategy of wild free-ranging and enclosure-housed Japanese raccoon dogs (*Nyctereutes procyonoides albus*). *International Journal of Biometeorology*, 53(2), 159–165. <https://doi.org/10.1007/S00484-008-0199-7/TABLES/2>
- Koper, N., & Manseau, M. (2012). A guide to developing resource selection functions from telemetry data using generalized estimating equations and

generalized linear mixed models. *Rangifer*, 32(2), 195–204.
<https://doi.org/10.7557/2.32.2.2269>

Kurose, N., Kaneko, Y., Abramov, A. v., Siriaroonrat, B., & Masuda, R. (2001). Low genetic diversity in Japanese populations of the Eurasian badger *Meles meles* (Mustelidae, Carnivora) revealed by mitochondrial cytochrome b gene sequences. *Zoological Science*, 18(8), 1145–1151.
<https://doi.org/10.2108/zsj.18.1145>

Lee, S. M., & Lee, E. J. (2019). Diet of the wild boar (*Sus scrofa*): Implications for management in forest-agricultural and urban environments in South Korea. *PeerJ*, 2019(10). <https://doi.org/10.7717/PEERJ.7835/>

Manly, B. F. J., McDonald, L. L., & Thomas, D. L. (1994). Review Reviewed Work(s): Resource Selection by Animals: Statistical Design and Analysis for Field Studies. *Source: Journal of Animal Ecology*, 63(3), 745–746.

Maruhashi, T., Saito, C., & Agetsuma, N. (1998). Home range structure and inter-group competition for land of Japanese macaques in evergreen and deciduous forests. *Primates* 1998 39:3, 39(3), 291–301.
<https://doi.org/10.1007/BF02573078>

Maruyama, T. E., Ueno, S., Mori, H., Kaneeda, T., & Moriguchi, Y. (2021). Factors Influencing Somatic Embryo Maturation in Sugi (Japanese Cedar, *Cryptomeria japonica* (Thunb. ex L.f.) D. Don).
<https://doi.org/10.3390/plants10050874>

Masaki, T., Takahashi, K., Sawa, A., Kado, T., Naoe, S., Koike, S., & Shibata, M. (2012). Fleshy fruit characteristics in a temperate deciduous forest of Japan: How unique are they? *Journal of Plant Research*, 125(1), 103–114.
<https://doi.org/10.1007/S10265-011-0423-0/FIGURES/6>

Meredith, M., & Ridout, M. (2021). *Overview of the overlap package*.

Meriggi, A., & Sacchi, O. (2009). Habitat requirements of wild boars in the northern Apennines (N Italy): A multi-level approach.
<Http://Dx.Doi.Org/10.1080/11250000109356382>, 68(1), 47–55.
<https://doi.org/10.1080/11250000109356382>



- Merow, C., Smith, M. J., & Silander, J. A. (2013). A practical guide to MaxEnt for modeling species' distributions: What it does, and why inputs and settings matter. *Ecography*, 36(10), 1058–1069. <https://doi.org/10.1111/J.1600-0587.2013.07872.X>
- Mitsuhashi, I., Sako, T., Teduka, M., Koizumi, R., Saito, M. U., & Kaneko, Y. (2018). Home range of raccoon dogs in an urban green area of Tokyo, Japan. *Journal of Mammalogy*, 99(3), 732–740. <https://doi.org/10.1093/JMAMMAL/GYY033>
- Niedballa, J., Wilting, A., Sollmann, R., Hofer, H., & Courtiol, A. (2019). Assessing analytical methods for detecting spatiotemporal interactions between species from camera trapping data. *Remote Sensing in Ecology and Conservation*, 5(3), 272–285. <https://doi.org/10.1002/rse2.107>
- Noda, H. M., Muraoka, H., & Nasahara, K. N. (2021). Plant ecophysiological processes in spectral profiles: perspective from a deciduous broadleaf forest. *Journal of Plant Research* 2021 134:4, 134(4), 737–751. <https://doi.org/10.1007/S10265-021-01302-7>
- O, A. F., Nichols, J. D., & Ullas Karanth, K. (2011). *Camera Traps in Animal Ecology: Methods and Analyses*. Springer. <https://doi.org/10.1007/978-4-431-99495-4>
- Ohashi, H., Saito, M., Horie, R., Tsunoda, H., Noba, H., Ishii, H., Kuwabara, T., Hiroshige, Y., Koike, S., Hoshino, Y., Toda, H., & Kaji, K. (2013). Differences in the activity pattern of the wild boar *Sus scrofa* related to human disturbance. *European Journal of Wildlife Research*, 59(2), 167–177. <https://doi.org/10.1007/s10344-012-0661-z>
- Osaki, A., Sashika, M., Abe, G., Shinjo, K., Fujimoto, A., Nakai, M., Shimozuru, M., & Tsubota, T. (2019). Comparison of feeding habits and habitat use between invasive raccoons and native raccoon dogs in Hokkaido, Japan. *BMC Ecology*, 19(1). <https://doi.org/10.1186/s12898-019-0249-5>
- Osugi, S., Trentin, B. E., & Koike, S. (2019). Impact of wild boars on the feeding behavior of smaller frugivorous mammals. *Mammalian Biology*, 97, 22–27. <https://doi.org/10.1016/J.MAMBIO.2019.03.013>

- Ridout, M. S., & Linkie, M. (2009a). Estimating overlap of daily activity patterns from camera trap data. *Journal of Agricultural, Biological, and Environmental Statistics*, 14(3), 322–337.
<https://doi.org/10.1198/jabes.2009.08038>
- Ridout, M. S., & Linkie, M. (2009b). Estimating overlap of daily activity patterns from camera trap data. *Journal of Agricultural, Biological, and Environmental Statistics*, 14(3), 322–337.
<https://doi.org/10.1198/jabes.2009.08038>
- Saeki, M., Johnson, P. J., & Macdonald, D. W. (2007). Movements and habitat selection of raccoon dogs (*Nyctereutes procyonoides*) in a mosaic landscape. *Journal of Mammalogy*, 88(4), 1098–1111. <https://doi.org/10.1644/06-MAMM-A-208R1.1/2/JMAMMAL-88-4-1098-FIG8.JPG>
- Saito, M., & Koike, F. (2013). Distribution of Wild Mammal Assemblages along an Urban-Rural-Forest Landscape Gradient in Warm-Temperate East Asia. *PLoS ONE*, 8(5), 65464. <https://doi.org/10.1371/journal.pone.0065464>
- Santos, M. J., Smith, A. B., Thorne, J. H., & Moritz, C. (2017). The relative influence of change in habitat and climate on elevation range limits in small mammals in Yosemite National Park, California, U.S.A. *Climate Change Responses* 2017 4:1, 4(1), 1–12. <https://doi.org/10.1186/S40665-017-0035-6>
- Schlichting, P. E., Fritts, S. R., Mayer, J. J., Gipson, P. S., & Dabbert, C. B. (2016). Determinants of variation in home range of wild pigs. *Wildlife Society Bulletin*, 40(3), 487–493. <https://doi.org/10.1002/WSB.662>
- Seki, Y., & Koganezawa, M. (2011). Factors influencing winter home ranges and activity patterns of raccoon dogs *Nyctereutes procyonoides* in a high-altitude area of Japan. *Acta Theriologica*, 56(2), 171–177. <https://doi.org/10.1007/S13364-010-0020-Y/FIGURES/4>
- Syfert, M. M., Smith, M. J., & Coomes, D. A. (2013). The Effects of Sampling Bias and Model Complexity on the Predictive Performance of MaxEnt Species Distribution Models. *PLoS ONE*, 8(2), 55158. <https://doi.org/10.1371/journal.pone.0055158>



- Tashima, S., Kaneko, Y., Anezaki, T., Baba, M., Yachimori, S., & Masuda, R. (2010). Genetic diversity within the Japanese badgers (*Meles anakuma*), as revealed by microsatellite analysis. *Mammal Study*, 35, 221–226.
- Tilman, D. (1987). The Importance of the Mechanisms of Interspecific Competition. *Source: The American Naturalist*, 129(5), 769–774.
<https://www.jstor.org/stable/2461735>
- Újváry, D., Schally, G., Buczkó, M., Szabó, L., & Szemethy, L. (2014). A simple method for the assessment of wild boars' (*Sus scrofa*) habitat use.
<https://doi.org/10.1016/j.jveb.2014.02.003>
- van Doormaal, N., Ohashi, H., Koike, S., & Kaji, K. (n.d.). Influence of human activities on the activity patterns of Japanese sika deer (*Cervus nippon*) and wild boar (*Sus scrofa*) in Central Japan. <https://doi.org/10.1007/s10344-015-0922-8>
- Vieira, E. M., & Paise, G. (2011). Temporal niche overlap among insectivorous small mammals. *Integrative Zoology*, 6(4), 375–386.
<https://doi.org/10.1111/J.1749-4877.2011.00266.X>
- Ward, O. G., & Wurster-Hill, D. H. (1990). *Nyctereutes procyonoides*. *Mamm Species*, 358(358), 1–5. <https://doi.org/10.2307/3504213>