



REFERENCES

- Al Khalili, S., Al Yaquobi, F., Al Abri, B., Al Thuhli, K., Al Marshoudi, S., Al Rawahi, B., & Al-Abri, S. (2022). Assessing Oman's knowledge, attitude and practice regarding tuberculosis: a cross-sectional study that calls for action. *International Journal of Infectious Diseases*, 124, S4–S11. <https://doi.org/10.1016/j.ijid.2022.06.002>
- Ali, S. M., Anjum, N., Naureen, F., Rashid, A., Tahir, A., Ishaq, M., & Usman, M. (2019). Satisfaction Level of Tuberculosis Patients Regarding Their Access to TB Care and Prevention Services, Delivered Through a Public–Private Mix Model in Pakistan. *Healthcare*, 7(4). <https://doi.org/10.3390/HEALTHCARE7040119>
- Alotaibi, B., Yassin, Y., Mushi, A., Maashi, F., Thomas, A., Mohamed, G., Hassan, A., & Yezli, S. (2019). Tuberculosis knowledge, attitude and practice among healthcare workers during the 2016 Hajj. *PLoS ONE*, 14(1), 1–15. <https://doi.org/10.1371/journal.pone.0210913>
- An, Y., Teo, A. K. J., Huot, C. Y., Tieng, S., Khun, K. E., Pheng, S. H., Leng, C., Deng, S., Song, N., Nonaka, D., & Yi, S. (2023). They do not have symptoms – why do they need to take medicines? Challenges in tuberculosis preventive treatment among children in Cambodia: a qualitative study. *BMC Pulmonary Medicine*, 23(1), 1–8. <https://doi.org/10.1186/s12890-023-02379-7>
- Article, O., Samal, J., & Dehury, R. K. (2017). *Impact of a Structured Tuberculosis Awareness Strategy on the Knowledge and Behaviour of the Families in a Slum Area in Chhattisgarh , India*. <https://doi.org/10.7860/JCDR/2017/24107.9489>
- Baloyi, D. P., Anthony, M. G., Meyerson, K. A., Mazibuko, S., Wademan, D., Viljoen, L., Myburgh, H., Preez, K., Osman, M., Charalambous, S., Hausler, H., & Hesseling, A. C. (2022). *Public Health Action*. 12(4), 159–164.
- Bastos, M. L., Campbell, J. R., Oxlade, O., Adjebimey, M., Trajman, A., Ruslami, R., Kim, H. J., Baah, J. O., Toelle, B. G., Long, R., Hoeppner, V., Elwood, K., Al-Jahdali, H., Apriani, L., Benedetti, A., Schwartzman, K., & Menzies, D. (2020). Health system costs of treating latent tuberculosis infection with four months of rifampin versus nine months of isoniazid in different settings. *Annals of Internal Medicine*, 173(3), 169–178. <https://doi.org/10.7326/M19-3741>
- Beyanga, M., Kidanya, B. R., Gerwing-Adima, L., Ochodo, E., Mshana, S. E., & Kasang, C. (2018). Investigation of household contacts of pulmonary tuberculosis patients increases case detection in Mwanza City, Tanzania. *BMC Infectious Diseases*, 18(1). <https://doi.org/10.1186/s12879-018-3036-6>



- CDC. (2012). Basic TB Facts. *CDC Factsheet*, 2–4.
- CDC. (2022). *Latent TB Infection and TB Disease / TB / CDC*. CDC.
- Chandra, D. K., Moll, A. P., Altice, F. L., Didomizio, E., Andrews, L., & Shenoi, S. V. (2022). Structural barriers to implementing recommended tuberculosis preventive treatment in primary care clinics in rural South Africa. *Global Public Health*, 17(4), 555–568. <https://doi.org/10.1080/17441692.2021.1892793>
- Cohen, A., Mathiasen, V. D., Schön, T., & Wejse, C. (2019). The global prevalence of latent tuberculosis: a systematic review and meta-analysis. *The European Respiratory Journal*, 54(3). <https://doi.org/10.1183/13993003.00655-2019>
- Cruz, A. T., & Starke, J. R. (2018). Completion rate and safety of tuberculosis infection treatment with shorter regimens. *Pediatrics*, 141(2). <https://doi.org/10.1542/peds.2017-2838>
- Datiko, D. G., Yassin, M. A., Theobald, S. J., & Cuevas, L. E. (2017). A community-based isoniazid preventive therapy for the prevention of childhood tuberculosis in Ethiopia. *International Journal of Tuberculosis and Lung Disease*, 21(9), 1002–1007. <https://doi.org/10.5588/ijtld.16.0471>
- Divala et al. (2022a). Public Health Action Campaign. *Public Health Action*, 12(1), Syzdikova, A., Zolfo, M., Malta, A., Diro, E., O.
- Divala et al. (2022b). Reasons for poor uptake of TB preventive therapy in South Africa D. *Public Health Action*, 12(1), Syzdikova, A., Zolfo, M., Malta, A., Diro, E., O.
- Fox, G. J., Nguyen, T. A., Coleman, M., Trajman, A., Velen, K., & Marais, B. J. (2021). Implementing tuberculosis preventive treatment in high-prevalence settings. *International Journal of Infectious Diseases*, 113, S13–S15. <https://doi.org/10.1016/J.IJID.2021.02.094>
- GLOBAL TB REPORT. (2022). Annual Report of Tuberculosis. In *Annual Global TB Report of WHO*.
- Gopichandran, V., Roy, P., Sitaram, A., & John, K. R. (2015). *Short Communication Impact of a Simple Educational Intervention on the Knowledge and Awareness of Tuberculosis among High School children in Vellore , India*. 35(1). <https://doi.org/10.4103/0970-0218.62589>
- Goroh, M. M. D., van den Boogaard, C. H. A., Lukman, K. A., Lowbridge, C., Juin, W. K., William, T., Jeffree, M. S., & Ralph, A. P. (2023). Factors affecting implementation of tuberculosis contact investigation and tuberculosis preventive therapy among children in Sabah, East Malaysia: A qualitative study. *PLoS ONE*, 18(5 May), 1–16.



Hopewell, P. C., & Jasmer, R. M. (2004). Overview of Clinical Tuberculosis. *Tuberculosis and the Tubercle Bacillus*, 13–31. <https://doi.org/10.1128/9781555817657.CH2>

Isaac, T., & Kizito, O. (2022). Factors Associated with Uptake of Isoniazid Preventive Therapy among Children under 5 Years Tuberculosis Contacts in Greater Kibaale, Uganda. *American Journal of Public Health Research*, 10(2), 76–89.

<https://doi.org/10.12691/ajphr-10-2-5>

Mandal, S., Bhatia, V., Sharma, M., Mandal, P. P., & Arinaminpathy, N. (2020). The potential impact of preventive therapy against tuberculosis in the WHO South-East Asian Region: A modelling approach. *BMC Medicine*, 18(1), 1–10.

<https://doi.org/10.1186/s12916-020-01651-5>

May, M., Id, D. G., Boogaard, C. H. A. Van Den, Id, A. L., Lowbridge, C., Ke, W., Id, J., William, T., Saffree, M., Id, J., & Ralph, A. P. (2023). *Factors affecting implementation of tuberculosis contact investigation and tuberculosis preventive therapy among children in Sabah , East Malaysia : A qualitative study*. 1–16.

<https://doi.org/10.1371/journal.pone.0285534>

Mebratu, W., Wedajo, S., Mohammed, S., Endawkie, A., & Damtew, Y. (2022). Prevalence and associated factors of tuberculosis among isoniazid users and non-users of HIV patients in Dessie, Ethiopia. *Scientific Reports*, 12(1), 1–10.

<https://doi.org/10.1038/s41598-022-16437-3>

Menzies, D., Adjibimey, M., Ruslami, R., Trajman, A., Sow, O., Kim, H., Obeng Baah, J., Marks, G. B., Long, R., Hoeppner, V., Elwood, K., Al-Jahdali, H., Gninafon, M., Apriani, L., Koesoemadinata, R. C., Kritski, A., Rolla, V., Bah, B., Camara, A., ... Benedetti, A. (2018). Four Months of Rifampin or Nine Months of Isoniazid for Latent Tuberculosis in Adults. *New England Journal of Medicine*, 379(5), 440–453.

<https://doi.org/10.1056/nejmoa1714283>

MOH-NSP. (2020). *National Strategic Plan for Ending TB 2020-24- EDITED - 20032020_FINAL_30March_withbudget*.

Murphy, A. L., & Gardner, D. M. (2019). Pilot Testing the Theoretical Framework of Acceptability in a Process Evaluation of a Community Pharmacy-Based Men's Mental Health Promotion Program. <Https://Doi.Org/10.1177/2158244019885129>, 9(4).

<https://doi.org/10.1177/2158244019885129>

Ngugi, S. K., Muiruri, P., Odero, T., & Gachuno, O. (2020). Factors affecting uptake and completion of isoniazid preventive therapy among HIV-infected children at a national



referral hospital, Kenya: A mixed quantitative and qualitative study. *BMC Infectious Diseases*, 20(1), 1–11. <https://doi.org/10.1186/s12879-020-05011-9>

NTP. (2020). *NTP Manual Timor-Leste*. 1–248.

Proctor, E., Silmere, H., Raghavan, R., Hovmand, P., Aarons, G., Bunger, A., Griffey, R., & Hensley, M. (2011). Outcomes for implementation research: Conceptual distinctions, measurement challenges, and research agenda. *Administration and Policy in Mental Health and Mental Health Services Research*, 38(2), 65–76.
<https://doi.org/10.1007/s10488-010-0319-7>

Rakhmawati, W., Fitri, S. Y. R., Sriati, A., & Hendrawati, S. (2022). Communication between parents and preschool-aged children about tuberculosis preventive treatment: A qualitative study. *Iranian Journal of Nursing and Midwifery Research*, 27(5), 370–376.
https://doi.org/10.4103/ijnmr.ijnmr_8_21

Reichler, M. R., Khan, A., Yuan, Y., Chen, B., McAuley, J., Mangura, B., Sterling, T. R., Team, T. E. S. C. T. O. 2, Bakhtawar, I., LeDoux, C., McAuley, J., Beison, J., Fitzgerald, M., Naus, M., Nakajima, M., Schluger, N., Hirsch-Moverman, Y., Moran, J., Blumberg, H., ... Luo, C. (2020). Duration of Exposure Among Close Contacts of Patients With Infectious Tuberculosis and Risk of Latent Tuberculosis Infection. *Clinical Infectious Diseases*, 71(7), 1627–1634. <https://doi.org/10.1093/CID/CIZ1044>

Rohman, H. (2019). *Pola_Spasial_dan_Aksesibilitas_Penggunaan*. 13(2), 49–55.

Sagili, K. D., Muniyandi, M., Shringarpure, K., Singh, K., Kirubakaran, R., Rao, R., Tonsing, J., Sachdeva, K. S., & Tharyan, P. (2022). Strategies to detect and manage latent tuberculosis infection among household contacts of pulmonary TB patients in high TB burden countries - a systematic review and meta-analysis. *Tropical Medicine and International Health*, 27(10), 842–863. <https://doi.org/10.1111/tmi.13808>

Salazar-Austin, N., Mulder, C., Hoddinott, G., Ryckman, T., Hanrahan, C. F., Velen, K., Chimoyi, L., Charalambous, S., & Chihota, V. N. (2022). Preventive Treatment for Household Contacts of Drug-Susceptible Tuberculosis Patients. *Pathogens*, 11(11), 1–13. <https://doi.org/10.3390/pathogens1111258>

Sekhon, M., Cartwright, M., & Francis, J. J. (2017). Acceptability of healthcare interventions : an overview of reviews and development of a theoretical framework. *BMC Health Services Research*, 1–13. <https://doi.org/10.1186/s12913-017-2031-8>

Sekhon, M., Cartwright, M., & Francis, J. J. (2022). Development of a theory-informed questionnaire to assess the acceptability of healthcare interventions. *BMC Health Services Research*, 22(1). <https://doi.org/10.1186/S12913-022-07577-3>



UNIVERSITAS
GADJAH MADA

Assessing the acceptability of TB preventive therapy among household contacts in Dili Municipality; Implementation research.

Etelvina De Jesus Da Costa , DR.dr.Yodi Mahendradhata.M.Sc, Ph.D, FRSRH ; DR. Dra. Ida Safitria Laksamawati, M

Universitas Gadjah Mada, 2023 | Diunduh dari <http://etd.repository.ugm.ac.id/>

Sharma, N., Basu, S., Khanna, A., Sharma, P., & Chandra, S. (2022). The intention to receive tuberculosis preventive therapy in adult household contacts of pulmonary TB patients in Delhi, India. In *Journal of Infection in Developing Countries* (Vol. 16, Issue 2).
<https://doi.org/10.3855/jidc.14910>

Sri Ayu Rahayu. (2019). DOI: <http://dx.doi.org/10.33846/sf10405> Pencegahan Tuberkulosis Paru dalam Keluarga: Kajian Literatur Sri Ayu Rahayu S. Paneo. *Jurnal Penelitian Kesehatan Suara Forikes*, 10(2), 270–274. <https://akper-sandikarsa.e-journal.id/JIKSH>

Suryavanshi, N., Murrill, M., Gupta, A., Hughes, M., Hesselink, A., Kim, S., Naini, L., Jones, L., Smith, B., Gupte, N., Dawson, R., Mave, V., Meshram, S., Mendoza-ticona, A., Sanchez, J., Kumarasamy, N., Comins, K., Conradie, F., Shenje, J., ... Nedsuwan, S. (2020). *Willingness to Take Multidrug-resistant Tuberculosis (MDR-TB) Preventive Therapy Among Adult and Adolescent Household Contacts of MDR-TB Index Cases : An International Multisite Cross-sectional Study*. 70. <https://doi.org/10.1093/cid/ciz254>

Tefera, F., Barnabee, G., Sharma, A., Feleke, B., Atnafu, D., Haymanot, N., O'Malley, G., & Feleke, G. (2019). Evaluation of facility and community-based active household tuberculosis contact investigation in Ethiopia: a cross-sectional study. *BMC Health Services Research*, 19(1), 1–9. <https://doi.org/10.1186/s12913-019-4074-5>

Triasih, R., Padmawati, R. S., Duke, T., Robertson, C., Sawyer, S. M., & Graham, S. M. (2016). A mixed-methods evaluation of adherence to preventive treatment among child tuberculosis contacts in Indonesia. *International Journal of Tuberculosis and Lung Disease*, 20(8), 1078–1083. <https://doi.org/10.5588/ijtld.15.0952>

Viney, K., Amaral, S., Marques, E. B., Siroka, A., Lopes, C., & Nery, S. V. (2019). Four of five tuberculosis patients experience catastrophic costs related to TB diagnosis and care in Timor-Leste. *The International Journal of Tuberculosis and Lung Disease : The Official Journal of the International Union against Tuberculosis and Lung Disease*, 23(11), 1191–1197. <https://doi.org/10.5588/ijtld.18.0765>

Weiner, B. J., Lewis, C. C., Stanick, C., Powell, B. J., Dorsey, C. N., Clary, A. S., Boynton, M. H., & Halko, H. (2017). Psychometric assessment of three newly developed implementation outcome measures. *Implementation Science*, 12(1), 1–12.
<https://doi.org/10.1186/s13012-017-0635-3>

WHO-TL. (2022). *Timor-Leste records a decline in Tuberculosis incidence, improved testing: Global TB Report*.

WHO. (2019). *Country Location type Incidence of tuberculosis (per 100 000 population per year)*. GLOBAL HEALTH OBSERVATORY DATA.



WHO. (2022). *Tuberculosis*.

WORLD BANK. (2014). *Poverty in Timor Leste 2014*.

Yetti, E., Tombeg, Z., & Hadi, A. J. (2023). Hubungan Sosial Budaya Dengan Upaya

Pencegahan TBC di Puskesmas Makale Kabupaten Tana Toraja. *JURNAL NERS*

(*Research & Learning in Nursing Science*), 7(2), 1364–1373.

Yuan, Y., Jin, J., Bi, X., Geng, H., Li, S., & Zhou, C. (2023). Factors associated with refusal
of preventive therapy after initial willingness to accept treatment among college students
with latent tuberculosis infection in Shandong, China. *BMC Infectious Diseases*, 23(1),
1–8. <https://doi.org/10.1186/s12879-023-08005-5>