



DAFTAR PUSTAKA

- Aboul-Maaty, N.A.-F. and Oraby, H.A.-S. (2019). Extraction of high-quality genomic DNA from different plant orders applying a modified CTAB-based method. *Bulletin of the National Research Centre*, [online] 43(1). doi:<https://doi.org/10.1186/s42269-019-0066-1>.
- Aguilar, M., Wang, X., Escalona, M., Yang, L. and Huang, L. (2022). Somatic embryogenesis of Arabica coffee in temporary immersion culture: Advances, limitations, and perspectives for mass propagation of selected genotypes. *Frontiers in Plant Science*, 13. doi:<https://doi.org/10.3389/fpls.2022.994578>.
- Barth, S., Geier, T., Klaus Eimert, Watillon, B., Sangwan, R.S. and Gleissberg, S. (2009). KNOX overexpression in transgenic Kohleria (Gesneriaceae) prolongs the activity of proximal leaf blastozones and drastically alters segment fate. *Planta*, 230(6), pp.1081–1091. doi:<https://doi.org/10.1007/s00425-009-0997-0>.
- Campos, N.A., Panis, B. and Carpentier, S.C. (2017). Somatic Embryogenesis in Coffee: The Evolution of Biotechnology and the Integration of Omics Technologies Offer Great Opportunities. *Frontiers in Plant Science*, 8. doi:<https://doi.org/10.3389/fpls.2017.01460>.
- Ciborowski, P. and Silberring, J. (2016). *Proteomic profiling and analytical chemistry : the crossroads*. 2nd ed. [online] Amsterdam ; Boston: Elsevier, pp.115–130. Available at: doi.org/10.1016/B978-0-444-63688-1.00007-0 [Accessed 24 Nov. 2023].
- Corthell, J.T. (2014). Agarose Gel Electrophoresis. *Basic Molecular Protocols in Neuroscience: Tips, Tricks, and Pitfalls*, [online] pp.21–25. doi:<https://doi.org/10.1016/b978-0-12-801461-5.00003-4>.
- Daúde, M.M., Freitas, N.C., Ságio, S.A., Paiva, L.V., Barreto, H.G. and Silva, S. (2020). Transcriptional analysis of WUSCHEL-related HOMEOBOX (WOX) genes in *Coffea arabica* L. *Biologia*, 75(9), pp.1483–1495. doi:<https://doi.org/10.2478/s11756-020-00460-8>.
- Drabik, A., Bodzoń-Kułakowska, A. and Silberring, J. (2016). Gel Electrophoresis. *Proteomic Profiling and Analytical Chemistry*, [online] pp.115–143. doi:<https://doi.org/10.1016/b978-0-444-63688-0-444-63688-1.00007-0>.
- Duverger, O. and Morasso, M.I. (2008). Role of homeobox genes in the patterning, specification, and differentiation of ectodermal appendages in mammals. *Journal of Cellular Physiology*, 216(2), pp.337–346. doi:<https://doi.org/10.1002/jcp.21491>.
- Emaus, M.N., Cagliero, C., Gostel, M.R., Johnson, G. and Anderson, J.L. (2022). Simple and efficient isolation of plant genomic DNA using magnetic ionic liquids. *Plant Methods*, 18(1). doi:<https://doi.org/10.1186/s13007-022-00860-8>.
- Gao, J., Yang, X., Zhao, W., Lang, T. and Tore Samuelsson (2015). Evolution, diversification, and expression of KNOX proteins in plants. *Frontiers Plant Science*, 6. doi:<https://doi.org/10.3389/fpls.2015.00882>.
- Garibyan, L. and Avashia, N. (2013). Polymerase Chain Reaction. *Journal of Investigative Dermatology*, [online] 133(3), pp.1–4. doi:<https://doi.org/10.1038/jid.2013.1>.
- Guerra, A.F., Santos, J. de F., Ferreira, L.T. and Rocha, O.C. (2021). Coffees of Brazil: research, sustainability and innovation. In: G. de CAPDEVILLE, ed., *Land-saving technologies*. [online] Embrapa, pp.61–74. Available at: <http://www.alice.cnptia.embrapa.br/alice/handle/doc/1139310> [Accessed 19 Dec. 2023].
- Gupta, N. (2019). DNA extraction and polymerase chain reaction. *Journal of Cytology*, 36(2), pp.116–117. doi:https://doi.org/10.4103/joc.joc_110_18.
- Hake, S., Smith, H.M.S., Holtan, H., Magnani, E., Mele, G. and Ramirez, J. (2004). THE ROLE OF KNOX GENES IN PLANT DEVELOPMENT. *Annual Review of Cell and Developmental Biology*, 20(1), pp.125–151. doi:<https://doi.org/10.1146/annurev.cellbio.20.031803.093824>.
- Han, Y., Zhang, C., Yang, H. and Jiao, Y. (2014). Cytokinin pathway mediates APETALA1 function in the establishment of determinate floral meristems in *Arabidopsis*. *Proceedings of the National Academy of Sciences*, 111(18), pp.6840–6845. doi:<https://doi.org/10.1073/pnas.1318532111>.
- Hay, A. and Tsiantis, M. (2009). A KNOX family TALE. *Current Opinion in Plant Biology*, 12(5), pp.593–598. doi:<https://doi.org/10.1016/j.pbi.2009.06.006>.



- Hay, A. and Tsiantis, M. (2010). KNOX genes: versatile regulators of plant development and diversity. *Development*, 137(19), pp.3153–3165. doi:<https://doi.org/10.1242/dev.030049>.
- Hiroshi Nagasaki, Sakamoto, T., Yutaka Satō and Matsuoka, M. (2001). Functional Analysis of the Conserved Domains of a Rice KNOX Homeodomain Protein, OSH15. *The Plant Cell*, 13(9), pp.2085–2098. doi:<https://doi.org/10.1105/tpc.010113>.
- Hofer, J., Gourlay, C., Michael, A. and Ellis, T.H.N. (2001). Expression of a class 1 knotted1-like homeobox gene is down-regulated in pea compound leaf primordia. *Plant Molecular Biology*, 45(4), pp.387–398. doi:<https://doi.org/10.1023/a:1010739812836>.
- Isbir, T., Kirac, D., Demircan, B. and Dalan, B. (2013). Gel Electrophoresis. *Brenner's Encyclopedia of Genetics*, pp.165–167. doi:<https://doi.org/10.1016/b978-0-12-374984-0.00580-5>.
- Jasinski, S., Piazza, P., Craft, J., Hay, A., Woolley, L., Rieu, I., Phillips, A., Hedden, P. and Tsiantis, M. (2005). KNOX Action in Arabidopsis Is Mediated by Coordinate Regulation of Cytokinin and Gibberellin Activities. *Current Biology*, 15(17), pp.1560–1565. doi:<https://doi.org/10.1016/j.cub.2005.07.023>.
- Jia, P., Zhang, C., Xing, L., Li, Y., Shah, K., Zuo, X., Zhang, D., An, N., Han, M. and Ren, X. (2020). Genome-Wide Identification of the MdKNOX Gene Family and Characterization of Its Transcriptional Regulation in Malus domestica. *Frontiers in Plant Science*, 11. doi:<https://doi.org/10.3389/fpls.2020.00128>.
- Kadri, K. (2019). Polymerase Chain Reaction (PCR): Principle and Applications. *Perspectives on Polymerase Chain Reaction*. [online] doi:<https://doi.org/10.5772/intechopen.86491>.
- Khan, N., Hu, C., Khan, W.A., Wang, W., Ke, H., Huijie, D., Zhishuo, Z. and Hou, X. (2018). Genome-wide Identification, Classification, and Expression Pattern of Homeobox Gene Family in Brassica rapa under Various Stresses. *Scientific Reports*, 8(1). doi:<https://doi.org/10.1038/s41598-018-34448-x>.
- Lee, P.Y., Costumbrado, J., Hsu, C.-Y. and Kim, Y.H. (2012). Agarose Gel Electrophoresis for the Separation of DNA Fragments. *Journal of Visualized Experiments*, [online] 3923(62). doi:<https://doi.org/10.3791/3923>.
- Lozano, E.V., Ibarra, J.E., Herrera-Ubaldo, H., de Folter, S. and Ponce, J.L.C. (2021). Osmotic stress-induced somatic embryo maturation of coffee Coffea arabica L., shoot and root apical meristems development and robustness. *Scientific Reports*, 11(1). doi:<https://doi.org/10.1038/s41598-021-88834-z>.
- Meisel, L. and Lam, E. (1996). The conserved ELK-homeodomain of KNOTTED-1 contains two regions that signal nuclear localization. *Plant Molecular Biology*, 30(1), pp.1–14. doi:<https://doi.org/10.1007/bf00017799>.
- Meng, L., Liu, X., He, C., Xu, B., Li, Y. and Hu, Y. (2020). Functional divergence and adaptive selection of KNOX gene family in plants. *Open Life Sciences*, 15(1), pp.346–363. doi:<https://doi.org/10.1515/biol-2020-0036>.
- Nobres, P., Patreze, C.M., Waltenberg, F.P., Correa, M.F., Tavano, E.C. da R., Mendes, B.M.J. and Alves-Ferreira, M. (2016). Characterization of the Promoter of the Homeobox Gene CaHB12 in Coffea arabica. *Tropical Plant Biology*, 9(1), pp.50–62. doi:<https://doi.org/10.1007/s12042-016-9159-2>.
- Oliveira, R.R. de, Cesarino, I., Mazzafera, P. and Dornelas, M.C. (2014). Flower development in Coffea arabica L.: new insights into MADS-box genes. *Plant Reproduction*, 27(2), pp.79–94. doi:<https://doi.org/10.1007/s00497-014-0242-2>.
- Pinto, R.T., Freitas, N.C., Cardoso, T.B., Paiva, L.V., de, D. and Pires, W. (2019). Genome-wide analysis, transcription factor network approach and gene expression profile of GH3 genes over early somatic embryogenesis in Coffea spp. *BMC Genomics*, 20(1). doi:<https://doi.org/10.1186/s12864-019-6176-1>.
- Rismayanti, A.Y. and Nafi'ah, H.H. (2021). Modifikasi Media Pada Induksi Kalus Kopi Arabika (Coffea Arabica L.) Berbuah Kuning. *Agro Wiralodra*, 4(2), pp.42–49.



UNIVERSITAS
GADJAH MADA

- Ruben, V., Lawrie, M.D. and Semarti, E. (2022). Isolation and Characterization of *Vanda Orchid Homeobox* Gene from *Vanda tricolor* var. *Suavis* Lindl. form Merapi. *Advances in biological sciences research*, 22. doi:<https://doi.org/10.2991/absr.k.220406.036>.
- Sakakibara, K., Ando, S., Yip, H.K., Tamada, Y., Hiwatashi, Y., Murata, T., Deguchi, H., Hasebe, M. and Bowman, J.L. (2013). KNOX2 Genes Regulate the Haploid-to-Diploid Morphological Transition in Land Plants. *Science*, 339(6123), pp.1067–1070. doi:<https://doi.org/10.1126/science.1230082>.
- Semarti, E., Purwantoro, A. and Indrianto, A. (2014). IN VITRO CULTURE OF ORCHIDS: THE ROLES OF CLASS-1 KNOX GENE IN SHOOT DEVELOPMENT. *Berkala Penelitian Hayati*, 20(1), pp.18–27. doi:<https://doi.org/10.23869/bphjbr.20.1.20144>.
- Serikawa, K.A., Martinez-Laborda, A. and Zambryski, P. (1996). Three knotted1-like homeobox genes in *Arabidopsis*. *Plant Molecular Biology*, 32(4), pp.673–683. doi:<https://doi.org/10.1007/bf00020208>.
- Setyawati, R. and Zubaidah, S. (2021). Optimasi Konsentrasi Primer dan Suhu Annealing dalam Mendeteksi Gen Leptin pada Sapi Peranakan Ongole (PO) Menggunakan Polymerase Chain Reaction (PCR). *Indonesian Journal of Laboratory*, 4(1), p.36. doi:<https://doi.org/10.22146/ijl.v4i1.65550>.
- Shara, P.N., Thomy, Z., Anhar, A., Harnelly, E. and RamLan, R. (2021). Morphological characterization of some *Coffea arabica* L. varieties in Gayo Experimental Garden Bener Meriah. *Journal of physics*, 1882(1), pp.012092–012092. doi:<https://doi.org/10.1088/1742-6596/1882/1/012092>.
- Shipunov, A., Fernández-Alonso, J.L., Hassemer, G., Alp, S., Lee, H.J. and Pay, K. (2021). Molecular and Morphological Data Improve the Classification of Plantagineae (Lamiales). *Plants*, 10(11), p.2299. doi:<https://doi.org/10.3390/plants10112299>.
- Sikand, K., Singh, J., Ebron, J.S. and Shukla, G.C. (2012). Housekeeping Gene Selection Advisory: Glyceraldehyde-3-Phosphate Dehydrogenase (GAPDH) and β-Actin Are Targets of miR-644a. *PLoS ONE*, 7(10), p.e47510. doi:<https://doi.org/10.1371/journal.pone.0047510>.
- Silalahi, D., Wirawan, I.G.P. and Sasadara, M.M.V. (2021). Optimization of annealing temperature for amplification of EhoscnOla locus in pranajawa (*Euchresta horsfieldii*) plant collected from mountains, urban and coastal areas in Bali. *IOP Conference Series: Earth and Environmental Science*, 913(1), p.012059. doi:<https://doi.org/10.1088/1755-1315/913/1/012059>.
- Speck, O. and Speck, T. (2021). Functional morphology of plants – a key to biomimetic applications. *New Phytologist*, 231(3). doi:<https://doi.org/10.1111/nph.17396>.
- Tsuda, K. and Hake, S. (2015). Diverse functions of KNOX transcription factors in the diploid body plan of plants. *Current Opinion in Plant Biology*, 27, pp.91–96. doi:<https://doi.org/10.1016/j.pbi.2015.06.015>.
- Vionita, S., Kardhinata, E.H. and Damanik, R.I. (2021). Morphology identification and description of coffee plants (*Coffea* sp) in Karo District. *International Conference on Agriculture, Environment and Food Security*, 782(4), pp.042051–042051. doi:<https://doi.org/10.1088/1755-1315/782/4/042051>.
- Vollbrecht, E., Veit, B., Sinha, N. and Hake, S. (1991). The developmental gene Knotted-1 is a member of a maize homeobox gene family. *Nature*, 350(6315), pp.241–243. doi:<https://doi.org/10.1038/350241a0>.
- Yu, H., Yang, S.H. and Goh, C.J. (2000). DOH1, a Class 1 knox Gene, Is Required for Maintenance of the Basic Plant Architecture and Floral Transition in Orchid. *The Plant Cell*, 12(11), pp.2143–2159. doi:<https://doi.org/10.1105/tpc.12.11.2143>.