

DAFTAR PUSTAKA

- Abbassy, M.A., 2015, Abushal, A. Differences In Dentofacial Characteristics Of Class I Malocclusion Between Saudi And Japanese Adult Females., *J. Orthod. Sci*, 4: 86–91
- Ahila, S. C., Sasikala, C., Kumar, B. M., Tah, R., Abinaya, K., 2016, Evaluation of the Correlation of Ramus Height, Gonial Angle, and Dental Height with Different Facial Forms in Individuals with Deep Bite Disorders, *Ann Med Health Sci Res*, 6(4): 232-8
- Al-Khateeb, S.N., 2009, Anteroposterior and Vertical Components of Class II division 1 and division 2 Malocclusion, *Angle Orthod*, 79(5): 860-2
- Alshahrani, I., Kamran, M. A., Alhaizaey, A., Abumelha, N., 2018, Evaluation of skeletal variations and establishment of Cephalometric Norms in Saudi Sub Population using Bjork Jarabak's analysis, *Pak J Med Sci*, 34(5): 1104-9
- Ardani, I.G.A.W., Sanjaya, M.L., Sjamsudin, J., 2018, Cephalometric Characteristic of Skeletal Class II Malocclusion in Javanese Population at Universitas Airlangga Dental Hospital, *Contemp Clin Dent*, 9(2): 342-5
- Baral P., 2013, Prevalence of malocclusion in permanent dentition in Aryan and Mongoloid races of Nepal-A Comparative study. *Pakistan Orthod J*, 5(2):57–9
- Basri, A., Mubarak, A., Siradjuddin, H. K., Abdullah, S.D., 2023, Penentuan Jumlah Klaster Terbaik pada K-Means dalam Melihat Pola Klastering Data Mahasiswa yang Telah Lulus, *JATI*, 3(1): 80-86
- Bratu, D. C., Balan, R., Szuhaneck, C., Pop, S., Bratu, E. A., Popa, G., 2014, Craniofacial morphology in patients with Angle Class II Division 2 malocclusion, *Rom J Morphol Embryol*, 55(3): 909-13
- Brezniak, N., Arad, A., Heller, M., Dinbar, A., Dinte, A., Wasserstein, A., 2002, Pathognomonic Cephalometric Characteristics of Angle Class II Division 2 Malocclusion, *Angle Orthod*, 72 (3): 251
- Chairani, C.N., Rahmi, E., 2016, Korelasi antara dimensi vertikal oklusi dengan panjang jari kelingking pada sub-ras Deutro Melayu, *Maj Ked Gi Ind*, 2(3): 156-8
- Currie, K., Sawchuk, D., Saltaji, H., Oh, H., Flores, C., Lagravere, M., 2017, Posterior cranial base natural growth and development: A systematic review, *Angle Orthod*, 87(6): 897-8
- Damayanti, A., Ayuningtyas, R., 2008, Karakteristik Fisik dan Pemanfaatan Pantai Karst Kabupaten Gunungkidul, *Makara Teknologi*, 12(2): 91-2

- Darwis, R., Editiawarni, T., 2018, Hubungan antara sudut interinsisial terhadap profil jaringan lunak wajah pada foto sefalometri, *J Ked GI Unpad*, 30(1): 15-9
- Dodda, K.K., Prasad, S.E., Kanuru, R.K., Nalluri, S., Mittapalli, Raghavendra, 2015, Diagnostic features of Angle's Class II div 2 malocclusion, *J Int Soc Prev Community Dent*, 5(6): 513-7
- English, J. D., Akyalcin, S., Peltomaki, T., Litschel, K., 2015, *Mosby's Orthodontic Review* Second Edition, Elsevier Inc: Missouri: 41
- Enikawati, M., Soenawan, H., Suharsini, M., Budihardjo, S.B., Sutadi H., Rizal, M.F., Fauziah, E., Wahano, N.A., Indriati, I.S., Maxillary and mandibular lengths in 10 to 16-year-old children (lateral cephalometry study), *J Phys Conf Ser*, 1073(2): 1-5
- Fitriyani, N., Ardani, I.G.A.W., Rusdiana, E., 2013, Garis estetik menurut Ricketts pada mahasiswa Fakultas Kedokteran Gigi Universitas Airlangga, *Maj Ked Gigi*, 46(2): 92-6
- Ghafari, J. G., Haddad, R. V., 2014, Cephalometric and dental analysis of Class II Division 2 reveals various subtypes of the malocclusion and the primacy of dentoalveolar components, *SeminOrthods*, 20(4): 272-86
- Gill, D. S., Naini, F. B., 2011, *Orthodontics Principles and Practice*, Blackwell Publishing, Singapore, 86
- Govinakovi, P. S., Al-Busaidi, I., Senguttuvan, V., 2017, Cephalometric Norms in an Omani Adult Population of Arab Descent, *Sultan Qaboos Univ Med J*, 18(2): 182-9
- H.S. Hwang, W.S. Kim, J.A. McNamara Jr., 2002, Ethnic differences in the soft tissue profile of Korean and European – American with normal occlusion and well balanced faces, *Angle Orthod.* 72 (1): 72–80
- Isiekwe, I., Dacosta, O., A cephalometric assessment of the nasolabial angle of an adult Nigerian population, *Nig Dent J*, 19(2): 80-3
- Jacob, 2000, *Buku Bacaan Antropologi Biologis*, Yogyakarta, Direktorat Jendral Pendidikan Tinggi, Departemen Pendidikan Nasional, 207
- Jacobson, A., 1975, The “Wits” appraisal of jaw disharmony, *Am J Orthod*, 67(2): 125
- Jain, N., Soni, D., 2021, An overview of class II division 2 malocclusion, *Int Journal of Health Sciences*, 5(S2): 214-7
- Kalha AS, Latif A, Govardhan SN., 2008, Soft-tissue cephalometric norms in a South Indian ethnic population. *Am J Orthod Dentofacial Orthop*, 133(6): 876-81
- Katyal, D., Balakrishnan, N., 2022, Evaluation of the accuracy and reliability of WebCeph – An artificial intelligence based online software, *APOS Trends Orthod*, 12(4): 261-76

- Kavitha, K., Karthik, 2012, Comparison of cephalometric norms of Caucasians and Non- Caucasians: a forensic aid in ethnic determination, *J. Forensic*, 4 (1): 53–5
- Khosravani, Fard., Fetрати, A., Rahimi, H., Asadi, E., 2010, Cephalometric characteristics of Class II Division 2 malocclusion: A comparison with normal children, *Iran J Orthod*, 5(1): 62-9
- Kumar A, Tandon P, Singh GK, Singh GP., 2019, Soft tissue growth changes from 8 to 16 years of age: A cross-sectional study. *Natl J Maxillofac Surg*, 10:161-7
- Laguhi VA, Gunawan PN, Anindita P., 2014, Gambaran maloklusi dengan menggunakan HMAR pada pasien di Rumah Sakit Gigi dan Mulut Universitas Sam Ratulangi Manado. *Jurnal e-GiGi (eG)*, 2(2): 1-7
- Lee, Kun-Feng; Tseng, Yu-Chuan; Chang, Hong-Po; and Chou, Szu-Ting, 2018, "Orthodontic Correction of Class II Division 2 Malocclusion," *Taiwan J Orthod*, 30 (3): 143-7
- Littlewood, S. J., Mitchell, L., 2013, *An introduction to orthodontics 4th ed*, New York, Oxford University Press Inc: 107-10
- Lubis, H.F., Simanjuntak, N.U., 2022, The relationship between maxillary and mandibular lengths of ethnic Bataks of chronological age 9-15 years, *Dent. J. (Maj Ked Gi)*, 55(2): 88-92
- Mageet, A. O., 2016, Classification of skeletal and dental malocclusion: Revisited, *Stoma Edu J*, 3:40
- Mahto, R. K., Kafle, D., Giri, A., Luintel, S., Karki, A., 2022, Evaluation of fully automated cephalometric measurements obtained from web-based artificial intelligence driven platform, *BMC Oral Health*, 22(132): 1-8
- Mangla, R., Singh, N., Dua, V., Padmanabhan, P., Khanna, M., 2011, Evaluation of mandibular morphology in different facial types, *Contemp Clin Dent*, 2(3): 200-6
- Maskey, S., Shrestha, R., 2019, Cephalometric Approach to Vertical Facial Height, *Orthodontic Journal of Nepal*, 9(1): 54-5
- Maurya, R.P., Sharma, V.P., Tandon, P., Nagar, A., Verma S.L., 2014, Soft-tissue characteristics of Class II division 2 malocclusion in North Indian adult population: A comparative study, *J Orthod Res*, 2: 97-104
- Mokhtar, K. I., Bakar, N. A., Ali, A. H., 2020, Genetics of malocclusion: A review, *Intl J OrofacHealth Sci*, 1(1): 4-10

- Mullane, M., Adams, R., Antoniou, C., Darendeliler, M.A., Peel, T., Vickers, D., 2020, *Class II Division 2 Malocclusion* Edition 3, Australian Society of Orthodontists, 1-4
- Novianty, S. I., Suhartono, B., 2021, Reference values for Down's cephalometric analysis in Papuans, *Maj Ked Gi Ind*, 7(1): 17-24
- Ocak, I., Soyulu, A. R., Aksu, M., 2022, Changes in Orbicularis Oris Superior and Masseter Muscle Activities After Upper Incisor Protrusion in Class II Division 2 Malocclusion: An Electromyographic Study, *Turk J Orthod*, 35(4):231-8
- Otuyemi, O., Afolabi, D., Oyewole, T., 2022, Rickett's E-line Profile Preferences among Nigerian Orthodontists, Orthodontic Trainees, and a Young Undergraduate Students' Population, *Niger J Clin Prac*, 25: 541-2
- Padharti, S. C., Vijayalakshmi, D., Apparao, H., 2019, Evaluation of Facial Height Ratios and Growth Patterns in Different Malocclusions in a Population of Dravidian Origin – A Cephalometric Study, *IOSR-JDMS*, 18(10):59-66
- Perovic, T., Blazej, Z., 2018, Male and Female Characteristics of Facial Soft Tissue Thickness in Different Orthodontic Malocclusions Evaluated by Cephalometric Radiography, *Medical Science Monitor*, 24: 3415-24
- Pitoyo, A.J., Triwahyudi, H., 2017, Dinamika Perkembangan Etnis di Indonesia dalam Konteks Persatuan Negara, *Populasi*, 25 (1): 65-6, 69
- Phulari, B. S., 2017, *Orthodontics Principles and Practice 2nd Edition*, Jaypee Brothers Medical Publisher, New Delhi, 79
- Proffit 2019, *Contemporary Orthodontics* 6th ed, Elsevier: Philadelphia, 174-6
- Quinzi, V., Paskay, L.C., D'Andrea, N., Albani, A., Monaco, A., Saccomanno, S., 2021, Evaluation of the Nasolabial Angle in Orthodontic Diagnosis: A Systematic Review, *Appl Sci*, 11 (2531): 1-18
- Sivakumar, A., Nalabothu, P., Thanh, H.N., Antonarakis, G.S., 2021, A Comparison of Craniofacial Characteristics between Two Different Adult Populations with Class II Malocclusion—A Cross- Sectional Retrospective Study. *Biology*, 10: 438
- Soemarmi, A., Indarti, E., Pujiyono, Diamantina, A., 2019, Konsep Negara Kepulauan dalam Upaya Perlindungan Wilayah Pengelolaan Perikanan Indonesia, *Jurnal Masalah-Masalah Hukum*, 48(3): 241-2
- Valle, L.F., Martin, C., Alarcon, J. A., Fernandez, J.C.P., Ortega, R., Linares, A. I., 2020, Novel Sub-Clustering of Class III Skeletal Malocclusion Phenotypes in a Southern European Population Based on Proportional Measurements, *J Clin Med*, 9 (3048): 2-10

- Topouzelis, N., Zaviriadis, A., Markoviski, E., 2011, Variation of Skeletal Cephalometric Variables in Class II Division 2 Patients with Age, *Balk J Stom*, 15: 35-40
- Uribe, L.M., Vela, K.C., Kummet, C., Dawson, D.V., Southard, T.E., 2013, Phenotypic diversity in white adults with moderate to severe Class III malocclusion, *Am J Orthod Dentofacial Orthop*, 144(1): 33-6
- Vieira, F. P., Pinzan, A., Janson, G., Fernandes, T. M. F., Sathler, R. C., Hentiques, R. P., 2014, Facial height in Japanese-Brazilian descendants with normal occlusion, *Dental Press J Orthod*, 10(5): 54-66
- Vojdani Z, Bahmanpour S, Momeni S, Vasaghi A, Yazdizadeh A, Karamifar A, Najafifar A, Setoodehmaram S, Mokhtar A., 2009, Cephalometry in 14-18 Years Old Girls and Boys of Shiraz-Iran High School. *Int. J. Morphol*, 27(1):101-4
- Xiong, X., Huang, Y., Liu, W., Wu, Y., Yi, Y., Wang, J., 2020, Distribution of Various Maxilla-Mandibular Positions and Cephalometric Comparison in Chinese Skeletal Class II Malocclusions, *J Contemp Dent Prac*, 21(8): 822-5
- Yadav, A. O., Walia, C. S., Borle, R. M., Chaoji, K. H., Rajan, R., Datarkar, A. N., 2012, Cephalometric norms for Central Indian population using Burstone and Legan analysis, *Indian J Dent Res*, 22 (1): 28-32
- Yassir, Y.A., Salman, A.R., Nabbat, S.A., The accuracy and reliability of WebCeph for cephalometric analysis, *J Taibah Univ Med Sc*, 17(1): 57-60
- Zawawi, K. H., 2012, Comparison of Wits appraisal among different ethnic groups, *J Orthod Sci*, 1(4): 88-91