



## DAFTAR PUSTAKA

- Albarakati, S.H., 2011, Soft tissue facial profile of adult Saudis, *Saudi Med J.*, 32(8): 836-42.
- Alex, A., Shashidhar, K., Kuttaappa M.N', Nayak, K.U.S., Menta, S.K., Anushree A., 2022, Comparative analysis of mandibular changes after orthodontic treatment with and without extraction of four premolars: a digital cephalometric study, *Seria Med.*, 68(2): 61-7.
- Almansob, Y.A.M., Jubari, M., Li, A.J., Liao, S.T., Mamdouh, A., Maudhah, A.A., Almansoub, H.A.M.M., Mao, J., 2019, Patient's facial soft tissue changes following the orthodontics treatment, *IOSR-JDMS.*, 18(4): 69-78.
- Ardani, I.G.A.W., Willyanti, I., Narmada, I.B., 2018, Correlation between vertical components and skeletal class II malocclusion in ethnic Javanese, *Dove Press J.*, 10(10): 297–302.
- Ardani, I.G.A., Wicaksono, A., Hamid, T., 2020, The occlusal plane inclination analysis for determining skeletal class III malocclusion diagnosis. *Clin Cosmet Investig Dent*, 12: 163-71.
- Ayoubi, A.A., Torre, D.D., Madlena, M., 2020, Craniofacial characteristics of syrian adolescents with class II division 1 malocclusion: a retrospective study. *PeerJ.*, 8: e9545.
- Baskaradoss J.K. dan Bhagavatula P., 2021, *Burt and Eklund's Dentistry, Dental Practice, and the Community*, 7th ed, Philadelphia: WB Saunders Company, pp. 208–17.
- Bhalajhi, S.I., 2015, *Orthodontics – The Art and Science*. 5th ed, New Delhi: Arya Medi Publishing, pp. 77-9, 91, 101.
- Bishara, S.E., 2006, Class II malocclusions: diagnostic and clinical considerations with and without treatment, *Semin Orthod*, 12:11-24.
- Brahmanta, A., 2017, *Monograf Gambaran Sefalometri Skeletal, Dental dan Jaringan Lunak*, Surabaya: Penerbit Kartika Mulya, Surabaya. h. 38.
- Cenzato, N., Nobili, A., Maspero, C., 2021 Prevalence of dental malocclusions in different geographical areas: scoping review, *Dent J.*, 9(10): 117-26.
- Cobourne, M.T. Dibiase, A.T., 2010, *Handbook of Orthodontics 1st ed*, Philadelphia: Elsevier, pp. 150-1.
- Cobourne, M.T., Fleming, P.S., DiBiase, A.T., Ahmad, S., 2012, *Clinical Cases in Orthodontics*, UK: Willey-Blackwell, p. 431.
- Currie, K., Sawchuk, D., Saltaji, H., Oh, H., Flores-Mir, C., Lagravere, M., 2017, posterior cranial base natural growth and development: a systematic review, *Angle Orthod*, 87(6): 897-910.
- de Frutos-Valle, L., Martín, C., Alrcón, J.A., Palma-Fernández, J.C., Ortega, R., Iglesias-Linares, A., 2020, Novel Sub-Clustering of Class III Skeletal



- Malocclusion Phenotypes in a Southern European Population Based on Proportional Measurements, *J Clin Med*, 9(9):3048.
- Devereux L., Moles D., Cunningham, S.J., McKnight, M., 2011, How important are lateral cephalometric radiographs in orthodontic treatment planning?, *Am J Orthod Dentofac Orthop*, 139(2): e175-e81.
- Elkaseh, A., Shayeb, M.A.L., Kuduruthullah, S., Elsubeih, E.S., 2021, Cephalometrics of Libyan adults, *ASJSUR*, 45(5): 1089-94.
- Esenlik, E. Sabuncuoglu, F.A., 2012, Alveolar and symphysis regions of patients with skeletal class II division 1 anomalies with different vertical growth patterns, *Eur J Dent*, 6(2): 123-32.
- Fitriyani, N., Ardani, I.G.A.W., Rusdiana, E., 2013, Garis estetik menurut Ricketts pada mahasiswa Fakultas Kedokteran Gigi Universitas Airlangga, *Dent J*, 46(2): 92-6.
- Gavrilovic, I. Gjorgova, J., 2006, Lip position in patients with class II division 1 malocclusion, *Balk J Stom*, 10(3): 183-6.
- Gill, D.S. Naini, F.B., 2011, *Orthodontics: Principle and Practice*, UK: Blackwell, pp. 29, 32, 33, 79-81, 85-6, 159, 161.
- Hajj N.E., Bassil-Nassif, N., Tauk, A., Mouhanna-Fattal, C., Bouzerhal, J.P., 2017, Maxillary and mandibular contribution to the establishment of class II malocclusion in an adult Lebanese population, *Int Orthod*, 15(4): 677-97.
- Hassan, A.H., 2011, Cephalometric characteristics of class II division 1 malocclusion in a Saudi population living in The Western Region, *Saudi Dental J*, 23(1): 23-7.
- Heasman, P., 2003, *Master Dentistry volume 2: Restorative Dentistry, Paediatric Dentistry, and Orthodontics*, Philadelphia: Elsevier, p. 232.
- Jacob, H.B., Buschang, P.H., 2014, Mandibular growth comparisons of class I and class II division 1 skeletofacial patterns, *Angle Orthod*, 84(5): 755-61.
- Jacobson, A. dan Jacobson, R.L., 2006, *Radiographic Cephalometry: From Basics to 3-D Imaging*, Michigan: Quintessence Pub, pp. 49, 206.
- Janson, G., Mendes, L.M., Junqueira, C.H.Z., Garib, D.G., 2016, Soft-tissue changes in Class II Malocclusion patients treated with extractions: a systematic review, *Eur J Orthod*, 38(6):631-7.
- Kamak, H. Celikoglu, M., 2012, Facial soft tissue thickness among skeletal malocclusions: is there a difference?, *Korean J Orthod*, 42(1): 23-31.
- Kharbanda, O.P., 2019, *Orthodontics: Diagnosis and Management of Malocclusion and Dentofacial Deformities*, Philadelphia: Elsevier, pp.271-4.
- Latorre, F.V., Julia, C.B., Chang, R.H., Manriquez, G., Munoz, A.D., 2022, Morphological distribution of class II malocclusions according to skeletal pattern in an adult sample of The School of Dentistry of the University of Chile, *Odontoestomat*, 24(1):1-10.



- Lee, Y.J., Park, J.T., Cha, J.Y., 2015, Perioral soft tissue evaluation of skeletal class II division 1 : a lateral cephalometric study, *Am J Orthod Dentofac Orthop*, 148(3): 405-13.
- Leonardi, R., Giordano, D., Maiorana, F., Spampinato, C., 2008, Automatic cephalometric analysis, *Angle Orthod*, 78(1):145-51.
- Lin, N.H., Soemantri, E.S., Gayatri, G., 2019, Changes in soft tissue facial profile of class II skeletal malocclusion patients with retrognathic mandible treated with twin block appliances, *PJD*, 31(1): 32-7.
- Lombardo, G., Vena, F., Negri, P., Pagano, S., Barilotti, C., Paglia, L., Colombo, S., Orso, M., Cianetti, S., 2020, Worldwide prevalence of malocclusion in the different stages of dentition: a systematic review and meta-analysis, *Eur J Paediatr Dent*, 21(2): 115-22.
- Lubis, H.F., 2014, Hubungan lebar mesiodistal gigi dengan kecembunginan profil jaringan lunak wajah pada mahasiswa Fakultas Kedokteran Gigi Universitas Sumatera Utara ras deutromelayu, *Dentika*, 18(1):58-62.
- Maan, A.S., Patil, A.K., 2019, Comparison of reliability and efficiency of Down's and Steiner's cephalometric analysis between digital and conventional methods, *Saudi J Oral Dent Res*, 4(3): 109-15.
- Maurya, R.P., Sharma, V.P., Tandon, P., Naar, A., Verma, S.L., 2014, Soft-tissue characteristics of class-II division-1 malocclusion in North Indian adult population: a cephalometric study, *J Orthod Res*, 2(2): 60-7.
- McNamara J.A., Jr., 1981, Components of class II malocclusion in children 8–10 years of age, *Angle Orthod*, 51(3): 177–202.
- Mitchell, L., 2007, *An introduction to orthodontics*, 3rd ed, New York: Oxford University Press, pp. 8-9.
- Mohammad, H.A., Abu Hassan, M.I., Hussain, S.F., 2011, Cephalometric evaluation for malaysian malay by steiner analysis, *Sci Res Essays*, 6(3):627-34.
- Oliver, B.M., 1982, The influence of lip thickness and strain on upper lip response to incisor retraction, *Am. J. Orthod*, 82(2): 141-8.
- Pancherz H., Zieber K., Hoyer B., 1997, Cephalometric characteristics of Class II division 1 and Class II division 2 malocclusions: a comparative study in children, *Angle Orthod*, 67(2): 111–20.
- Perovic, T. dan Blazej, Z., 2018, Male and female FSTT in orthodontic malocclusion using cephalometric radiography, *Med Sci Monit*, 24: 3415-24.
- Perillo L., Padricelli G., Isola G., Femiano F., Chiodini P., Matarese G., 2012, Class II malocclusion division 1: a new classification method by cephalometric analysis, *Eur J Paediatr Dent*, 13(3):192-6.
- Prince, S.T.T., Dilip, S., Sangeetha, D., Ravi, K., Krishnaraj, R., 2023, Reproducibility of linear and angular cephalometric measurements obtained



- by an artificial-intelligence assisted software (WebCeph) in comparison with digital software (AutoCEPH) and manual tracing method, *Dental Press J Orthod*, 28(1):e2321214.
- Proffit, W.R., Fields, H.W., Larson, B.E., Sarver, D.M., 2019, *Contemporary Orthodontics*, 6th ed. Elsevier, Philadelphia, pp. 4, 156, 157, 174-7.
- Rosenblum R.E., 1995, Class II malocclusion: mandibular retrusion or maxillary protrusion?, *Angle Orthod*, 65(1): 49–62.
- Sidlauskas, A., 2005, The effects of the Twin-block appliance treatment on the skeletal and dentoalveolar changes in Class II Division 1 malocclusion, *Medicina (Kaunas)*, 41(5): 392–400.
- Sidlauskas, A., Svalkauskiene, V., Sidlauskas, M., 2006, Assesment of skeletal and dental pattern of class II division 1 malocclusion with relevance to clinical practice, *Stomatol*, 8(1): 3-8.
- Siriwat, P.P dan Jarabak, J.R., 1985, Malocclusion and facial morphology is there a relationship?, *Angle Orthod*, 55(2): 127-38.
- Sivakumar, A., Nalabothu, P., Thanh, H.N., Antonarakis, G.S., 2021, Comparison of craniofacial characteristics between two different adult populations with class II malocclusion – a cross-sectional retrospective study, *Biologi*, 10(5): 438-47.
- Staley, R.N. Reske, N.T., 2011, *Essentials of Orthodontics Diagnosis and Treatment*, Iowa: Blackwell Publishing, p. 7.
- Thiesen, G., Pletsch, G., Zastrow, M.D., do Valle, C.V.M., do Valle-Corotti, K.M., Patel, M.P., Conti, P.C.R., 2013, Comparative analysis of the anterior and posterior length and deflection angle of the cranial base, in individuals with facial pattern I, II and III, *Dental Press J Orthod*, 18(1): 69-75.
- Uysal, T., Yagci, A., Aldrees, A.M., Ekizer, E., 2011, Ethnic differences in dentofacial relationships of Turkish and Saudi young adults with normal occlusions and well-balanced faces, *Eur. J. Orthod.*, 34(2012): 296-301.
- Widyanti, A., Susanti, L., Sutalaksana, I.Z., Muslim, K., 2015, Ethnic differences in Indonesian anthropometry data: evidence from three different largest ethnics, *J Ergon*, 47:72-8.
- Xie, L., Wang, P., Wu, J., 2020, Soft and hard tissue changes following treatment of class II division 1 malocclusion with twin-block and myofunctional appliance: A Pilot Study, *CJPRS*, 2(4): 217-27.
- Xiong, X., Huang, Y., Liu, W., Wu, Y., Yi, Y., Wang, J., 2020, Distribution of various maxilla-mandibular positions and cephalometric comparison in chinese skeletal class II malocclusions, *J Contemp Dent Pract*, 21(8): 822-8.
- Yassir, Y.A., Salman, A.R., Nabbat, S.A., 2022, The accuracy and reliability of WebCeph for cephalometric analysis. J. Taibah Univ, *Medical Sci*, 17(1): 57-66.