

DAFTAR PUSTAKA

- Abbassy, M.A., Abushal, A. (2015) Differences in dentofacial characteristics of class I malocclusion between Saudi and Japanese adult females. *J Orthod Sci.*, 4: 86–91.
- Alhammadi, M.S., Halboub, E., Fayed, M.S., Labib, A., El-Saaidi, C.M.S. (2018) Global distribution of malocclusion traits: A systematic review, *Dental Press J Orthod.*, 23(6): e1–e10.
- Alhammadi, M.S. (2019) Dentoalveolar compensation in different anterioposterior and vertical skeletal malocclusions, *J Clin Exp Dent.*, 11(8): e745–e53.
- Alshahrani, I., Kamran, M.A., Alhaizaey, A., Abumelha, N., (2018) Evaluation of skeletal variations and establishment of cephalometric norms in saudi sub population using bjork jarabak's analysis, Pak. *J Med Sci.*, 34(5): 1104–9.
- Ananta, A., Arifin, E.N., Bakhtiar, 2005, Ethnicity and ageing in Indonesia 2000–2050, *Asian Popul.*, 1(2): 227–43.
- Ardani, I.G.A.W., Sanjaya, M.L., Sjamsudin, J. (2018) Cephalometric characteristic of skeletal class II malocclusion in Javanese population at Universitas Airlangga Dental Hospital, *Contemp Clin Dent.*, 9: S342-6.
- Ardani, I.G.A.W., Wicaksono, A. Hamid, T. (2020) The occlusal plane inclination analysis for determining skeletal class III malocclusion diagnosis, *Clin Cosmet Investig Dent.*, 12, pp. 163–71.
- Arifin, E.N., Ananta E., Utami, D.R.W.W., Handayani, NB., Pramono A. (2012) Quantifying Indonesia's Ethnic Diversity, *Asian Popul.*, 11(3): 233–56.
- Auconi, P., Scazzocchio, M., Defraia, E., Mcnamara, J.A., Franchi, L. (2014) prediction of class III treatment outcomes through orthodontic data mining, *Eur J Orthod.*, 37(3): 257–67.
- Auconi, Pietro *et al.* (2014) Forecasting craniofacial growth in individuals with class III malocclusion by computational modelling, *Eur. J. Orthod.*, 36(2): 207–16.
- Bahrou S, Hasan AA, Khalil F. (2018) Facial Proportion In Different Mandibular Rotation In Class I Individual. *IAJD.*, 5(1): 14-7.
- Bhalajhi (2018) *Orthodontics: The Art and Science*, Arya Medi Publishing House Pvt: New Delhi.

- Budipramana, M., Budhy, T.I. Ardani, I.G.A.W. (2021) Gonial angle characteristics of class iii malocclusion in javanese ethnic, *Pesqui Bras Odontopediatria Clin Integr.*, 21: 1-8.
- Cárdenas, Y.A.R., Guillen, L.E.A., Tume, Z.C.D., Mora, G.A.R., Castillo, A.A.D. (2017) Influence of the components of Björk-Jarabak cephalometric analysis on the facial profile, *IJO.*, 28 (4): 35-8.
- Chairani, C.N, Rahmi, E. (2016) Korelasi antara dimensi vertikal oklusi dengan panjang jari kelingking pada sub-ras Deutro Melayu, *MajKedGIl.*, 2(3): 155-63.
- Choi, H.J., Kim, J.Y., Yoo, S.E., Kwon, J.H., Park, K. (2010) Cephalometris characteristics of Korean children with class III malocclusion in the deciduous dentition, *Angle Orthod.*, 80 (1): 86-90.
- Fatani, N.H., Hammam, M.B., Oraif, H., Taher, S., Taj, W., Bukhari, O. (2019) Prevalence of malocclusion among Schoolchildren in Makkah, Saudi Arabia, *Open Access Maced. J Med Sci.*, 7 (5): 856–61.
- Farooq, M.U., Khan, M.A., Imran, S., Sameera, A., Qureshi, A., Ahmed, S.A., Kumar, S., Rahman, M.A.U, (2016) Assessing the reliability of digitalized cephalometric analysis in comparison with manual cephalometric analysis, *J Clin Diagnostic Res.*, 10(10): ZC20– ZC3.
- de Frutos-Valle, L., Martin, C., Alrcon, J.A., Palma-Fernandez, J.C., Ortega, R., Iglesias-Linare, A.I., (2019) Subclustering in skeletal class III phenotypes of different ethnic origins: A Systematic review, *J Evid Base Dent Pract.*, 19 (1): 34–52.
- de Frutos-Valle, L., Martin, C., Alrcon, J.A., Palma-Fernandez, J.C., Ortega, R., Iglesias-Linare, A.I., (2020) Novel sub-clustering of class III skeletal malocclusion phenotypes in a southern european population based on proportional measurements, *J Clin Med.*, 9, 3048: 1–13.
- de Frutos-Valle, L., Martin, C., Alarcon, J.A., Palma-Fernandez, J.C., (2020) Sub-clustering in skeletal class III malocclusion phenotypes via principal component analysis in a southern European population, *Sci Rep.*, 10 (1): 1–12.
- Garg, R., Alexander, M. (2015) Are we similar to Caucasians: orthognathic surgery for North Indians. *J. Maxillofac Oral Surg.*, 14 (2): 271-7.
- Goyal, V., Kapoor, D.N., Kumar, S., Sagar, M. (2011) Maturation of permanent teeth in different facial types: A comparative study, *Indian J Dent Res.*, 22(5): 627–32.

- Grabner LW., Vanarsdall RL., Katherine W.L., Huang G.J. (2013) *Orthodontic current principles and techniques*, 6th ed, America: Elsevier.
- Hardy, D. K., Cubas, Y. P., Orellana, M. F. (2012) Prevalence of Angle class III malocclusion: A systematic review and meta-analysis, *Open J Epidemiol.*, 2 (04): 75–82.
- Grewal, H., Sharma, H., Aggarwal, N., (2013) A cephalometric comparison of horizontal and vertical skeletal parameters in North and South Indian population groups, *JPFA.*, 27 (1): 14-17.
- Holdaway RA. (1983) A soft-tissue cephalometric analysis and its use in orthodontic treatment planning. Part I. *Am J Orthod.*, 84(1):1-28.
- Jacobson, A. (2007) Radiograph cephalometry, *J Chem Inf Model.*, 53(9): 1689–1699.
- Jeelani W., Fida M., Shaikh A. (2015) Facial soft tissue thickness among various vertical facial patterns in adult Pakistani subjects, *Forensic Sci Int.*, 257: 517e1-517.e6
- Juneja, M., Garg, P., Kaur, R., Manocha, P., Prateek, Batra, S., Singh, P., Singh, S., Jindal P. (2021) A review on cephalometric landmark detection techniques, *BSPC.*, 66: 102486: 1-14.
- Kale, B. Buyukcavus, M. (2020) Comparison of three-dimensional soft tissue evaluations between skeletal and pseudo class III malocclusions, *Sci Rep.*, 10(14717): 1–11.
- Katyal, D. Balakrishnan, N. (2022) Evaluation of the accuracy and reliability of WebCeph – An artificial intelligence-based online software, *APOS Trends Orthod.*, 12 (4): 271-6.
- Kavadia S, Sidiropoulou-Chatzigianni S, Pappa G, Markovitsi El, Kalklamanos El. (2017) Soft Tissue Characteristics and Gender Dimorphism in Class III Malocclusion: Cephalometric Study in Adults Greeks. *Balk J Dent Med.*, 21:162-6.
- Kolokitha, O.E., Georgiadis, T. (2019) Differential diagnosis of Skeletal Class III, *Balk J Dent Med.*, 23(3): 55–62.
- Komalawati., Indriaty, E., Supartinah, A. (2013) Profil jaringan lunak dan keras wajah lelaki dan perempuan dewasa etnis Aceh berdasarkan keturunan campuran Arab, Cina, dan Hindia, *Cakradoanya Dent J.*, 5(2): 542-618.
- Krull, J.T., Krull, G.E., Dean, J.A. (2016) Cephalometrics and Facial Aesthetics: The Key to Complete Treatment Planning, in *McDonald and Avery's Dentistry for the Child and Adolescent*: Tenth Edition. Elsevier Inc.

- Larasasu, A., Koesbardiati, T., Yudianto, A. (2018) Estimasi Tinggi Badan Ras Mongoloid, *BIOPASCA*, 20(2): 107–20.
- Lazi, H., Efendi, R., Purwandari, E. P. (2017) Deteksi warna kulit menggunakan model warna Cielab Neural Network untuk identifikasi ras manusia (studi kasus ras: Kaukasoid, Mongoloid, dan Negroid), *Rekursif*, 5(2): 121–33.
- Li, C., Cai, Y., Chen, S., Chen, F. (2016) Classification and characterization of class III malocclusion in Chinese individuals, *Head & Face Med.*, 12(1): 1–8.
- Lubis, MM., Fulvian, J., 2021, Perbedaan tinggi vertikal wajah pada maloklusi kelas I dan II skeletal, *PJDRS*, 5(1): 51-6.
- Mahto, R.K., Kafle, D., Giri, A., Luintel S., Karki, A. (2022) Evaluation of fully automated cephalometric measurements obtained from web-based artificial intelligence driven platform, *BMC Oral Health*, 22(132): 1-8.
- Munandar, S., (1995) Cephalometric analysis of Deutro-Malay Indonesians, *Aust. Dent. J.*, 40 (6): 381-8.
- Nanda, R.S., Tosun, Y.S. (2019) *Biomechanis in Orthodontics, Principle, and Practice*. Chicago: Quintessence Publishing Co.
- de Oliveira, T.C.P., Copello F.M., Silva I.M.C, Nojima L.I, Nojima M.C.G. (2021) Dentofacial and skeletal pattern in African descendants from Southeastern Brazil: Clinical prospective study, *Dental Press J Orthod.*, 26(3): 1–25.
- Proffit, W.R., Fields, H. W., Sarver, D. M. (2019) *Contemporary Orthodontics*, 6th edition, St. Louis Missouri: Mosby Elsevier.
- Phulari, B.S. (2013) *An Atlas on Cephalometric LandMarks*, Jaypee Brothers Medical Publishers: New Delhi.
- Phulari, B.S., (2017) *Orthodontics principles and practice*, 2nd Edition. Jaypee Brothers Medical Publishers: New Delhi.
- Rahoojo A., Naz, S., Rahoojo, M.A., Memon, AB., Sandeep, Rahoojo, M.A., (2022) Soft Tissue Lip Profile with Steiner(S) and Ricketts (E) Lip Analysis in Patient's with Skeletal and Dental Class I Occlusion, *P J M H S*, 16 (1): 786-788.
- Rathore, AS, Dhar V, Arora R, Diwanji A. (2012) Cephalometric Norms for Mewari Children using Steiner's Analysis, *Int J Clin Pediatr Dent.*, 5(3): 173–7.
- Rubika, J., Felicita, A.S. and Sivambiga, V. (2015) Gonial angle as an indicator for the prediction of growth pattern, *World J. Dent.*, 6(3): 161–163.

- Sanggarnjanavanich S, Sekiya T, Nomura Y, Nakayama T, Hanada N, Nakamura Y. (2014) Cranial-base morphology in adults with skeletal Class III malocclusion. *Am. J. Orthod. Dentofacial Orthop.*, 146: 82-91.
- Shindy, R. A., Sahelangi, O. P. (2020) Gambaran hasil analisis sefalometri pada pasien ras Deutro Melayu usia 8-12 tahun menggunakan analisis Rickett, *J Kedokteran Gigi*, 2: 19-22.
- Siddika A., Rahman S, A., Alam M, K. (2020) Ricketts' cephalometric analysis for Saudi population. *Pesqui Bras Odontopediatria Clín Integr.*, 20: e5364.
- Uribe, L.M.M., Vela, K.C., Kummet, C., Dawson D.V., Southarf, T.E. (2013) Phenotypic diversity in white adults with moderate to severe Class III malocclusion, *Am J Orthod Dentofacial Orthop.*, 144(1): 32-42.
- Vasconcelos, M.B., Vercelino, C.R.M.P., Gurgel, J.dA., Bramante, F.dS. (2014) Cephalometric characteristics of Class III malocclusion In Brazilian individuals, *Braz J Oral Sci.*, 13(4): 314-8.
- Vieira FP, Pinzan A, Janson G, Fernandes TMF, Sathler RC, Henriques RP. (2014) Facial height in Japanese-Brazilian descendants with normal occlusion. *Dent Press J Orthod.*, 19(5):54-66.
- Wu, TY., Chang, TF., Wu, CH. (2022) True vertical changes in patients with skeletal class III malocclusion after nonsurgical orthodontic treatment—a retrospective study comparing different vertical facial patterns, *J Dent Sci.*, 17 (3): 1096-1101.
- Yang, I., Choi, J. Baek, S. (2022) Characterization of phenotype of skeletal class III Malocclusion in Korean Adult Patients Treated with Orthognatic Surgery using Cluster Analysis, *Angle Orthod.*, 92(4): 537-46.
- Yassir, Y.A., Salman, A.R., Nabbat, S.A. (2022) The accuracy and reliability of WebCeph for cephalometric analysis, *J. Taibah Univ. Med. Sci.*, 17(1): 57-66.
- Zawawi, K.H., Alsulaimani, F.F., Al-Dharrab, A.A., Afify, A.R., Al-Zahrani, M.S., Baeshen, H.A. (2021) Morphological features of Class I, II and III malocclusions of Saudi adolescents, *Saudi J. Biol. Sci.*, 28(6): 3534-3539.
- Zere, E., Chaudari, P.K., Sharan, J., Dhingra, K., Tiwari, N. (2018) Developing class III malocclusions: challenges and solutions, *Clin. Cosmet. Investig. Dent.*, 10: 99-116.