

REFERENCES

- Aguilera-Pena, M. P., Cardenas-Cruz, A. F., Baracaldo, I., Garcia-Cifuentes, E., Ocampo-Navia, M. I., & Coral, E. J. (2021). Ischemic stroke in young adults in Bogota, Colombia: a cross-sectional study. *Neurological Sciences*, 42(2), 639-645. <https://doi.org/10.1007/s10072-020-04584-2>
- Aroor, S., Singh, R., & Goldstein, L. B. (2017). BE-FAST (Balance, Eyes, Face, Arm, Speech, Time): Reducing the Proportion of Strokes Missed Using the FAST Mnemonic. *Stroke*, 48(2), 479–481. <https://doi.org/10.1161/STROKEAHA.116.015169>
- Billinger, S. A., Arena, R., Bernhardt, J., Eng, J. J., Franklin, B. A., Johnson, C. M., ... & Council on Clinical Cardiology (2014). Physical activity and exercise recommendations for stroke survivors: a statement for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke*, 45(8), 2532–2553. <https://doi.org/10.1161/STR.0000000000000022>
- Billinger, S. A., Boyne, P., Coughenour, E., Dunning, K., & Matlage, A. (2015). Does aerobic exercise and the FITT principle fit into stroke recovery?. *Current neurology and neuroscience reports*, 15(2), 519. <https://doi.org/10.1007/s11910-014-0519-8>
- Birawa, A. B. P. and Amalia, L. (2015). Stroke pada Usia Muda. *cdkjournal*, 42(10), 736-739. <http://dx.doi.org/10.55175/cdk.v42i10.955>.
- Boot, E., Ekker, M. S., Putaala, J., Kittner, S., De Leeuw, F. E., & Tuladhar, A. M. (2020). Ischaemic stroke in young adults: a global perspective. *Journal of neurology, neurosurgery, and psychiatry*, 91(4), 411–417. <https://doi.org/10.1136/jnnp-2019-322424>
- Brott, T., Adams, H. P., Jr, Olinger, C. P., Marler, J. R., Barsan, W. G., Biller, J., Spilker, J., Holleran, R., Eberle, R., & Hertzberg, V. (1989). Measurements of acute cerebral infarction: a clinical examination scale. *Stroke*, 20(7), 864–870. <https://doi.org/10.1161/01.str.20.7.864>
- Bursac, Z., Gauss, C. H., Williams, D. K., & Hosmer, D. W. (2008). Purposeful selection of variables in logistic regression. *Source code for biology and medicine*, 3(17), 1-8. <https://doi.org/10.1186/1751-0473-3-17>
- Chen, R., Ovbiagele, B., & Feng, W. (2016). Diabetes and Stroke: Epidemiology, Pathophysiology, Pharmaceuticals and Outcomes. *The American journal of the medical sciences*, 351(4), 380–386. <https://doi.org/10.1016/j.amjms.2016.01.011>
- Choudhury, M. J. H., Chowdhury, M. T. I., Nayeem, A., & Jahan, W. A. (2015). Modifiable and Non-Modifiable Risk Factors of Stroke: A Review Update. *Journal of National Institute of*



- Neurosciences Bangladesh*, 1(1), 22–26. <https://doi.org/10.3329/jninb.v1i1.22944>
- Dahlberg, K., Stenberg, E., Liang, Y., Nilsson, U., & Jaensson, M. (2022). The General Self-Efficacy Scale in a population planned for bariatric surgery in Sweden: a psychometric evaluation study. *BMJ open*, 12(11), e061509. <https://doi.org/10.1136/bmjopen-2022-061509>
- Ding, M., Wang, R., Kalpouzos, G., Laukka, E. J., Li, Y., Johnell, K., Fratiglioni, L., & Qiu, C. (2021). Cerebral Small Vessel Disease Associated With Atrial Fibrillation Among Older Adults: A Population-Based Study. *Stroke*, 52(8), 2685–2689. <https://doi.org/10.1161/STROKEAHA.120.031573>
- Divani, A. A., Majidi, S., Barrett, A. M., Noorbaloochi, S., & Luft, A. R. (2011). Consequences of stroke in community-dwelling elderly: the health and retirement study, 1998 to 2008. *Stroke*, 42(7), 1821–1825. <https://doi.org/10.1161/STROKEAHA.110.607630>
- Djukanovic, I., Carlsson, J., & Årestedt, K. (2017). Is the Hospital Anxiety and Depression Scale (HADS) a valid measure in a general population 65-80 years old? A psychometric evaluation study. *Health and quality of life outcomes*, 15(1), 1-10. <https://doi.org/10.1186/s12955-017-0759-9>
- Duvekot, M. H. C., Venema, E., Rozeman, A. D., Moudrous, W., Vermeij, F. H., ... PRESTO investigators (2021). Comparison of eight prehospital stroke scales to detect intracranial large-vessel occlusion in suspected stroke (PRESTO): a prospective observational study. *The Lancet. Neurology*, 20(3), 213–221. [https://doi.org/10.1016/S1474-4422\(20\)30439-7](https://doi.org/10.1016/S1474-4422(20)30439-7)
- Ezema, C. I., Akusoba, P. C., Nweke, M. C., Uchewoke, C. U., Agono, J., & Usoro, G. (2019). Influence of Post-Stroke Depression on Functional Independence in Activities of Daily Living. *Ethiopian journal of health sciences*, 29(1), 841–846. <https://doi.org/10.4314/ejhs.v29i1.5>
- GBD 2019 Stroke Collaborators (2021). Global, regional, and national burden of stroke and its risk factors, 1990-2019: a systematic analysis for the Global Burden of Disease Study 2019. *The Lancet. Neurology*, 20(10), 795–820. [https://doi.org/10.1016/S1474-4422\(21\)00252-0](https://doi.org/10.1016/S1474-4422(21)00252-0)
- Murie-Fernández, M., & Marzo, M. M. (2020). Predictors of Neurological and Functional Recovery in Patients with Moderate to Severe Ischemic Stroke: The EPICA Study. *Stroke research and treatment*, 2020, 1419720. <https://doi.org/10.1155/2020/1419720>
- Frost, Y., Weingarden, H., Zeilig, G., Nota, A., & Rand, D. (2015). Self-Care Self-Efficacy Correlates with Independence in Basic Activities of Daily Living in Individuals with Chronic Stroke. *Journal of stroke and cerebrovascular diseases : the official journal of National Stroke Association*, 24(7), 1649–1655.



<https://doi.org/10.1016/j.jstrokecerebrovasdis.2015.03.054>

- Gyawali, P., Chow, W. Z., Hinwood, M., Kluge, M., English, C., Ong, L. K., Nilsson, M., & Walker, F. R. (2020). Opposing Associations of Stress and Resilience With Functional Outcomes in Stroke Survivors in the Chronic Phase of Stroke: A Cross-Sectional Study. *Frontiers in neurology*, *11*, 230. <https://doi.org/10.3389/fneur.2020.00230>
- Mickey, R. M., & Greenland, S. (1989). The impact of confounder selection criteria on effect estimation. *American journal of epidemiology*, *129*(1), 125–137. <https://doi.org/10.1093/oxfordjournals.aje.a115101>
- Haast, R. A., Gustafson, D. R., & Kiliaan, A. J. (2012). Sex differences in stroke. *Journal of cerebral blood flow and metabolism : official journal of the International Society of Cerebral Blood Flow and Metabolism*, *32*(12), 2100–2107. <https://doi.org/10.1038/jcbfm.2012.141>
- Harvey R. L. (2015). Predictors of Functional Outcome Following Stroke. *Physical medicine and rehabilitation clinics of North America*, *26*(4), 583–598. <https://doi.org/10.1016/j.pmr.2015.07.002>
- Hasanah, M., Gofir, A., & Setyopranoto, I. (2019) ‘Neurorehabilitasi motorik pasca stroke’, *Berkala NeuroSains*, *18*(2), 51–56. <https://journal.ugm.ac.id/bns/article/view/54998>.
- Haselbach, D., Renggli, A., Carda, S., & Croquelois, A. (2014). Determinants of neurological functional recovery potential after stroke in young adults. *Cerebrovascular diseases extra*, *4*(1), 77–83. <https://doi.org/10.1159/000360218>
- Hidayati, N. and Irmawati, F. (2020) *Anatomi Fisiologi Manusia Sistem Kardiovaskular Berbasis SETS-PBL (Science, Enviromental, Technology, Society-Problem Based Learning)*. I. Edited by T. A. Prayitno. Malang: Media Nusa Kreatif.
- Huang, H. C., Chang, C. H., Lee, T. H., Chang, Y. J., Ryu, S. J., Chang, T. Y., Huang, K. L., Liu, C. H., & Chang, H. J. (2013). Differential trajectory of functional recovery and determinants for first time stroke survivors by using a LCGA approach: a hospital based analysis over a 1-year period. *European journal of physical and rehabilitation medicine*, *49*(4), 463–472.
- Hui, C., Tadi, P., & Patti, L. (2022). Ischemic Stroke. In *StatPearls*. StatPearls Publishing.
- Jang, D. E., & Shin, J. H. (2019). Self-Care Performance of Middle-Aged Stroke Patients in Korea. *Clinical nursing research*, *28*(3), 263–279. <https://doi.org/10.1177/1054773817740670>
- Jang, S. Y., Shin, Y., Kim, D. Y., & Sohn, M. K. (2015). Effect of obesity on functional outcomes at 6 months post-stroke among elderly Koreans: A prospective multicentre study. *BMJ Open*, *5*(12), 1–6. <https://doi.org/10.1136/bmjopen-2015-008712>.



- Katan, M., & Luft, A. (2018). Global Burden of Stroke. *Seminars in neurology*, 38(2), 208–211.
<https://doi.org/10.1055/s-0038-1649503>
- Kemenkes RI. (2013). Pedoman Pengendalian Stroke. *Pedoman Pengendalian Stroke*, 1–66.
<http://p2ptm.kemkes.go.id/uploads/2016/10/Pedoman-Pengendalian-Stroke.pdf>
- Kemenkes RI. (2014). Situasi kesehatan jantung. *Pusat data dan informasi kementerian kesehatan RI*, 3. <https://doi.org/10.1017/CBO9781107415324.004>.
- Kemenkes RI. (2018). Stroke Dont Be The One. 10. <https://p2ptm.kemkes.go.id/kegiatan-p2ptm/pusat/-/begini-cara-mengenali-gejala-stroke>
- Kernan, W. N., Inzucchi, S. E., Sawan, C., Macko, R. F., & Furie, K. L. (2013). Obesity: a stubbornly obvious target for stroke prevention. *Stroke*, 44(1), 278–286.
<https://doi.org/10.1161/STROKEAHA.111.639922>
- Knoflach, M., Matosevic, B., Rücker, M., Furtner, M., Mair, A., Wille, G., Zangerle, A., Werner, P., Ferrari, J., Schmidauer, C., Seyfang, L., Kiechl, S., Willeit, J., & Austrian Stroke Unit Registry Collaborators (2012). Functional recovery after ischemic stroke--a matter of age: data from the Austrian Stroke Unit Registry. *Neurology*, 78(4), 279–285.
<https://doi.org/10.1212/WNL.0b013e31824367ab>
- Kobyłańska, M., Kowalska, J., Neustein, J., Mazurek, J., Wójcik, B., Belza, M., Cichosz, M., & Szczepańska-Gieracha, J. (2018). The role of biopsychosocial factors in the rehabilitation process of individuals with a stroke. *Work (Reading, Mass.)*, 61(4), 523–535.
<https://doi.org/10.3233/WOR-162823>
- Kossi, O., Batcho, C. S., Adoukonou, T., & Thonnard, J. L. (2016). Functional recovery after stroke in Benin: A six-month follow-up study. *Journal of rehabilitation medicine*, 48(8), 671–675.
<https://doi.org/10.2340/16501977-2128>
- Külzer, A. M., Scolari, C. C., & Gus, M. (2008). Relationship between usual physical, cognitive and social activities and functional recovery at hospital discharge after acute stroke. *Journal of rehabilitation medicine*, 40(3), 195–199. <https://doi.org/10.2340/16501977-0149>
- Kusyani, A. and Khayudin, B. A. (2022). *Asuhan Keperawatan Stroke untuk Mahasiswa dan Perawat Profesional*. Edited by Guepedia/Br. Bogor: Guepedia.
- Lewin, A., Jöbges, M., & Werheid, K. (2013). The influence of self-efficacy, pre-stroke depression and perceived social support on self-reported depressive symptoms during stroke rehabilitation. *Neuropsychological rehabilitation*, 23(4), 546–562.
<https://doi.org/10.1080/09602011.2013.794742>
- Lee, K. B., Lim, S. H., Kim, K. H., Kim, K. J., Kim, Y. R., Chang, W. N., Yeom, J. W., Kim, Y. D.,

- & Hwang, B. Y. (2015). Six-month functional recovery of stroke patients: a multi-time-point study. *International journal of rehabilitation research. Internationale Zeitschrift fur Rehabilitationsforschung. Revue internationale de recherches de readaptation*, 38(2), 173–180. <https://doi.org/10.1097/MRR.0000000000000108>
- Sheffler, L. R., Knutson, J. S., Gunzler, D., & Chae, J. (2012). Relationship between body mass index and rehabilitation outcomes in chronic stroke. *American journal of physical medicine & rehabilitation*, 91(11), 951–956. <https://doi.org/10.1097/PHM.0b013e31826458c6>
- Murie-Fernández, M., & Marzo, M. M. (2020). Predictors of Neurological and Functional Recovery in Patients with Moderate to Severe Ischemic Stroke: The EPICA Study. *Stroke research and treatment*, 2020, 1419720. <https://doi.org/10.1155/2020/1419720>
- Novrianto, R., Maretih, A. K. E. and Wahyudi, H. (2019). Validitas Konstruk Instrumen General Self Efficacy Scale Versi Indonesia. *Jurnal Psikologi*, 15(1), 1-9. <https://doi.org/10.24014/jp.v15i1.6943>.
- Ohyama, K., Watanabe, M., Nosaki, Y., Hara, T., Iwai, K., & Mokuno, K. (2020). Correlation Between Skeletal Muscle Mass Deficit and Poor Functional Outcome in Patients with Acute Ischemic Stroke. *Journal of stroke and cerebrovascular diseases : the official journal of National Stroke Association*, 29(4), 104623. <https://doi.org/10.1016/j.jstrokecerebrovasdis.2019.104623>
- Paciaroni, M., & Acciarresi, M. (2019). Poststroke Fatigue. *Stroke*, 50(7), 1927–1933. <https://doi.org/10.1161/STROKEAHA.119.023552>
- Parmar, P. (2018). Stroke: Classification and diagnosis. *Clinical Pharmacist*, 10(1). <https://doi.org/10.1211/CP.2018.20204150>.
- Pongantung, H., Lorica, J., Rosdewi, & Solon, M. (2020). Construct validity indonesian version of barthel index for post stroke. *Indian Journal of Public Health Research and Development*, 11(3), 2007–2011. <https://www.embase.com/search/results?subaction=viewrecord&id=L2004452298&from=export>.
- Puolakka, T., Virtanen, P., Kinnunen, J., Kuisma, M., & Strbian, D. (2021). Prehospital identification of large vessel occlusion using the FAST-ED score. *Acta neurologica Scandinavica*, 144(4), 400–407. <https://doi.org/10.1111/ane.13474>
- Putra, M. D. K., Rahayu, W. and Umar, J. (2019). Indonesian-language version of general self-efficacy scale-12 using Bayesian confirmatory factor analysis: A construct validity testing. *Jurnal Penelitian dan Evaluasi Pendidikan*, 23(1), 12–25.



<https://doi.org/10.21831/pep.v23i1.20008>.

- Proding, B., O'Connor, R. J., Stucki, G., & Tennant, A. (2017). Establishing score equivalence of the Functional Independence Measure motor scale and the Barthel Index, utilising the International Classification of Functioning, Disability and Health and Rasch measurement theory. *Journal of rehabilitation medicine*, 49(5), 416–422. <https://doi.org/10.2340/16501977-2225>
- Riskesdas (2018) *Laporan Provinsi Jawa Tengah Riskesdas 2018, Kementerian Kesehatan RI*.
- Rudy, M., Widyadharma, I. P. E. and Adnyana, I. M. O. (2015). Reliability indonesian version of the hospital anxiety and depression scale (HADS) of stroke patients in sanglah general hospital denpasar. *Research Gate*, 2, 1–23. <https://doi.org/10.13140/RG.2.1.3604.5928>.
- Sacco, R. L., Kasner, S. E., Broderick, J. P., Caplan, L. R., Connors, J. J., ... Council on Nutrition, Physical Activity and Metabolism (2013). An updated definition of stroke for the 21st century: a statement for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke*, 44(7), 2064–2089. <https://doi.org/10.1161/STR.0b013e318296aeca>
- Singh, S., Shankar, R., & Singh, G. P. (2017). Prevalence and Associated Risk Factors of Hypertension: A Cross-Sectional Study in Urban Varanasi. *International journal of hypertension*, 2017, 5491838. <https://doi.org/10.1155/2017/5491838>
- Sok, S. R., Yim, E., Sim, H., & Sim, H. J. (2021). Relationships among Self-Care Competency, Presence of Depressive Symptom, and Health-Related Quality of Life of Korean Stroke Patients. *Clinical nursing research*, 30(5), 670–679. <https://doi.org/10.1177/1054773820913984>
- Syafni, A. N. (2020). Rehabilitasi Medik Pasien Pasca Stroke', *Jurnal Ilmiah Kesehatan Sandi Husada*. 12(2), 873–877. <https://doi.org/10.35816/jiskh.v10i2.428>.
- Synhaeve, N. E., Arntz, R. M., van Alebeek, M. E., van Pamel, J., Maaijwee, N. A., Rutten-Jacobs, L. C., Schoonderwaldt, H. C., de Kort, P. L., van Dijk, E. J., & de Leeuw, F. E. (2016). Women have a poorer very long-term functional outcome after stroke among adults aged 18-50 years: the FUTURE study. *Journal of neurology*, 263(6), 1099–1105. <https://doi.org/10.1007/s00415-016-8042-2>
- Teh, W. L., Abidin, E., Vaingankar, J. A., & Seow, L. S. E. (2018). Prevalence of stroke, risk factors, disability and care needs in older adult in Singapore: Results from the WiSE study. *BMJ Open*, 8(3), 1–9. <https://doi.org/10.1136/bmjopen-2017-020285>.
- Thomas, S. E., Plumber, N., Venkatapathappa, P., & Gorantla, V. (2021). A Review of Risk Factors

- and Predictors for Hemorrhagic Transformation in Patients with Acute Ischemic Stroke. *International journal of vascular medicine*, 2021, 1-12. <https://doi.org/10.1155/2021/4244267>
- Torrisi, M., De Cola, M. C., Buda, A., Carioti, L., Scaltrito, M. V., Bramanti, P., Manuli, A., De Luca, R., & Calabrò, R. S. (2018). Self-Efficacy, Poststroke Depression, and Rehabilitation Outcomes: Is There a Correlation?. *Journal of stroke and cerebrovascular diseases : the official journal of National Stroke Association*, 27(11), 3208–3211. <https://doi.org/10.1016/j.jstrokecerebrovasdis.2018.07.021>
- Uzunkaya, F. and Idil Soyulu, A. (2021). Predictors of full functional recovery in endovascularly treated patients with aneurysmal subarachnoid hemorrhage. *Turkish Journal of Medical Sciences*, 51(4), 2000–2006. <https://doi.org/10.3906/sag-2103-3>.
- Wang, C. Y., Miyoshi, S., Chen, C. H., Lee, K. C., Chang, L. C., Chung, J. H., & Shi, H. Y. (2020). Walking ability and functional status after post-acute care for stroke rehabilitation in different age groups: a prospective study based on propensity score matching. *Aging*, 12(11), 10704–10714. <https://doi.org/10.18632/aging.103288>
- Wang, Y., Hafeez, A., Meng, F., Zhang, R., Wang, X., Chen, X., Kong, Q., Du, H., & Ma, X. (2016). The correlation of D-dimer levels with patient outcomes in acute ischemic cerebrovascular disease complicating coronary heart disease. *Neurological research*, 38(6), 524–532. <https://doi.org/10.1080/01616412.2016.1187829>
- Xing, Y., Yang, S., Dong, F., Wang, M., Feng, Y., & Zhang, F. (2018). The beneficial role of early exercise training following stroke and possible mechanisms. *Life Sciences*. Elsevier, 198(139), 32–37. <https://doi.org/10.1016/j.lfs.2018.02.018>.
- Yahya, T., Jilani, M. H., Khan, S. U., Mszar, R., Hassan, S. Z., Blaha, M. J., Blankstein, R., Virani, S. S., Johansen, M. C., Vahidy, F., Cainzos-Achirica, M., & Nasir, K. (2020). Stroke in young adults: Current trends, opportunities for prevention and pathways forward. *American journal of preventive cardiology*, 3, 100085. <https://doi.org/10.1016/j.ajpc.2020.100085>