

## DAFTAR PUSTAKA

1. Kemenkes RI. (2020). Pedoman Pencegahan dan Pengendalian COVID-19. Jakarta: Kemenkes RI.
2. WHO. (2022). Data Pasien COVID-19. Jakarta: WHO.
3. Iriani. (2022). Prevalensi, Onset, dan Durasi Gangguan Olfaktori dan Gangguan Gustatori pada Kasus COVID-19. *Journal of Clinical Medicine*, 9(2)141-146.
4. CDC. (2021). *Data and Statistics on Venous Thromboembolism* / CDC. [online] Available at: <<https://www.cdc.gov/ncbddd/dvt/data.html>> [Accessed 31 October 2021].
5. Li et al. (2020). Coronavirus disease 2019 (COVID-19): current status and future perspectives. *International Journal of Antimicrobial Agents*, 55(5), p.105951.
6. Du Z et., al. (2020). Detection of antibodies against SARS-CoV-2 in patients with COVID-19. *J. Med. Virol.* 2020. doi:10.1002/jmv.25820.
7. Mason. (2020). Pathogenesis of COVID-19 from a cell biology perspective. *European Respiratory Journal*, 55(4), p.2000607.
8. WHO. (2020). Coronavirus Disease (COVID-19): Situation Report - 142.
9. Tong et.al. (2020). The prevalence of olfactory and gustatory dysfunction in COVID-19 patients: A systematic review and meta-analysis. *Otolaryngology Head and Neck Surgery*, 163(1), 3–11.
10. Mehraeen. (2021). Transmission Modes of COVID-19: A Systematic Review. *National Library of Medicine*, 21(6).
11. Hajikhani et al. (2020). Olfactory and gustatory dysfunction in COVID-19 patients: A meta-analysis study. *Physiological Reports*, 8(1).
12. PERHATI-KL. (2020). Terapi Gangguan Pernafasan. Jakarta: PERHATI.
13. Yasser B. Hennawi, et.al. (2023). Olfactory and Gustatory Dysfunctions Following COVID-19 Infection: Factors That Affect Their Duration in Saudi Arabia
14. Mubarak, et.al. (2021). Prevalence of anosmia among COVID-19 patients in Taif City, Kingdom of Saudi Arabia. *Saudi Med J*; Vol. 42 (1)
15. Jain, et.al. (2020). Olfactory and taste dysfunction in coronavirus disease 2019 patients: its prevalence and outcomes. *Laryngol Otol* 2020;1–5. <https://doi.org/10.1017/S0022215120002467>
16. Kandakure. (2022). Prevalence and Recovery from Newly Onset Anosmia and Ageusia in Covid 19 Patients at our Tertiary Care Centre. *Indian J Otolaryngol Head Neck Surg*, 74(2): S3399–S3406; <https://doi.org/10.1007/s12070-021-02540-w>.
17. A. Faycal. (2020). Prevalence and factors associated with symptom persistence: A prospective study of 429 mild COVID-19 outpatients. *Infectious Diseases Now* 52 (2022) 75–81

18. Hasan, et.al. (2023). The prevalence of olfactory dysfunction and its associated factors in patients with COVID-19 infection: A Hospital-based study. *Bangladesh Journal of Medical Science* Vol. 22 No. 01
19. Ramasamy, et.al. (2021). Olfactory and Gustatory Dysfunctions as a Clinical Manifestation of Coronavirus Disease 2019 in a Malaysian Tertiary Center. *Annals of Otology, Rhinology & Laryngology* Vol. 130(5) 513–519
20. Poerbonegoro. (2021). Cross-sectional study on the proportion of smell and taste disturbances in hospitalized COVID-19 patients. *Annals of Medicine and Surgery* 71.
21. Hoang *et al.* (2020). Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China', *The Lancet*, 395(10223), pp. 497–506. doi: 10.1016/S0140-6736(20)30183-5.
22. Simone Meini *et al.* (2020). Olfactory and gustatory dysfunctions in 100 patients hospitalized for COVID-19: sex differences and recovery time in real-life. *European Archives of Oto-Rhino-Laryngology* (2020) 277:3519–352. <https://doi.org/10.1007/s00405-020-06102-8>
23. Raid M. Al-Ani, et.al. (2020). Prevalence of Anosmia and Ageusia in Patients with COVID-19 at a Primary Health Center, Doha, Qatar. *Indian J Otolaryngol Head Neck Surg*
24. Yadav. (2021). Olfactory and Gustatory Dysfunction in COVID-19 Patients: A Tertiary Care Institute Experience in India. *Indian J Otolaryngol Head Neck Surg*, 74(Suppl 2):S2703–S2709; <https://doi.org/10.1007/s12070-020-02064-9>
25. Foster, et.,al. (2020). Smell loss is a prognostic factor for lower severity of coronavirus disease 2019. *Ann. Allergy Asthma Immunol*; 125(4): 481–483.
26. Mutiawati. (2021). Anosmia and dysgeusia in SARS-CoV-2 infection: incidence and effects on COVID-19 severity and mortality, and the possible pathobiology mechanisms - a systematic review and meta-analysis. *F1000Research*, 10, 40. <https://doi.org/10.12688/f1000research.28393.1>
27. Ningrum. (2022). Anosmia dan ageusia sebagai faktor prognosis baik pada pasien COVID19 rawat inap di RSUP Sanglah Denpasar. *Intisari Sains Medis*, 13(3): 566–574.
28. Butowt. (2020). SARS-CoV-2: Olfaction, brain infection and the urgent need for clinical samples allowing earlier virus detection. *ACS Chem. Neurosci.* 11 (9), 1200–1203.
29. Brychea et al. (2020). Massive transient damage of the olfactory epithelium associated with infection of sustentacular cells by SARS-CoV-2 in golden Syrian hamsters. *BioRxiv*, DOI: 10.1101/2020.06.16.151704
30. Chung, et.al. (2021). Neurosensory rehabilitation and olfactory network recovery in COVID-19-related olfactory dysfunction. *Brain Sci*, 11(2):1–8.



31. Leedman. (2021). Olfactory dysfunction at six months after coronavirus disease 2019 infection. *J Laryngol Otol*135:839–43.