



DAFTAR PUSTAKA

- Adonis Muganza, N., Olivier, M., André Kabamba, M., Charles Wembonyama, M., Oscar Numbi, L., Stanis Okitotsho, W. 2021. Predictors of mortality in neonatal sepsis in a resource-limited setting. *J. Adv. Pediatr. Child Heal.*, 4(1):057–061.
- Boeddha, N.P., Kerklaan, D., Dunbar, A., van Puffelen, E., Nagtzaam, N.M.A., Vanhorebeek, I., van den Berghe, G., Hazelzet, J.A., Joosten, K.F., Verbruggen, S.C., Dik, W.A., Driessen, G.J. 2018. HLA-DR Expression on Monocyte Subsets in Critically Ill Children. *Pediatr. Infect. Dis. J.*, 37(10):1034–1040.
- Cajander, S., Tina, E., Bäckman, A., Magnuson, A., Strålin, K., Söderquist, B., Källman, J. 2016. Quantitative real-time polymerase chain reaction measurement of HLA-DRA gene expression in whole blood is highly reproducible and shows changes that reflect dynamic shifts in monocyte surface HLA-DR expression during the course of sepsis. *PLoS One*, 11(5):1–13.
- Carey, R.N., Durham, A.P., Walter W. Hauck, PhD Anders Kallner, MD, PhD Marina V. Kondratovich, P., Jonathan Guy Middle, PhD James F. Pierson-Perry Merle B. Smith, MBA, MS, M., Aparna Srinivasan, P. 2014. User verification of precision and estimation of bias. *In: Clin. Lab. Stand. Inst. UW I-TECH Deborah Confer*, 1–81. Available at: https://clsi.org/media/1431/ep15a3_sample.pdf.
- Chen, X., Yin, Y., Zhang, J. 2011. Sepsis and immune response. *World J. Emerg. Med.*, 2(2):117.
- Doughty, C., Oppermann, L., Hartmann, N.-U., Dreschers, S., Gille, C., Orlikowsky, T. 2022. Monocytes in Neonatal Bacterial Sepsis: Think Tank or Workhorse? *BioChem*, 2(1):27–43.
- Draz, I., Dwedar, R., Hagag, S., Salama, N. 2020. Prevalence and Prognosis of Thrombocytopenia in Blood Culture Proven Neonatal Sepsis. *Egypt. J. Med. Microbiol.*, 29(1):181–186.
- Drewry, A.M., Ablordeppey, E.A., Murray, E.T., Beiter, E.R., Walton, A.H., Hall, M.W., Hotchkiss, R.S. 2016. Comparison of monocyte human leukocyte antigen-DR expression and stimulated tumor necrosis factor alpha production as outcome predictors in severe sepsis: a prospective observational study. *Crit. Care*, 1–10.
- Edmond, K., Zaidi, A. 2010. New approaches to preventing, diagnosing, and treating neonatal sepsis. *PLoS Med.*, 7(3):1–8.
- Ershad, M., Mostafa, A., Dela Cruz, M., Vearrier, D. 2019. Neonatal Sepsis. *Curr. Emerg. Hosp. Med. Rep.*, 7(3):83–90.
- Fleischmann, C., Reichert, F., Cassini, A., Horner, R., Harder, T., Markwart, R., Tröndle, M., Savova, Y., Kisson, N., Schlattmann, P., Reinhart, K., Allegranzi, B., Eckmanns, T. 2021. Global incidence and mortality of neonatal sepsis: A systematic review and meta-analysis. *Arch. Dis. Child.*, 106(8):745–752.
- García, M.A.A., González Yebra, B., López Flores, A.L., Guaní Guerra, E. 2012.



- The Major Histocompatibility Complex in Transplantation. *J. Transplant.*, 2012(June):1–7.
- Genel, F., Atlihan, F., Ozsu, E., Ozbek, E. 2010. Monocyte HLA-DR expression as predictor of poor outcome in neonates with late onset neonatal sepsis. *J. Infect.*, 60(3):224–228.
- Glaser, M.A., Hughes, L.M., Jnah, A., Newberry, D., Harris-Haman, P.A. 2021. Neonatal Sepsis: A Review of Pathophysiology and Current Management Strategies. *Adv. Neonatal Care*, 21(1):49–60.
- Gonzalez, A.C., Spearman, P.W., Stoll, B.J. 2011. Neonatal infectious diseases. *Evidenced-Based Handb. Neonatol.*, 60(2):373–396.
- Grifoni, A., Moore, E., Voic, H., Sidney, J., Phillips, E., Jadi, R., Mallal, S., De Silva, A.D., De Silva, A.M., Peters, B., Weiskopf, D., Sette, A. 2019. Characterization of magnitude and antigen specificity of HLA-DP, DQ, and DRB3/4/5 restricted DENV-specific CD4+ T cell responses. *Front. Immunol.*, 10(JULY):1–10.
- Haan, T.R. de, Beckers, L., de Jonge, R.C.J., Spanjaard, L., van Toledo, L., Pajkrt, D., van Wassenaeer-Leemhuis, A.G., van der Lee, J.H. 2013. Neonatal Gram Negative and Candida Sepsis Survival and Neurodevelopmental Outcome at the Corrected Age of 24 Months. *PLoS One*, 8(3):1–7.
- Harum, N.A., Utomo, M.T., -, A., Gunawan, P.I. 2021. the Correlation Between Apgar Score and Gestational Age With Neonatal Sepsis and Associated Mortality. *J. Widya Med.*, 7(2):141–156. Available at: <http://journal.wima.ac.id/index.php/JWM/article/view/3388>.
- Hjelmgren, H., Ygge, B.M., Nordlund, B., Andersson, N. 2022. Nurses' experiences of blood sample collection from children: a qualitative study from Swedish paediatric hospital care. *BMC Nurs.*, 21(1):1–11.
- Joshi, I., Carney, W.P., Rock, E.P. 2023. Utility of monocyte HLA-DR and rationale for therapeutic GM-CSF in sepsis immunoparalysis. *Front. Immunol.*, 14(February):1–15.
- Jung, N., Kim, C., Kim, H., Seo, Y., Hwang, J., Yang, M., Ahn, S.Y., Sung, S.I., Chang, Y.S. 2023. Changes to Blood-Sampling Protocol to Reduce the Sampling Amount in Neonatal Intensive Care Units: A Quality Improvement Project. *J. Clin. Med.*, 12(17):2–9.
- Kanakoudi-Tsakalidou, F., Debonera, F., Drossou-Agakidou, V. 2001. Flow cytometric measurement of HLA-DR expression on circulating monocytes in healthy and sick neonates using monocyte negative selection. *Clin. Exp. Immunol.*, 123(3):402–407.
- Kardana, I.M. 2011. Incidence and factors associated with mortality of neonatal sepsis. *Paediatr. Indones.*, 51(3):144.
- Kliegman, R.M., Geme, J.W.S., Blum, N.J., Shah, S.S., Tasker, R.C., Wilson, K.M., Behrman, R.E. 2020. Epidemiology of Infections. In: *Nelson Textb. Pediatr.* 21 Ed. 21st edn. Philadelphia: Elsevier, 4249–4278.
- Lai, M.Y., Tsai, M.H., Lee, C.W., Chiang, M.C., Lien, R., Fu, R.H., Huang, H.R., Chu, S.M., Hsu, J.F. 2015. Characteristics of neonates with culture-proven bloodstream infection who have low levels of C-reactive protein (≤ 10 mg/L). *BMC Infect. Dis.*, 15(1):1–9.



- Leal, Y.A., Álvarez-Nemegyei, J., Velázquez, J.R., Rosado-Quiab, U., Diego-Rodríguez, N., Paz-Baeza, E., Dávila-Velázquez, J. 2012. Risk factors and prognosis for neonatal sepsis in southeastern Mexico: analysis of a four-year historic cohort follow-up. *BMC Pregnancy Childbirth*, 12(48):1–9.
- Leventogiannis, K., Kyriazopoulou, E., Antonakos, N., Kotsaki, A., Tsangaris, I., Markopoulou, D., Grondman, I., Rovina, N., Theodorou, V., Antoniadou, E., Koutsodimitropoulos, I., Dalekos, G., Vlachogianni, G., Akinosoglou, K., Koulouras, V., Komnos, A., Kontopoulou, T., Prekates, A., Koutsoukou, A., et al. 2022. Toward personalized immunotherapy in sepsis: The PROVIDE randomized clinical trial. *Cell Reports Med.*, 3(11):3–5.
- Livaditi, O., Kotanidou, A., Psarra, A., Dimopoulou, I., Sotiropoulou, C., Augustatou, K., Papasteriades, C., Armaganidis, A., Roussos, C., Orfanos, S.E., Douzinas, E.E. 2006. Neutrophil CD64 expression and serum IL-8: Sensitive early markers of severity and outcome in sepsis. *Cytokine*, 36(5–6):283–290.
- Logothetis, D., Giourou, E., Starakis, I., Lekkou, A., Theodorou, G., Leotsinidis, M., Karakantza, M., Gogos, C. 2015. HLA-DR monocyte antigen expression as predictors of outcome in patients with community-acquired infections presenting with fever. *Crit. Care*, 19(Suppl 1):46.
- Macey, M.G. 2007. Principles of flow cytometry. *Flow Cytom. Princ. Appl.*, 1–15.
- Manzoli, T.F., Troster, E.J., Ferranti, J.F., Sales, M.M. 2016. Prolonged suppression of monocytic human leukocyte antigen-DR expression correlates with mortality in pediatric septic patients in a pediatric tertiary Intensive Care Unit. *J. Crit. Care*, 33:84–89.
- Martua, Y.S. 2021. Analisis Faktor – Faktor yang Berhubungan Dengan Kejadian Sepsis Neonatorum di RSUD Taluk Kuantan. *J. Ilm. Kesehatan.*, 13(1):55–63.
- McEnroe, R.J., Durham, A.P., Goldford, M.D., Marina V. Kondratovich, P., Samir Lababidi, P., Robert Magari, P., Jonathan Guy Middle, P., Pierson-Perry, J.F., Jeffrey E. Vaks, P. 2014. Evaluation of Precision of Quantitative Measurement Procedures; Approved Guideline. *In: Clin. Lab. Stand. Inst. Third*, 37–54.
- Milton, R., Gillespie, D., Dyer, C., Taiyari, K., Carvalho, M.J., Thomson, K., Sands, K., Portal, E.A.R., Hood, K., Ferreira, A., Hender, T., Kirby, N., Mathias, J., Nieto, M., Watkins, W.J., Bekele, D., Abayneh, M., Solomon, S., Basu, S., et al. 2022. Neonatal sepsis and mortality in low-income and middle-income countries from a facility-based birth cohort: an international multisite prospective observational study. *Lancet Glob. Heal.*, 10(5):e661–e672.
- Mohakud, N.K., Mishra, J.P., Nayak, M.K., Mishra, J., Pradhan, L., Panda, S.S., Bahera, M.R., Pugulia, R. 2022. Bacteriological Profile and Outcome of Culture-Positive Neonatal Sepsis in a Special Newborn Care Unit Setting, Odisha. *Cureus*, 14(5):1–4.
- Msanga, D.R., Parpia, F., Konje, E.T., Hokororo, A., Mshana, S.E. 2021. High Mortality among Premature Neonates with Positive Blood Culture Neonatal Sepsis in a Tertiary Hospital, Tanzania: A Call for Action. *Children*, 8(11):1037.



- Odabasi, I.O., Bulbul, A. 2020. Neonatal Sepsis. *Dep. Neonatol. Sariyer Hamidiye Etfal Train. Res. Hosp. Istanbul, Turkey*, 54(2):142–158.
- Pawar, A., Raut, A., Kalrao, V., Jacob, J., Godha, I., Thomas, R. 2016. Etiology and Clinical Outcomes of Neonatal and Pediatric Sepsis. *Arch. Pediatr. Infect. Dis.*, 4(2):2–6.
- Perry, S.E., Mostafa, S.M., Wenstone, R., Shenkin, A., McLaughlin, P.J. 2012. HLA-DR regulation and the influence of GM-CSF on transcription, surface expression and shedding. *Int. J. Med. Sci.*, 1(3):126–136.
- Pfortmueller, C.A., Meisel, C., Fux, M., Schefold, J.C. 2017. Assessment of immune organ dysfunction in critical illness: utility of innate immune response markers. *Intensive Care Med. Exp.*, 5(1):8–10.
- Ree, I.M.C., Fustolo-Gunnink, S.F., Bekker, V., Fijnvandraat, K.J., Steggerda, S.J., Lopriore, E. 2017. Thrombocytopenia in neonatal sepsis: Incidence, severity and risk factors. *PLoS One*, 12(10):1–10.
- Roble, A.K., Ayehubizu, L.M., Olad, H.M. 2022. Neonatal Sepsis and Associated Factors Among Neonates Admitted to Neonatal Intensive Care Unit in General Hospitals, Eastern Ethiopia 2020. *Clin. Med. Insights Pediatr.*, 16:1–8.
- Roeslani, R.D., Amir, I., Nasrulloh, M.H., Suryani. 2013. Faktor Resiko pada Sepsis Neonatorum Awitan Dini. *Sari Pediatr.*, 14(6):363–368.
- Romli, M.I., Yuniati, T., Hilmanto, D. 2021. Presepsin level as risk factor for mortality in premature infants with neonatal sepsis. *Paediatr. Indones.*, 61(3):165–170.
- Rusmawatingtyas, D., Rahmawati, A., Makrufardi, F., Mardhiah, N., Murni, I.K., Uiterwaal, C.S.P.M., Savitri, A.I., Kumara, I.F., Nurnaningsih. 2021. Factors associated with mortality of pediatric sepsis patients at the pediatric intensive care unit in a low-resource setting. *BMC Pediatr.*, 21(1):1–10.
- Salendu, P.M. 2013. Sepsis Neonatorum Dan Pneumonia Pada Bayi Aterm. *J. Biomedik*, 4(3):175–179.
- Salsabila, K., Toha, N.M.A., Rundjan, L., Pattanittum, P., Sirikarn, P., Rohsiswatmo, R., Wandita, S., Hakimi, M., Lumbiganon, P., Green, S., Turner, T. 2022. Early-onset neonatal sepsis and antibiotic use in Indonesia: a descriptive, cross-sectional study. *BMC Public Health*, 22(1):1–12.
- Sheikh, A., Javed, T., Afzal, M., Sheikh, C. 2010. Course and Complications of Early Onset Neonatal Sepsis : A Descriptive Study. *Annals*, 16(4):307–310.
- Simonsen, K.A., Anderson-Berry, A.L., Delair, S.F., Dele Davies, H. 2014. Early-onset neonatal sepsis. *Clin. Microbiol. Rev.*, 27(1):21–47.
- Song, J., Park, D.W., Moon, S., Cho, H.J., Park, J.H., Seok, H., Choi, W.S. 2019. Diagnostic and prognostic value of interleukin-6, pentraxin 3, and procalcitonin levels among sepsis and septic shock patients: A prospective controlled study according to the Sepsis-3 definitions. *BMC Infect. Dis.*, 19(1):1–11.
- Tillinger, W., Jilch, R., Waldhoer, T., Reinisch, W., Junger, W. 2013. Monocyte human leukocyte Antigen-DR expression - A tool to distinguish intestinal bacterial infections from inflammatory bowel disease? *Shock*, 40(2):89–94.
- Turhan, E.E., Gürsoy, T., Ovalı, F. 2015. Factors which affect mortality in neonatal



- sepsis. *Turk Pediatr. Ars.*, 50(3):170–175.
- Utomo, M.T. 2010a. Neonatal Sepsis in Low Birth Weight Infants in Dr. Soetomo General Hospital. *Indones. J. Trop. Infect. Dis.*, 1(2):86.
- Utomo, M.T. 2010b. Risk Factors of Neonatal Sepsis: A Preliminary Study in Dr. Soetomo Hospital. *Indones. J. Trop. Infect. Dis.*, 1(1):23.
- Vizcarra-Jiménez, D., Copaja-Corzo, C., Hueda-Zavaleta, M., Parihuana-Travezaño, E.G., Gutierrez-Flores, M., Rivarola-Hidalgo, M., Benites-Zapata, V.A. 2022. Predictors of Death in Patients with Neonatal Sepsis in a Peruvian Hospital. *Trop. Med. Infect. Dis.*, 7(11):342.
- Weiss, S.L., Peters, M.J., Alhazzani, W., Agus, M.S.D., Flori, H.R., Inwald, D.P., Nadel, S., Schlapbach, L.J., Tasker, R.C., Argent, A.C., Brierley, J., Carcillo, J., Carrol, E.D., Carroll, C.L., Cheifetz, I.M., Choong, K., Cies, J.J., Cruz, A.T., De Luca, D., et al. 2020. Surviving sepsis campaign international guidelines for the management of septic shock and sepsis-associated organ dysfunction in children. *Intensive Care Med.*, 46(1):10–67.
- Wibowo, T., Haksari, E.L., Wandita, S., Anggraini, A., Safrida, E.N. 2019. *Panduan Praktik Klinis Sepsis Neonatorum*.
- Wu, J.F., Ma, J., Chen, J., Ou-Yang, B., Chen, M.Y., Li, L.F., Liu, Y.J., Lin, A.H., Guan, X.D. 2011. Changes of monocyte human leukocyte antigen-DR expression as a reliable predictor of mortality in severe sepsis. *Crit. Care*, 15(5):1–8.
- Wynn, J.L., Wong, H.R. 2010. Pathophysiology and Treatment of Septic Shock in Neonates. *Clin. Perinatol.*, 37(2):439–479.
- Wynn, J.L., Wong, H.R. 2017. Pathophysiology of Neonatal Sepsis. *In: Fetal Neonatal Physiol.* Elsevier, 1536–1552.
- Yetti, H., Herlina, H., Mayetti, M., Abdiana, A., Ihsan, I. 2019. CD64 Index as a Predictor of Outcome for Children with Systemic Inflammatory Response Syndrome (SIRS). *In: Proc. Proc. 1st EAI Int. Conf. Med. Heal. Res. IComHER Novemb. 13-14th 2018, Padang, West Sumatera, Indones.* EAI, 1–8.
- Zhuang, Y., Peng, H., Chen, Yuanzhuo, Zhou, S., Chen, Yanqing. 2017. Dynamic monitoring of monocyte HLA-DR expression for the diagnosis , prognosis , and prediction of sepsis 3 . Sepsis and the Immune Response : Reduced Expression of HLA-DR. *Front. Biosci.*, 1(301):1344–1354.