

DAFTAR PUSTAKA

- Adi Nugroho, Suwarman, A. Muthalib Nawawi. (2013). Hubungan antara Rasio Neutrofil-Limfosit dan Skor Sequential Organ Failure Assesment pada Pasien yang Dirawat di Ruang Intensive Care Unit. *Jurnal Anestesi Perioperatif [JAP]*. 2013;1(3) : 189-96]
- Ahmed, M. A. S. & Mohammed, S. A., (2018). Neutrophyl-to-lymphocyte ratio as a prognostic marker in critically-ill septic patients. *Research and Opinion in Anesthesia and Intensive Care*, Volume 5, pp. 279-286.
- Ali AT. (2013). Risk factors for endometrial cancer. *Ceska Gynekologie*. Vol. 78(5):548
- Ambaraningrum SL, Yetti Hernaningsih, Edward Kusuma, Hartono Kahar. (2022). Cut-Off Value of Procalcitonin in Sepsis and Septic Shock patients at Dr. Soetomo Hospital. *Indonesianjournalofclinicalpathology*. 2022 March; 28(2): 179-184
- American College of Obstetricians and Gynecologist (ACOG). (2018). Practice Bulletin No. 195. *Obstetrics & Gynecology*, 131(6), e172–e189.
- Angela MK, Wichmann MW, Ayala A, Cloffi WG, Chaudry IH. (1997). Testosterone receptor blockade after hemorrhage in males. *Arch Surg*, 132 : 1207-1214
- Angus, D.C., W.T. Linde-Zwirble, J. Lidicker, G. Clermont, J. Carcillo and M.R. Pinsky, 2001. Epidemiology of severe sepsis in the United States: Analysis of incidence, outcome and associated costs of care. *Critical Care Med.*, 29: 1303-1310
- Annborn M, Josef Dankiewicz, David Erlingec, Sabine Hertel, Malin Rundgren, J. Gustav Smith, Joachim Struck, Hans Friberg. (2013). Procalcitonin after cardiac arrest - an indicator of severity of illness, ischemia-reperfusion injury and outcome. *Resuscitation*. 84 : 782-787
- Arif, S.K. Rukka A.B.S. and Wahyuni S.. (2017). Comparison of Neutrophils-lymphocytes Ratio and Procalcitonin Parameters in Sepsis Patient Treated in Intensive Care Unit Dr. Wahidin Hospital, Makassar, Indonesia. *J. Med. Sci.*, 17 (1): 17-21 January-March, 2017 DOI: 10.3923/jms.2017.17.21
- Ashitha , L. V. , Vanimaya, Shilpa Ravindran, R. Saikant, S. Lakshmi, R. Kartik and Manoj. G. (2017). Procalcitonin: a promising diagnostic marker for sepsis and antibiotic therapy. *Journal of Intensive Care*. .5:51.
- Becker K., Richard Snider and Eric S. Nylen. (2010). Procalcitonin in sepsis and systemic inflammation: a harmful biomarker and a therapeutic target. *Br J Pharmacol*. 159, 253–264

- Bone R.C., et al. (1992). Definitions for sepsis and organ failure and guidelines for the use of innovative therapies in sepsis. The ACCP/SCCM Consensus Conference Committee. American College of Chest Physicians/Society of Critical Care Medicine. *The ACCP/SCCM Consensus Conference Committee. American College of Chest Physicians/Society of Critical Care Medicine*, 101 : 1644-1655
- Brady, J. Shahd Horie and John G. Laffey (2020). Role of the adaptive immune response in sepsis. *Intensive care medicine*. 8 : 20
- Carlota, G., Adaia Albasanz-Puig^{1,3}, Guillermo Cuervo^{1,3} and Jordi Carratalà (2021). Understanding and Managing Sepsis in Patients With Cancer in the Era of Antimicrobial Resistance. *Frontiers in Medicine*, 8 :1-15
- Chandra, A., Enkhbaatar, P., Nakano, Y., Traber, LD., Traber, D.L ., 2006. Sepsis: Emerging role of nitric oxide and selectins. *CLINICS*, 61(1), pp. 71-6
- Chen XH, Yin YJ, Zhang JX. (2011). Sepsis and Immune Response. *World J Emerg Med*. 2:88-92
- Claudia, G. Heilmann E., Molitor A., Schuetz P. (2020). Role of procalcitonin use in the management of sepsis. *Journal of Thoracic Disease*. 12 : S5-S15
- Cornelis, P., et al. (2010). Lymphocytopenia and neutrophil-lymphocyte count ratio predict bacteremia better than conventional infection markers in an emergency care unit. *critical care*, 8192.
- Da Silva, FP, V.Nizet. (2009). Cell death during sepsis: Integration of disintegration in the inflammatory response to overwhelming infection. *Apoptosis*. 14:509-521
- Deborah, J. S., et al. (2011). The Pathogenesis of Sepsis. *National Institutes of Health*, 19-48.
- Domonkos, T., et al. (2016). Early procalcitonin kinetics and appropriateness of empirical antimicrobial therapy in critically ill patients: A prospective observational study. *J Crit Care*.
- Dumas, E. Catherine Martel, Paul-Eduard Neagoe, Arnaud Bonnefoy, Martin G. Sirois. (2012), Angiopoietin-1 but not angiopoietin-2 promotes neutrophil viability: Role of interleukin-8 and platelet-activating factor, *Biochimica et Biophysica Acta - Molecular Cell Research. Elsevier B.V.*, vol. 1823, no. 2, pp. 358–367
- Fabiano, P. D., Nizet V. (2009). Cell death during sepsis: integration of disintegration in the inflammatory response to overwhelming infection. *Cell Death and Disease*. 14:509–521

- Fathi, M., Markazi-Moghaddam, N., & Ramezankhani, A. (2018). A systematic review on risk factors associated with sepsis in patients admitted to intensive care units. *Australian Critical Care*. p 1-10
- Girard, T. D., Opal, S. M., & Ely, E. W. (2005). Insights into Severe Sepsis in Older Patients: From Epidemiology to Evidence-Based Management. *Clinical Infectious Diseases*, 40(5), 719–727
- Glauben, L., et al. (2014). Chronic inflammation and cytokines in the tumor microenvironment. *Journal Immunology Research*. 149185, 19
- Glenn, L., and Whipple A. (2010). The sirens' song of empowerment: a case study of health promotion and the New Zealand Prostitutes Collective. *Glob Health Promot*, 17 : 33-8
- Grivennikov, S. I., Greten, F. R. and Karin, M. (2011). Immunity, Inflammation, and Cancer. *Cell*, vol. 140, no. 6, pp. 883–899
- Gudiol Carlota, Adaia Albasanz-Puig, Guillermo Cuervo and Jordi Carratalà. (2021). Understanding and Managing Sepsis in Patients With Cancer in the Era of Antimicrobial Resistance. *Front. Med.* 8:636547
- Guyton AC, Hall JE. 2006. Fisiologi kedokteran. Edisi ke-9. Dalam: Setiawan I, penyunting. Resistensi tubuh terhadap infeksi: I. Leukosit, granulosit, system makrofag-monosit, dan inflamasi. Jakarta. EGC
- Hanahan, D. and Weinberg, R. A. 2011, 'Hallmarks of cancer: The next generation', *Cell*. Elsevier Inc., vol. 144, no. 5, pp. 646–674
- Hotchkiss R, Karl I. (2003). The pathophysiology and treatment of sepsis. *N Eng J Med*. 2003; 348:138-50
- Hui, B. H., et al. (2017). Procalcitonin-guided antibiotic therapy in intensive care unit patients: a systematic review and meta-analysis. *Ann Intensive Care*, 7:114.
- Irene M.R, Gómez-Laguna J, and Carrasco L. (2013). Impact of PRRSV on activation and viability of antigen presenting cells. *World Journal of Virology*. 2(4): 146-151
- Jager CPC, Wijk PTL, Mathoera RB, Leuvink JJ, Poll T, Wever PC. (2010). Lymphocytopenia and neutrophil-lymphocyte count ratio predict bacteremia better than conventional infection markers in an emergency care unit. *Crit Care Med*. 2010;14:1–8.
- James, M. O. (2007). Sepsis. *The American Journal of Medicine*, 1012-1022.
- Jigeun, Y., et al. (2020). Incidence and clinical course of septic shock in neutropenic patients during chemotherapy for gynecological cancers. *Journal of Gynecologic Oncology*. 1(5):e62

- Jilma, B., Blann, A., Pernerstorfer, T., Stohlawetz, P., Eichler, H.-G., Vondrovec, B., Wagner, O. F. (1999). Regulation of Adhesion Molecules during Human Endotoxemia. *American Journal of Respiratory and Critical Care Medicine*, 159(3), 857–863
- Khaled Ali Esmaeil Ali Shalaby, Tarek Elsayed Gouda, Afaf Abdel Hafez Abdel Mageed. Neutrophil -To- Lymphocyte Ratio [NLR] as A Promising Prognostic Marker in Critically Ill Septic Patients. *The Egyptian Journal of Hospital Medicine* (January 2022) Vol. 86, Page 242-249
- Kaml, G. J. & Davis, K. A., (2016). Surgical critical care for the patient with sepsis and multiple organ dysfunction. *Anesthesiology Clin*, Volume 34, pp. 681-96
- Kaponis Apostolos, Theodoros Filindris and George Decavalas. (2012). Septic Shock in Obstetrics and Gynecology. *Patra University School of Medicine, Patra Greece*. 15 : 312-334
- Kementerian Kesehatan Republik Indonesia (Kemenkes RI). 2018. Penyakit Paru Obstruktif Kronik (PPOK). Direktorat Jenderal Pencegahan dan Pengendalian Penyakit
- Kenneth, B. (2010). Procalcitonin in sepsis and systemic inflammation: a harmful biomarker and a therapeutic target. *Br J Pharmacol*.
- Kingsley, S. M. (2017). Role of microRNAs in sepsis. *Inflammation research*.
- Lambden Simon , Pierre Francois Laterre , Mitchell M. Levy and Bruno Francois. 2019. The SOFA score—development, utility and challenges of accurate assessment in clinical trials. *BMC Critical Care* 23:374
- Levi MM, Fink MP, Marshall JC, et al. 2003. International Sepsis Definition Conference. *Crit Care Med*. 31 : 1250-1256
- Liu, X. (2016). Prognostic Significance of Neutrophil-to-Lymphocyte Ratio in Patients with Sepsis: A Prospective Observational Study. *Mediators of Inflammation*. 8191254,1-8
- Ljungstrom, L.L., D. Karlsson, A. Pernestig, R. Andersson and G. Jacobsson, 2015. Neutrophil to lymphocyte count ratio performs better than procalcitonin as a biomarker for bacteremia and severe sepsis in the emergency department. *Crit. Care*, 19: 66-66
- Lortet-Tieulent J, Ferlay J, Bray F, Jemal A. (2017). International patterns and trends in endometrial cancer incidence, 1978–2013. *JNCI*. 2017;110(4):354– 361
- Lu, A., Haifeng Li, Yuming Zheng, Minzhong Tang, Jun Li, Huihui Wu, et al. (2017). 'Prognostic Significance of Neutrophil to Lymphocyte Ratio, Lymphocyte to Monocyte Ratio, and Platelet to Lymphocyte Ratio in Patients with Nasopharyngeal Carcinoma.', *BioMed research international*, vol. 2017, p. 3047802

- Ljungstrom L, Anna-Karin Pernestig , Gunnar Jacobsson, Rune Andersson, Barbara Usener, and Diana Tilevik. (2017). Diagnostic accuracy of procalcitonin, neutrophil-lymphocyte count ratio, C-reactive protein, and lactate in patients with suspected bacterial sepsis. *PLoS ONE* 12(7): e0181704
- Madhutandra, S. (2012). Gynecological Malignancies: Epidemiological Characteristics. *Asian Pacific J Cancer Prev*, 13 : 2997-3004.
- Martin, A. et al. (2013). Procalcitonin after cardiac arrest - an indicator of severity of illness, ischemia-reperfusion injury and outcome. *Resuscitation*. 84 : 782-787
- Mervyn, S. et al (2016). The Third International Consensus Definitions for Sepsis and. *JAMA*, 315(8): 801–810
- National Cancer Institute. 2018. Medullary Thyroid Cancer (MTC). *National Institute of Health (NIH)*. Tersedia di www.cancer.gov. Diakses pada 10 Juni 2022
- Newton, S., et al. (2004). Sepsis-Induced Changes in Macrophage Co-Stimulatory Molecule Expression. *National Institutes of Health*. 5(4): 375–383
- O'Brien J., Naeem A. Ali, Scott K. Aberegg, Edward Abraham. (2007). Sepsis. *The American Journal of Medicine*, 2007, 120 : 1012-1022.
- Oczkowski S, Al Mubarak Y, Alhazzani W, Rhodes A, Kübler A, Jankowski M. (2022). Sepsis and Septic Shock. *McMaster Textbook of Internal Medicine*. Kraków: Medycyna Praktyczna .
<https://empendium.com/mcmtextbook/chapter/B31.II.18.7> Accessed March 10, 2022
- Pablo Raul, Jorge Monserrat, Alfredo Prieto, and Melchor Alvarez-Mon. (2014). Role of Circulating Lymphocytes in Patients with Sepsis. *BioMed Research International*. Volume 2014
- Perhimpunan Endokrinologi Indonesia (Perkeni). (2021). Pedoman Pengelolaan dan Pencegahan Diabetes Melitus Tipe 2 Dewasa di Indonesia. Penerbit PB Perkeni
- Perhimpunan Kardiologi Indonesia (Perki). 2015. Pedoman Tatalaksana Sindrom Koroner Akut. Perki Edisi ketiga, hal. 2.
- Perhimpunan Kardiologi Indonesia (Perki). 2020. Pedoman Tatalaksana Gagal Jantung. Perki Edisi kedua, hal. 22.
- Rainer, T. H., Chan, T. Y. F., & Cocks, R. A. (1999). Do peripheral blood counts have any prognostic value following trauma? *Injury*, 30(3), 179–185.
- Rajab, R., 2012. Hubungan kadar laktat dengan disfungsi organ multipel pada pasien sepsis di Makassar. Ph.D. Thesis, Universitas Hasanuddin, Makassar
- Riche F, Gayat E, Barthélémy R et al. (2015): Reversal of neutrophil-to-lymphocyte count ratio in early versus late death from septic shock. *Critical Care*, 19(1): 1-10.

- Rello J., Francisco Valenzuela-Sa´nchez . Maria Ruiz-Rodriguez. Sepsis: A Review of Advances in Management. *Adv Ther* 34:2393–2411
- Romdhoni A. 2009. SIRS/Sepsis Dan Syok Septik Pada Penderita Tumor Ganas Kepala dan Leher. *Jurnal THT-KL*. Vol.2, No.1 : 48 - 61
- Sakr Yasser, Cristina Elia , Luciana Mascia , Bruno Barberis , Silvano Cardellino , Sergio Livigni , et al. (2013). The influence of gender on the epidemiology of and outcome from severe sepsis. *Critical Care* 2013, 17:R50
- Shen Xiao-Fei, Ke Cao b, Jin-peng Jiang C. , Wen-Xian Guan , Jun-Feng Du. 2017. Neutrophil dysregulation during sepsis: an overview and update. *J. Cell. Mol. Med.* Vol 21, No 9, 2017 pp. 1687-1697
- Shiferaw, B. (2016). The Role of Procalcitonin as a Biomarker in Sepsis. *Journal of Infectious Disease and Epidemiology*. 2 : 006
- Singer M., et al. (2016). The Third International Consensus Definitions for Sepsis and Septic Shock (Sepsis-3). *JAMA*. 315(8):801-810
- Song J, Park DW, Moon S, Cho HJ, Park JH, Seok H, Choi WS. Diagnostic and prognostic value of interleukin-6, pentraxin 3, and procalcitonin levels among sepsis and septic shock patients: A prospective controlled study according to the sepsis-3 definitions. *BMC Infect Dis*, 2019; 19(1): 968
- Subedi, K., Singh M., and Ojha N. (2017). Histopathological Pattern of Gynecological Malignancy at Tribhuvan University Teaching Hospital, Nepal. *Nepal Journal of Obstetrics and Gynaecology*, 2:44-47.
- Wataru, Y., et al (2017). Clinical Statistics of Gynecologic Cancers in Japan. *Journal of Gynecologic Oncology*. 28(2):e32
- Wibowo A.A.A. (2020). Hubungan Antara Neutrophyl-Lymphocyte Ratio dan Kadar Prokalsitonin Plasma Pada Pasien Sepsis Yang Dirawat Di Ruang Intensif Rsup Dr Sardjito. Tesis. Departemen Anestesiologi Dan Terapi Intensif Fakultas Kedokteran, Kesehatan Masyarakat Dan Keperawatan Universitas Gadjah Mada
- World Health Organization. (2020). The Global Cancer Observatory. <https://gco.iarc.fr/today/data/factsheets/populations/360-indonesia-factsheets.pdf>
- Xing-hai Chen, Yong-jie Yin, Jing-xiao Zhang.(2011). Sepsis and immune response. *World J Emerg Med*;2:88–92. 3.
- Xu, G. G, Guo X.G., Ji T. X., and Chen Q (2014). Neutrophil count to lymphocyte count ratio is a potential diagnostic index for bacteremia in adult. *Life Science Journal*, 172-177.

- Xuan , L., Shen Y., Wang H., Ge Q., Fei A., and Pan S. (2016). Prognostic Significance of Neutrophil-to-Lymphocyte Ratio in. *Mediators of Inflammation*. Volume 2016, Article ID 8191254, 8 pages
- Yoo J., Jung Y., Jung Hwan Ahn, Youn J. C., Keun H., Hur S. Incidence and clinical course of septic shock in neutropenic patients during chemotherapy for gynecological cancers. *Journal of Gynecol Oncol*. 2020 Sep;31(5):e62
- Zahorec. (2001). Ratio of neutrophil to lymphocyte counts — rapid and simple parameter of systemic inflammation and stress in critically ill. *Bratisl Lek Listy*; 102 (1): 5–14