



## DAFTAR PUSTAKA

- Ahmed WAS, Hamdy MA. (2018). Optimal management of umbilical cord prolapse. *International Journal of Women's Health*, 10, 459–465. <https://doi.org/10.2147/IJWH.S130879>. *International Journal of Women's Health*, 10, 459–465.
- Aiken AR, Aiken CE, Alberry MS, Brockelsby JC, SSott, JG. (2015). Management of fetal malposition in the second stage of labor: A propensity score analysis. *American Journal of Obstetrics and Gynecology*, 212(3), 355.e1-355.e7. <https://doi.org/10.1016/j.ajog.2014.10.023>
- Akinlusi FM, Rabiu KA, Durojaiye IA, Adewunmi AA, Ottun TA, Oshodi YA. (2018). Caesarean delivery-related blood transfusion: Correlates in a tertiary hospital in Southwest Nigeria. *BMC Pregnancy and Childbirth*, 18(1). <https://doi.org/10.1186/s12884-017-1643-7>
- Al-Zirqi I, Daltveit AK, Forsén L, Stray-Pedersen B, Vangen S. (2017). Risk factors for complete uterine rupture. *American Journal of Obstetrics and Gynecology*, 216(2), 165.e1-165.e8. <https://doi.org/10.1016/j.ajog.2016.10.017>
- Allen VM, O'Connell CM, Liston RM, Baskett TF. (2003). Maternal morbidity associated with cesarean delivery without labor compared with spontaneous onset of labor at term. *Obstetrics and Gynecology*, 102(3), 477–482. [https://doi.org/10.1016/S0029-7844\(03\)00570-2](https://doi.org/10.1016/S0029-7844(03)00570-2)
- American College of Obstetricians and Gynecologists (ACOG). Obstetric Care Consensus No.11. (2022). Pregnancy at Age 35 Years or Older. *Obstet Gynecol*, 140(2), 348–366.
- Ashar H, Kusrini I. (2020). *Determinant of the Increased Sectio Caesarea Labor Rates of Indonesia in 2017*. 22(Ishr 2019), 268–272. <https://doi.org/10.2991/ahsr.k.200215.051>
- Baskett TF. (2015). Preparedness for Emergency “Crash” Caesarean Section. *Journal of Obstetrics and Gynaecology Canada*, 37(12), 1116–1117. [https://doi.org/10.1016/S1701-2163\(16\)30078-0](https://doi.org/10.1016/S1701-2163(16)30078-0)
- Battula SP, Mohammed NH, Datta S. (2021). Antepartum haemorrhage. *Obstetrics, Gynaecology and Reproductive Medicine*, 31(4), 117–123. <https://doi.org/10.1016/j.ogr.2021.02.001>
- Benli AR, Cetin Benli N, Usta AT, Atakul T, Koroglu M. (2015). Effect of Maternal Age on Pregnancy Outcome and Cesarean Delivery Rate. *Journal of Clinical Medicine Research*, 7(2), 97–102. <https://doi.org/10.14740/jocmr1904w>
- Bergholt T, Skjeldestad FE, Pyykönen A, Rasmussen SC, Tapper AM, Bjarnadóttir RI, Smárason A, Másdóttir BB, Klungsøyr K, Albrechtsen S, Källén K, Gissler M, Løkkegaard ECL. (2020). Maternal age and risk of cesarean section in women with induced labor at term—A Nordic register-based study. *Acta Obstetricia et Gynecologica Scandinavica*, 99(2), 283–289. <https://doi.org/10.1111/aogs.13743>



- Berguna JSN, Basabih M. (2023). Evaluation of Decision-to-Delivery Interval and Its Effect on Maternal and Neonatal Outcomes in Category-1 and Category-2 Emergency Caesarean Section Deliveries: A Systematic Review. *JMMR (Jurnal Medicoeticolegal Dan Manajemen Rumah Sakit)*, 12(2), 233–252. <https://doi.org/10.18196/jmmr.v12i2.57>
- Betrán AP, Ye J, Moller AB, Zhang J, Gülmezoglu AM, Torloni MR. (2016). The increasing trend in caesarean section rates: Global, regional and national estimates: 1990-2014. *PLoS ONE*, 11(2), 1–12. <https://doi.org/10.1371/journal.pone.0148343>
- Bishop D, Dyer RA, Maswime S, Rodseth RN, Dyk D, Kluyts HL, Tumukunde J, Madzimbamuto FD, Elkhogia AM, Ndonga AKN, Ngumi ZWW, Omigbodun AO, Amanor-Boadu SD, Zoumenou E, Basenero A, Munlemvo DM, Youssouf C, Ndayisaba G, Antwi-Kusi A, Zubia NZ. (2019). Maternal and neonatal outcomes after caesarean delivery in the African Surgical Outcomes Study: a 7-day prospective observational cohort study. *The Lancet Global Health*, 7(4), e513–e522. [https://doi.org/10.1016/S2214-109X\(19\)30036-1](https://doi.org/10.1016/S2214-109X(19)30036-1)
- Bloom SL, Leveno KJ, Spong CY, Gilbert S, Hauth JC, Landon MB, Varner MW, Moawad AH, Caritis SN, Harper M, Wapner RJ, Sorokin Y, Miodovnik M, O'sullivan MJ, Sibai BM, Langer O, Gabbe SG. (2006). Decision-to-Incision Times and Maternal and Infant Outcomes. In *Obstet Gynecol* (Vol. 108).
- Butwick AJ, Walsh EM, Kuzniewicz M, Li SX, Escobar GJ. (2017). Patterns and predictors of severe postpartum anemia after Cesarean section. *Transfusion*, 57(1), 36–44. <https://doi.org/10.1111/trf.13815>
- Caughey AB, Cahill AG, Guise JM, Rouse DJ. (2014). Safe prevention of the primary cesarean delivery. *American Journal of Obstetrics and Gynecology*, 210(3), 179–193. <https://doi.org/10.1016/j.ajog.2014.01.026>
- Chami HW, Gueye M, Ndiaye MD, Wade M, Diouf A, Diakhate A, Ndour SB, Niang N, Mbaye M. (2019). Relation between Obstetric Outcome and Parity. *Open Journal of Obstetrics and Gynecology*, 09(06), 894–903. <https://doi.org/10.4236/ojog.2019.96087>
- Chow KM, Hkjgom, SLM. (2015). Maternal and Fetal Outcomes in Extremely Urgent Caesarean Delivery in Relation to the Decision-to-delivery Interval. In *Hong Kong J Gynaecol Obstet Midwifery* (Vol. 15, Issue 1).
- Cunningham FG, Leveno KJ, Dashe JS, Hoffman BL, Spong CY, Casey BM. *Williams Obstetrics 26 Edition*. United State, McGraw Hill. 2022:1405-1474
- Dai J, Shi Y, Wu Y, Guo L, Lu D, Chen Y, Wang Y, Lai H, Kong X. (2023). The interaction between age and parity on adverse pregnancy and neonatal outcomes. *Frontiers in Medicine*, 10. <https://doi.org/10.3389/fmed.2023.1056064>
- Dessu S, Samuel S, Gebremeskel F, Basazin A, Tariku Z, Markos M. (2021). Determinants of post cesarean section surgical site infection at public hospitals in Dire Dawa administration, Eastern Ethiopia: Case control study. *PLoS ONE*, 16(4 April). <https://doi.org/10.1371/journal.pone.0250174>
- Dorjey Y, Tshomo Y, Wangchuk D, Bhandari P, Dorji C, Pradhan D, Pemo R. (2023). Evaluation of decision to delivery interval and its effect on feto-



- maternal outcomes in Category-I emergency cesarean section deliveries in Phuentsholing General Hospital, 2020: A retrospective cross-sectional study. *Health Science Reports*, 6(1). <https://doi.org/10.1002/hsr2.1050>
- Dupuis O, Sayegh I, Decullier E, Dupont C, Clément HJ, Berland M, Rudigoz RC. (2008). Red, orange and green Caesarean sections: A new communication tool for on-call obstetricians. *European Journal of Obstetrics and Gynecology and Reproductive Biology*, 140(2), 206–211. <https://doi.org/10.1016/j.ejogrb.2008.04.003>
- Ergani SY, İskender D, Aksan A, İbanoğlu MC, Çakır BT, Kınay T, Erkaya S, İskender CT, Çelen Ş, Çağlar AT, Engin-üstün Y. (2023). Postoperative care in the caesarean intensive care unit: experience from a tertiary maternity hospital. *Journal of the Turkish German Gynecology Association*, 24(1), 42–47. <https://doi.org/10.4274/jtgga.galenos.2022.2021-9-27>
- Fan D, Wu S, Liu L, Xia Q, Wang W, Guo X, Liu Z. (2017). Prevalence of antepartum hemorrhage in women with placenta previa: A systematic review and meta-analysis. *Scientific Reports*, 7. <https://doi.org/10.1038/srep40320>
- Fawcus S, Moodley J. (2013). Postpartum haemorrhage associated with caesarean section and caesarean hysterectomy. *Best Practice and Research: Clinical Obstetrics and Gynaecology*, 27(2), 233–249. <https://doi.org/10.1016/j.bpobgyn.2012.08.018>
- Frick AP. (2021). Advanced maternal age and adverse pregnancy outcomes. In *Best Practice and Research: Clinical Obstetrics and Gynaecology* (Vol. 70, pp. 92–100). Baillière Tindall Ltd. <https://doi.org/10.1016/j.bpobgyn.2020.07.005>
- Fuhrmann L, Pedersen TH, Atke A, Møller AM, Ostergaard D. (2015). Multidisciplinary team training reduces the decision-to-delivery interval for emergency Caesarean section. *Acta Anaesthesiologica Scandinavica*, 59(10), 1287–1295. <https://doi.org/10.1111/aas.12572>
- Gabbai D, Attali E, Ram S, Amikam U, Ashwal E, Hiersch L, Gamzu R, Yoge Y. (2022). Prediction model for prolonged hospitalization following cesarean delivery. *European Journal of Obstetrics and Gynecology and Reproductive Biology*, 274, 23–27. <https://doi.org/10.1016/j.ejogrb.2022.04.026>
- Ganap EP, Hakimi M, Hadijono S, Emilia O. (2016). Impact Of Obstetrician/Gynecologist Hospitalists On Response Time Of Obstetric Emergency Operation In Sardjito General Hospital Yogyakarta Indonesia: A Retrospective Cohort Study. In *Jurnal Kesehatan Reproduksi* (Vol. 3, Issue 3).
- Gelebo KG, Neme D, Destaw B, Aweke Z, Kasa, SM. (2023). The effect of preoperative anemia on perioperative outcomes among patients undergoing emergency surgery: A multicenter prospective cohort study. *Helijon*, 9(7). <https://doi.org/10.1016/j.heliyon.2023.e17804>
- Gunawan T, Attamimi A, Praduatmo H. (2018). Hubungan Response Time Seksio Sesarea Emergensi Kategori 1 Dengan Luaran Perinatal di RSUP Dr. Sardjito. In *Jurnal Kesehatan Reproduksi* (Vol. 5, Issue 1).
- Gupta S, Naithani U, Madhanmohan C, Singh A, Reddy P, Gupta A. (2017). Evaluation of decision-to-delivery interval in emergency cesarean section: A



- 1-year prospective audit in a tertiary care hospital. *Journal of Anaesthesiology Clinical Pharmacology*, 33(1), 64–70. <https://doi.org/10.4103/0970-9185.202197>
- Hadiati DR, Hakimi M, Nurdianti DS, Ota E. (2014). Skin preparation for preventing infection following caesarean section. *Cochrane Database of Systematic Reviews*, 2014(9). <https://doi.org/10.1002/14651858.CD007462.pub3>
- Harde M, Dave S, Wagh S, Gujjar P, Bhadade R, Bapat A. (2014). Prospective evaluation of maternal morbidity and mortality in post-cesarean section patients admitted to postanesthesia intensive care unit. *Journal of Anaesthesiology Clinical Pharmacology*, 30(4), 508–513. <https://doi.org/10.4103/0970-9185.142844>
- Hassan B, Mandar O, Alhabardi N, Adam I. (2022). Length of Hospital Stay After Cesarean Delivery and Its Determinants Among Women in Eastern Sudan. *International Journal of Women's Health*, 14, 731–738. <https://doi.org/10.2147/IJWH.S356855>
- Hillemanns P, Hasbargen U, Strauss A, Schulze A, Genzel-Boroviczeny O, Hepp H. (2003). Maternal and neonatal morbidity of emergency caesarean sections with a decision-to-delivery interval under 30 minutes: Evidence from 10 years. *Archives of Gynecology and Obstetrics*, 268(3), 136–141. <https://doi.org/10.1007/s00404-003-0527-4>
- Hirani BA, Mchome BL, Mazuguni NS, Mahande MJ. (2017). The decision delivery interval in emergency caesarean section and its associated maternal and fetal outcomes at a referral hospital in northern Tanzania: a cross-sectional study. *BMC Pregnancy and Childbirth*, 17(1), 411. <https://doi.org/10.1186/s12884-017-1608-x>
- Iqbal K, Iqbal A, Rathore SS, Ahmed J, Ali SA, Farid E, Hasanain M, Azeem Q, Qadar LT, Memon FR, Azim D. (2022). Risk factors for blood transfusion in Cesarean section: A systematic review and meta-analysis. In *Transfusion Clinique et Biologique* (Vol. 29, Issue 1, pp. 3–10). Elsevier Masson s.r.l. <https://doi.org/10.1016/j.trcli.2021.09.010>
- Jardine J, Blotkamp A, Gurol-Urganci I, Knight H, Harris T, Hawdon J, van der Meulen J, Walker K, Pasupathy D. (2020). Risk of complicated birth at term in nulliparous and multiparous women using routinely collected maternity data in England: cohort study. *BMJ (Clinical Research Ed.)*, 371, m3377. <https://doi.org/10.1136/bmj.m3377>
- Jauniaux E, Grobman W. (2016). *Textbook of Caesarean Section* (E. Jauniaux & W. Grobman, Eds.; 1st ed.). Oxford University Press. [www.ebook2book.ir](http://www.ebook2book.ir)
- Jharaik H, Dhiman B, Verma SK, Sharma A. (2019). Consequences of antepartum hemorrhage and its maternal and perinatal outcome. *International Journal of Reproduction, Contraception, Obstetrics and Gynecology*, 8(4), 1480. <https://doi.org/10.18203/2320-1770.ijrcog20191203>
- Kitaw TM, Limenh SK, Chekole FA, Getie SA, Gemedu BN, Engda AS. (2021). Decision to delivery interval and associated factors for emergency cesarean section: a cross-sectional study. *BMC Pregnancy and Childbirth*, 21(1). <https://doi.org/10.1186/s12884-021-03706-8>



- Kolas T, Øian P, Skjeldestad FE. (2010). Risks for peroperative excessive blood loss in cesarean delivery. *Acta Obstetricia et Gynecologica SSandinavica*, 89(5), 658–663. <https://doi.org/10.3109/00016341003605727>
- Kwek K, Yeap ML, Tan KH, Tee JCS, Yeo GSH. (2005). Crash caesarean section-decision-to-delivery interval, *Acta Obstet Gynecol SSand*, 84: 914–915.
- Lagrew DC, Bush MC, McKeown AM, Lagrew NG. (2006). Emergent (crash) cesarean delivery: Indications and outcomes. *American Journal of Obstetrics and Gynecology*, 194(6), 1638–1643. <https://doi.org/10.1016/j.ajog.2006.03.007>
- Larsson C, Djufveldt E, Lindam A, Tunón K, Nordin P. (2021). Surgical complications after caesarean section: A population-based cohort study. *PLoS ONE*, 16(10 October). <https://doi.org/10.1371/journal.pone.0258222>
- Leung TY, Lao TT. (2013). Timing of caesarean section according to urgency. *Best Practice and Research: Clinical Obstetrics and Gynaecology*, 27(2), 251–267. <https://doi.org/10.1016/j.bpobgyn.2012.09.005>
- Lucas DN, Yentis SM, Kinsella SM, Holdcroft A, May AE, Wee M, Robinson PN. (2000). Urgency of caesarean section: A new classification. *Journal of the Royal Society of Medicine*, 93(7), 346–350. <https://doi.org/10.1177/014107680009300703>
- Lumbiganon P, Laopaiboon M, Gülmезoglu AM, Souza JP, Taneepanichskul S, Ruyan P, Attygalle DE, Shrestha N, Mori R, Hinh ND, Bang HT, Rathavy T, Chyun K, Cheang K, Festin M, Udomprasertgul V, Germar MJV, Yanqiu G, Roy M, Villar J. (2010). Method of delivery and pregnancy outcomes in Asia: the WHO global survey on maternal and perinatal health 2007–08. *The Lancet*, 375(9713), 490–499. [https://doi.org/10.1016/S0140-6736\(09\)61870-5](https://doi.org/10.1016/S0140-6736(09)61870-5)
- Magann EF, Evans S, Hutchinson M, Collins R, Lanneau G, Morrison JC. (2005). Postpartum hemorrhage after cesarean delivery: An analysis of risk factors. *Southern Medical Journal*, 98(7), 681–685. <https://doi.org/10.1097/01.SMJ.0000163309.53317.B8>
- Mardiawan D, Prawitasari S. (2023). Achievement of decision-to-delivery interval in emergency cesarean section category 1 during the COVID-19 pandemic: a scoping review. In *Bali Medical Journal* (Vol. 12, Issue 1, pp. 838–842). Sanglah General Hospital. <https://doi.org/10.15562/bmj.v12i1.3976>
- Massarello KC, Horta BL, Silveira MF. (2017). Maternal complications and cesarean section without indication: systematic review and meta-analysis. *Revista de Saude Publica*, 51, 105. <https://doi.org/10.11606/S1518-8787.2017051000389>
- Mishra, N., Gupta, R., & Singh, N. (2018). Decision Delivery Interval in Emergency and Urgent Caesarean Sections: Need to Reconsider the Recommendations? *Journal of Obstetrics and Gynecology of India*, 68(1), 20–26. <https://doi.org/10.1007/s13224-017-0991-6>
- Molina G, Weiser TG, Lipsitz SR, Esquivel MM, Uribe-Leitz T, Azad T, Shah N, Semrau K, Berry WR, Gawande AA, Haynes AB. (2015). Relationship between cesarean delivery rate and maternal and neonatal mortality. *JAMA* -



- Journal of the American Medical Association*, 314(21), 2263–2270.  
<https://doi.org/10.1001/jama.2015.15553>
- Motomura K, Ganchimeg T, Nagata C, Ota E, Vogel JP, Betran AP, Torloni MR, Jayaratne K, Jwa SS, Mittal S, Recidoro ZD, Matsumoto K, Fujieda M, Nafiou I, Yunis K, Qureshi Z, Souza JP, Mori R. (2017). Incidence and outcomes of uterine rupture among women with prior caesarean section: WHO Multicountry Survey on Maternal and Newborn Health. *Scientific Reports*, 7(March), 1–9. <https://doi.org/10.1038/srep44093>
- Mylonas I, Friese K. (2015). Indications for and risks of elective cesarean section. In *DeutSches Arzteblatt International* (Vol. 112, Issues 29–30, pp. 489–495). DeutScher Arzte-Verlag GmbH. <https://doi.org/10.3238/arztebl.2015.0489>
- Ohuma EO, Jabin N, Young MF, Epie T, Martorell R, Peña-Rosas JP, Garcia-Casal MN, Kennedy SH, Victora CG, Craik R, Ash S, Barros FC, Barsosio HC, Berkley JA, Carvalho M, Fernandes M, Cheikh Ismail L, Lambert A, Lindgren CM, Villar J. (2023). Association between maternal haemoglobin concentrations and maternal and neonatal outcomes: the prospective, observational, multinational, INTERBIO-21st fetal study. *The Lancet Haematology*, 10(9), e756–e766. [https://doi.org/10.1016/S2352-3026\(23\)00170-9](https://doi.org/10.1016/S2352-3026(23)00170-9)
- Pallasmaa N, Ekblad U, Aitokallio-Tallberg A, Uotila J, Raudaskoski T, Ulander VM, Hurme S. (2010). Cesarean delivery in Finland: Maternal complications and obstetric risk factors. *Acta Obstetricia et Gynecologica Scandinavica*, 89(7), 896–902. <https://doi.org/10.3109/00016349.2010.487893>
- RCOG. (2010). Good Practice No. 11 CLASSIFICATION OF URGENCY OF CAESAREAN SECTION – A CONTINUUM OF RISK. *Good Practice No. 11*, 1–4.
- Shaikh AG, Shaikh VF, Nasim MT, Farid E, Ali SA, Azim D, Afzal H, Iqbal K, Iqbal A. (2022). Risk Factors for Blood Transfusion during Caesarean Section. *Pakistan Journal of Medical and Health Sciences*, 16(4), 339–342. <https://doi.org/10.5335/pjmhs22164339>
- Sharshiner R, Silver RM. (2015). Management of Fetal Malpresentation. *CLINICAL OBSTETRICS AND GYNECOLOGY*, 58(2), 246–255. [www.clinicalobgyn.com](http://www.clinicalobgyn.com)
- Staboulidou I, Beslic J, Kuehnle E, Kaisenberg, C, Hillemanns P, SShippert C. (2018). Neonatal and Maternal Short-Term Outcome after Emergency Caesarean Section in Comparison to Elective and Second Stage Caesareans: Results of A Retrospective 10-Year Survey. *International Journal of Women's Health and Wellness*, 4(2). <https://doi.org/10.23937/2474-1353/1510080>
- Temesgen MM, Gebregzi AH, Kasahun HG, Ahmed SA, Woldegerima YB. (2020). Evaluation of decision to delivery time interval and its effect on feto-maternal outcomes and associated factors in category-1 emergency caesarean section deliveries: Prospective cohort study. *BMC Pregnancy and Childbirth*, 20(1). <https://doi.org/10.1186/s12884-020-2828-z>
- Tempest N, Navaratnam K, Hapangama DK. (2015). Management of delivery when malposition of the fetal head complicates the second stage of labour. *The*



- Obstetrician & Gynaecologist, 17(4), 273–280.*  
<https://doi.org/10.1111/tog.12217>
- Torloni MR, Betran AP, Souza JP, Widmer M, Allen T, Gulmezoglu M, Merialdi, M. (2011). Classifications for cesarean section: A systematic review. *PLoS ONE, 6(1)*. <https://doi.org/10.1371/journal.pone.0014566>
- Vargas M, Marra A, Buonanno P, Iacobazzo C, SShiavone V, Servillo G. (2019). Obstetric admissions in ICU in a tertiary care center: A 5-years retrospective study. *Indian Journal of Critical Care Medicine, 23(5)*, 213–219. <https://doi.org/10.5005/jp-journals-10071-23163>
- Wang Y, Liu D, Wu X, Zheng C, Chen X. (2023). Effect of in situ simulation training for emergency caesarean section on maternal and infant outcomes. *BMC Medical Education, 23(1)*. <https://doi.org/10.1186/s12909-023-04772-6>
- WHO. (2011). Haemoglobin concentrations for the diagnosis of anaemia and assessment of severity. Vitamin and Mineral Nutrition Information System. Geneva, World Health Organization, 2011 (WHO/NMH/NHD/MNM/11.1 <http://www.who.int/vmnis/indicators/haemoglobin.pdf>
- Wong L, Kwan AHW, Lau SL, Sin WTA, Leung TY. (2021). Umbilical cord prolapse: revisiting its definition and management. *American Journal of Obstetrics and Gynecology, 225(4)*, 357–366. <https://doi.org/10.1016/j.ajog.2021.06.077>
- Wong TCT, Lau CQH, Tan EL, Kanagalingam D. (2017). Decision-to-delivery intervals and total duration of surgery for caesarean sections in a tertiary general hospital. *Singapore Medical Journal, 58(6)*, 332–337. <https://doi.org/10.11622/smedj.2016098>
- Wormer KC, Jamil RT, Bryant SB. Acute Postpartum Hemorrhage. [Updated 2023 May 8]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2023 Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK499988/>
- Yang XJ, Sun SS. (2017). Comparison of maternal and fetal complications in elective and emergency cesarean section: a systematic review and meta-analysis. *Archives of Gynecology and Obstetrics, 296(3)*, 503–512. <https://doi.org/10.1007/s00404-017-4445-2>
- Yeni CM, Hasanuddin, Maharani CR, Maulida NF. (2022). Evaluasi Response Time Seksio Emergensi Kategori I Terhadap Luaran Maternal dan Neonatal Dengan Tersedianya Alur Pelayanan Seksio Sesarea Emergensi Di RSUDZA Juli-Oktober 2021. *Journal of Medical Science, 3(1)*, 10–18. <https://doi.org/10.55572/jms.v3i1.57>