

DAFTAR PUSTAKA

- Abdellah, E. M., Ali, T. H., Abdou, D. A. M., Hassanein, N. M., Fadel, M., Karam El-Din, A. A., & El-Ghonemy, D. H. (2020). Enhancement of lipid productivity from a promising oleaginous fungus *Aspergillus* sp. strain EM2018 for biodiesel production: Optimization of culture conditions and identification. *Grasas y Aceites*, *71*(3), 371. <https://doi.org/10.3989/gya.0345191>
- Barbedo, J. (2013). Automatic object counting in Neubauer chambers. *Anais de XXXI Simpósio Brasileiro de Telecomunicações. XXXI Simpósio Brasileiro de Telecomunicações*. <https://doi.org/10.14209/sbrt.2013.224>
- Bardhan, P., Baruah, J., Raj, G. V. S. B., Kalita, E., & Mandal, M. (2021). Optimization of culture conditions for biomass and lipid production by oleaginous fungus *Penicillium citrinum* PKB20 using response surface methodology (RSM). *Biocatalysis and Agricultural Biotechnology*, *37*, 102169. <https://doi.org/10.1016/j.cbab.2021.102169>
- Candiotti, L. V., De Zan, M. M., Cámara, M. S., & Goicoechea, H. C. (2014). Experimental design and multiple response optimization. Using the desirability function in analytical methods development. *Talanta*, *124*, 123–138. <https://doi.org/10.1016/j.talanta.2014.01.034>
- Carvalho, A. K. F., Bento, H. B. S., Rivaldi, J. D., & de Castro, H. F. (2018). Direct transesterification of *Mucor circinelloides* biomass for biodiesel production: Effect of carbon sources on the accumulation of fungal lipids and biofuel properties. *Fuel*, *234*, 789–796. <https://doi.org/10.1016/j.fuel.2018.07.029>
- Cavalcanti Luna, M. A., Vieira, E. R., Okada, K., Campos-Takaki, G. M., & Nascimento, A. E. do. (2015). Copper-induced adaptation, oxidative stress and its tolerance in *Aspergillus niger* UCP1261. *Electronic Journal of Biotechnology*, *18*(6), 418–427. <https://doi.org/10.1016/j.ejbt.2015.09.006>
- Chatzifragkou, A., Makri, A., Belka, A., Bellou, S., Mavrou, M., Mastoridou, M., Mystrioti, P., Onjaro, G., Aggelis, G., & Papanikolaou, S. (2011). Biotechnological conversions of biodiesel derived waste glycerol by yeast and fungal species. *Energy*, *36*(2), 1097–1108. <https://doi.org/10.1016/j.energy.2010.11.040>
- Chenthamarakshan, A., Parambayil, N., Miziriya, N., Soumya, P. S., Lakshmi, M. S. K., Ramgopal, A., Dileep, A., & Nambisan, P. (2017). Optimization of laccase production from *Marasmiellus palmivorus* LA1 by Taguchi method of Design of experiments. *BMC Biotechnology*, *17*(1), 12. <https://doi.org/10.1186/s12896-017-0333-x>
- Chen, N., Xu, C., Guo, X., & Shim, H. (2022). Effects of sodium and magnesium supplement on lipid production and wastewater treatment by *Rhodospiridium toruloides*. *Renewable Energy*, *199*, 919–928. <https://doi.org/10.1016/j.renene.2022.09.043>
- Enshaieh, M., Nahvi, I., & Madani, M. (2014). Improving microbial oil production with standard and native oleaginous yeasts by using Taguchi design. *International Journal of Environmental Science & Technology*, *11*(3), 597–604. <https://doi.org/10.1007/s13762-013-0373-2>
- Fazili, A. B. A., Shah, A. M., Zan, X., Naz, T., Nosheen, S., Nazir, Y., Ullah, S., Zhang, H., & Song, Y. (2022). *Mucor circinelloides*: a model organism for oleaginous fungi and its potential applications in bioactive lipid production.

- Microbial Cell Factories*, 21(1), 29. <https://doi.org/10.1186/s12934-022-01758-9>
- Franken-Luykx, J. M. M. (1967). Spectrophotometric determination of nitrogen in wort. *Journal of the Institute of Brewing*, 73(2), 187–189. <https://doi.org/10.1002/j.2050-0416.1967.tb03034.x>
- Gusakov, A. V., Kondratyeva, E. G., & Sinitsyn, A. P. (2011). Comparison of two methods for assaying reducing sugars in the determination of carbohydrase activities. *International Journal of Analytical Chemistry*, 2011, 283658. <https://doi.org/10.1155/2011/283658>
- Hashem, A. H., Abu-Elreesh, G., El-Sheikh, H. H., & Suleiman, W. B. (2022). Isolation, identification, and statistical optimization of a psychrotolerant *Mucor racemosus* for sustainable lipid production. *Biomass Conversion and Biorefinery*. <https://doi.org/10.1007/s13399-022-02390-8>
- Haura, A., & Ilmi, M. (2024). The Potential of *Mucor irregularis* Isolated From Fruits in Producing Microbial Lipid. *Maandblad Voor Accountancy En Bedrijfseconomie*, 53(1), 19–26. <https://doi.org/10.55230/mabjournal.v53i1.2747>
- Hinshaw, J. V. (2005). The Flame Ionization Detector. *LCGC North America*.
- Ibrahim, A. G., Baazeem, A., Al-Zaban, M. I., Fawzy, M. A., Hassan, S. H. A., & Koutb, M. (2023). Sustainable Biodiesel Production from a New Oleaginous Fungus, *Aspergillus carneus* Strain OQ275240: Biomass and Lipid Production Optimization Using Box–Behnken Design. *Sustainability*, 15(8), 6836. <https://doi.org/10.3390/su15086836>
- Jeennor, S., Laoteng, K., Tanticharoen, M., & Cheevadhanarak, S. (2006). Comparative fatty acid profiling of *Mucor rouxii* under different stress conditions. *FEMS Microbiology Letters*, 259(1), 60–66. <https://doi.org/10.1111/j.1574-6968.2006.00242.x>
- Jernejc, K., & Legisa, M. (2002). The influence of metal ions on malic enzyme activity and lipid synthesis in *Aspergillus niger*. *FEMS Microbiology Letters*, 217(2), 185–190. <https://doi.org/10.1111/j.1574-6968.2002.tb11473.x>
- Jones, A. D., Boundy-Mills, K. L., Barla, G. F., Kumar, S., Ubanwa, B., & Balan, V. (2019). Microbial lipid alternatives to plant lipids. *Methods in Molecular Biology*, 1995, 1–32. https://doi.org/10.1007/978-1-4939-9484-7_1
- Kasim, M. S., Che Haron, C. H., Ghani, J. A., Mohamad, E., Izamshah, R., Mohd, A. A., & Saedon, J. B. (2014). Multi-Objective Optimization Using Box-Behnken of Response Surface Methodology for High-Speed Machining of Inconel 718. *Applied Mechanics and Materials*, 629, 487–492. <https://doi.org/10.4028/www.scientific.net/AMM.629.487>
- Knothe, G., Matheaus, A. C., & Ryan, T. W. (2003). Cetane numbers of branched and straight-chain fatty esters determined in an ignition quality tester☆. *Fuel*, 82(8), 971–975. [https://doi.org/10.1016/S0016-2361\(02\)00382-4](https://doi.org/10.1016/S0016-2361(02)00382-4)
- Minitab, LLC., M. S. (n.d.). *What is the signal-to-noise ratio in a Taguchi design?* - Minitab. Retrieved May 5, 2024, from <https://support.minitab.com/en-us/minitab/help-and-how-to/statistical-modeling/doe/supporting-topics/taguchi-designs/what-is-the-signal-to-noise-ratio/>
- Minitab, LLC, M. S. (n.d.). *Lack-of-fit and lack-of-fit tests* - Minitab. Retrieved May 5, 2024, from <https://support.minitab.com/en-us/minitab/help-and-how-to>

- to/statistical-modeling/regression/supporting-topics/regression-models/lack-of-fit-and-lack-of-fit-tests/
- Narkprasom, N., Assavarachan, R., & Wongputtisin, P. (2013). Optimization of Reducing Sugar Production from Acid Hydrolysis of Sugarcane Bagasse by Box Behnken Design. *Journal of Medical and Bioengineering*, 2(4), 238–241. <https://doi.org/10.12720/jomb.2.4.238-241>
- Ochsenreither, K., Glück, C., Stressler, T., Fischer, L., & Syldatk, C. (2016). Production strategies and applications of microbial single cell oils. *Frontiers in Microbiology*, 7, 1539. <https://doi.org/10.3389/fmicb.2016.01539>
- Okafor, N., & Okeke, B. C. (2017). *Modern Industrial Microbiology and Biotechnology* (2nd ed., p. 488). CRC Press.
- Passoth, V. (2017). Lipids of yeasts and filamentous fungi and their importance for biotechnology. In A. A. Sibirny (Ed.), *Biotechnology of yeasts and filamentous fungi* (pp. 149–204). Springer International Publishing. https://doi.org/10.1007/978-3-319-58829-2_6
- Priyadarshi, D., & Paul, K. K. (2019). Optimisation of biodiesel production using taguchi model. *Waste and Biomass Valorization*, 10(6), 1547–1559. <https://doi.org/10.1007/s12649-017-0158-9>
- Qiao, W., Tao, J., Luo, Y., Tang, T., Miao, J., & Yang, Q. (2018). Microbial oil production from solid-state fermentation by a newly isolated oleaginous fungus, *Mucor circinelloides* Q531 from mulberry branches. *Royal Society Open Science*, 5(11), 180551. <https://doi.org/10.1098/rsos.180551>
- Ramos, M. J., Fernández, C. M., Casas, A., Rodríguez, L., & Pérez, A. (2009). Influence of fatty acid composition of raw materials on biodiesel properties. *Bioresource Technology*, 100(1), 261–268. <https://doi.org/10.1016/j.biortech.2008.06.039>
- Ratledge, C. (2004). Fatty acid biosynthesis in microorganisms being used for Single Cell Oil production. *Biochimie*, 86(11), 807–815. <https://doi.org/10.1016/j.biochi.2004.09.017>
- Raven, S., Francis, A., Srivastava, C., Kezo, S., & Tiwari, A. (2019). Fungal biofuels: innovative approaches. In A. N. Yadav, S. Singh, S. Mishra, & A. Gupta (Eds.), *Recent Advancement in White Biotechnology Through Fungi: Volume 2: Perspective for Value-Added Products and Environments* (pp. 385–405). Springer International Publishing. https://doi.org/10.1007/978-3-030-14846-1_13
- Rohde, J. K., Fuh, M. M., Evangelakos, I., Pauly, M. J., Schaltenberg, N., Siracusa, F., Gagliani, N., Tödter, K., Heeren, J., & Worthmann, A. (2022). A Gas Chromatography Mass Spectrometry-Based Method for the Quantification of Short Chain Fatty Acids. *Metabolites*, 12(2). <https://doi.org/10.3390/metabo12020170>
- Somashekar, D., Venkateshwaran, G., Sambaiyah, K., & Lokesh, B. R. (2003). Effect of culture conditions on lipid and gamma-linolenic acid production by mucoraceous fungi. *Process Biochemistry*, 38(12), 1719–1724. [https://doi.org/10.1016/S0032-9592\(02\)00258-3](https://doi.org/10.1016/S0032-9592(02)00258-3)
- Tang, X., Chen, H., Chen, Y. Q., Chen, W., Garre, V., Song, Y., & Ratledge, C. (2015). Comparison of Biochemical Activities between High and Low Lipid-Producing Strains of *Mucor circinelloides*: An Explanation for the High

- Oleaginicuity of Strain WJ11. *Plos One*, 10(6), e0128396.
<https://doi.org/10.1371/journal.pone.0128396>
- Tan, Y. H., Abdullah, M. O., Nolasco-Hipolito, C., & Ahmad Zauzi, N. S. (2017). Application of RSM and Taguchi methods for optimizing the transesterification of waste cooking oil catalyzed by solid ostrich and chicken-eggshell derived CaO. *Renewable Energy*, 114, 437–447.
<https://doi.org/10.1016/j.renene.2017.07.024>
- Walker, G. M. (1994). The roles of magnesium in biotechnology. *Critical Reviews in Biotechnology*, 14(4), 311–354. <https://doi.org/10.3109/07388559409063643>
- Xia, C., Zhang, J., Zhang, W., & Hu, B. (2011). A new cultivation method for microbial oil production: cell pelletization and lipid accumulation by *Mucor circinelloides*. *Biotechnology for Biofuels*, 4, 15. <https://doi.org/10.1186/1754-6834-4-15>
- Yang, Y., Heidari, F., & Hu, B. (2019). Fungi (Mold)-Based Lipid Production. *Methods in Molecular Biology*, 1995, 51–89. https://doi.org/10.1007/978-1-4939-9484-7_3
- Yu, Y., Zhang, L., Li, T., Wu, N., Jiang, L., Ji, X., & Huang, H. (2018). How nitrogen sources influence *Mortierella alpina* aging: From the lipid droplet proteome to the whole-cell proteome and metabolome. *Journal of Proteomics*, 179, 140–149. <https://doi.org/10.1016/j.jprot.2018.03.014>
- Zhang, X., Peng, Y., Li, X., Ma, G., & Chen, X. (2015). Screening of iron-enriched fungus from natural environment and evaluation of organically bound iron bioavailability in rats. *Food Science and Technology (Campinas)*, 35(1), 58–65. <https://doi.org/10.1590/1678-457X.6454>
- Zhang, X.-Y., Li, B., Huang, B.-C., Wang, F.-B., Zhang, Y.-Q., Zhao, S.-G., Li, M., Wang, H.-Y., Yu, X.-J., Liu, X.-Y., Jiang, J., & Wang, Z.-P. (2022). Production, biosynthesis, and commercial applications of fatty acids from oleaginous fungi. *Frontiers in Nutrition*, 9, 873657.
<https://doi.org/10.3389/fnut.2022.873657>