

DAFTAR PUSTAKA

- Abbo, L.M., & Grossi, P.A. (2019). Surgical site infections: Guidelines from the American Society of Transplantation Infectious Diseases Community of Practice. *Clin. Transplant.*, 33: 289 - 294
- Abraham, C., & Sheeran, P. (2014). The health belief model, *Cambridge Handbook of Psychology, Health and Medicine, Second Edition*.
- Agnesha, F., & Rahardjo, S. (2018). Tromboemboli Pada Kehamilan. *J. Komplikasi Anestesi*, 3: 75-78
- Ahlqvist, V.H., Persson, M., Magnusson, C., & Berglind, D. (2019). Elective and nonelective cesarean section and obesity among young adult male offspring: A Swedish population-based cohort study. *PLoS Med.*, 16 (12): e1002996
- Ajuzieogu, O. V., Amucheazi, A., Ezike, H.A., Achi, J., & Abam, D.S. (2014). The efficacy of chewing gum on postoperative ileus following cesarean section in Enugu, South East Nigeria: A randomized controlled clinical trial. *Niger. J. Clin. Pract.*, 17: 739 - 742
- Ajzen, I. (2012). The theory of planned behavior, *Handbook of Theories of Social Psychology.*, 26: 1113 - 1127
- Akalpler, O., & Okumus, H. (2018). Gum chewing and bowel function after caesarean section under spinal anesthesia. *Pakistan J. Med. Sci.*, 34: 1242–1247.
- Akiboye, F., Adderley, N.J., Martin, J., Gokhale, K., Rudge, G.M., Marshall, T.P., *et al.* (2020). Impact of the Diabetes Inpatient Care and Education (DICE) project on length of stay and mortality. *Diabet. Med.*, 37: 277 - 285
- Allen, T.K., & Habib, A.S. (2018). Inadvertent Perioperative Hypothermia Induced by Spinal Anesthesia for Cesarean Delivery Might Be More Significant Than We Think: Are We Doing Enough to Warm Our Parturients? *Anesth. Analg.*, 126: 7 - 9.
- Aluri, S., & Wrench, I.J. (2014). Enhanced recovery from obstetric surgery: A UK survey of practice. *Int. J. Obstet. Anesth.*, 23 : 157–160.
- Alvarado, Y., Layson, S., Menchaca, A., Moore, M., & Strong, N. (2019). Implementation of Enhanced Recovery After Surgery to Improve Surgical Outcomes in Women Requiring Cesareans. *J. Obstet. Gynecol. Neonatal Nurs.*, 48 : S24–S25.
- Amiruddin, A., Emilia, O., Prawitasari, S., & Prawirodihardjo, L. (2018). Hubungan Kepatuhan Tim Bedah dalam Penerapan Surgery Safety Checklist dengan Infeksi Luka Operasi dan Lama Rawat Inap pada Pasien Seksio Sesarea di Rumah Sakit Umum Daerah Kabupaten Barru. *J. Kesehat. Reproduksi*, 5: 145 - 148.
- Ammann, E.M., Kalsekar, I., Yoo, A., Scamuffa, R., Hsiao, C.W., Stokes, A.C., *et al.* (2019). Assessment of obesity prevalence and validity of obesity diagnoses coded in claims data for selected surgical populations: A retrospective, observational

study. *Medicine (Baltimore)*, 98 : e16438.

- Andersen, L.F., Wessel, I., Gotfredsen, K., & Øzhayat, E.B. (2020). Patients' and health professionals' experience of the Danish fast track treatment pathway for head and neck cancer patients receiving oral rehabilitation. *Acta Odontol. Scand.*, 78 : 362–369.
- Armbrust, R., Hinkson, L., Von Weizsäcker, K., & Henrich, W. (2016). The Charité cesarean birth: A family orientated approach of cesarean section. *J. Matern. Neonatal Med.*, 29: 163 - 168.
- Arroyo-Fernández, F.J., Calderón Seoane, J.E., & Torres Morera, L.M. (2020). Strategies of analgesic treatment after cesarean delivery. Current state and new alternatives. *Rev. Esp. Anesthesiol. Reanim.*, 67 : 167–175.
- Arzola, C., & Wiecezorek, P.M. (2011). Efficacy of low-dose bupivacaine in spinal anaesthesia for Caesarean delivery: Systematic review and meta-analysis. *Br. J. Anaesth.*, 107: 308 - 311.
- Ayeleke, R.O., Mourad, S., Marjoribanks, J., Calis, K.A., & Jordan, V. (2017). Antibiotic prophylaxis for elective hysterectomy. *Cochrane Database Syst. Rev.*,
- Azwar, S. (2013). Sikap Manusia: Teori dan Pengukurannya. *Yogyakarta: Pustaka Pelajar*.
- Azwar, S. (2011). Reliabilitas Dan Validitas. *Yogyakarta: Pustaka Pelajar*.
- Babazade, R., Vadhera, R.B., Krishnamurthy, P., Varma, A., Doulatram, G., Saade, G.R., *et al.* (2020). Acute postcesarean pain is associated with in-hospital exclusive breastfeeding, length of stay and post-partum depression. *J. Clin. Anesth.*, 62 : 109697.
- Baeriswyl, M., Kirkham, K.R., Kern, C., & Albrecht, E. (2015). The analgesic efficacy of ultrasound-guided transversus abdominis plane block in adult patients: A meta-analysis. *Anesth. Analg.*, 121: 1640 - 1654.
- Baluku, M., Bajunirwe, F., Ngonzi, J., Kiwanuka, J., & Ttendo, S. (2020). A randomized controlled trial of enhanced recovery after surgery versus standard of care recovery for emergency cesarean deliveries at Mbarara Hospital, Uganda. *Anesth. Analg.*, 130: 769–776.
- Beg, T., Daoud, B., & Angelo, T. (2021). Recent updates in obstetric analgesia. *Indian J. Pain*, 35: 186 - 194.
- Begum, T., Rahman, A., Nababan, H., Emdadul Hoque, D.M., Khan, A.F., Ali, T., *et al.* (2017). Indications and determinants of caesarean section delivery: Evidence from a population-based study in Matlab, Bangladesh. *PLoS One*, 12: e0188074.
- Bell, B.R., Bastien, P.E., & Douketis, J.D. (2015). Prevention of venous thromboembolism in the Enhanced Recovery After Surgery (ERAS) setting: an evidence-based review. *Can. J. Anesth.*, 62: 194.

- Bellei, E.A., Biduski, D., Cechetti, N.P., & De Marchi, A.C.B. (2018). Diabetes Mellitus m-Health Applications: A Systematic Review of Features and Fundamentals. *Telemed. e-Health.*, 24: 839 - 852.
- Bernard, H., & Foss, M. (2014). Patient experiences of enhanced recovery after surgery (ERAS). *Br. J. Nurs.*, 23 : 100–106.
- Bilku, D.K., Dennison, A.R., Hall, T.C., Metcalfe, M.S., & Garcea, G. (2014). Role of preoperative carbohydrate loading: A systematic review. *Ann. R. Coll. Surg. Engl.*, 96: 15 - 22.
- Blumenfeld, Y.J., El-Sayed, Y.Y., Lyell, D.J., Nelson, L.M., & Butwick, A.J. (2015). Risk Factors for Prolonged Postpartum Length of Stay Following Cesarean Delivery. *Am. J. Perinatol.*, 32 : 825–832.
- Bohiltea, R.-E., Turcan, N., & Cîrstoiu, M.M. (2019). The management of thromboembolic disease in pregnancy and puerperium. *Obstet. Ginecol.*, 67: 208 - 2012.
- Bohner, G., & Dickel, N. (2011). Attitudes and Attitude Change. *Annu. Rev. Psychol.*, 19: 257 - 276.
- Bollag, L., & Nelson, G. (2020). Enhanced Recovery After Cesarean (ERAC) – Beyond The Pain Scores. *Int. J. Obstet. Anesth.*, 43 : 36–38.
- Bollag, L., Tiouririne, M., Lim, G., Carvalho, B., Zakowski, M., Bhambhani, S., *et al.* (2019). Society of obstetric anesthesia and Perinatology (SOAP) enhanced recovery after cesarean (ERAC) consensus statement. *Anesth Analg.*, 132: e77.
- Bowden, S.J., Dooley, W., Hanrahan, J., Kanu, C., Halder, S., Cormack, C., *et al.* (2019). Fast-track pathway for elective caesarean section: a quality improvement initiative to promote day 1 discharge. *BMJ open Qual.*, 8 : e000465.
- Brown, D., & Xhaja, A. (2018). Nursing Perspectives on Enhanced Recovery After Surgery. *Surg. Clin. North Am.*, 98: 1211 - 1221.
- Caughey, Aaron B, Wood, S.L., Macones, G.A., Wrench, I.J., Huang, J., Norman, M., *et al.* (2018). Guidelines for intraoperative care in cesarean delivery: Enhanced Recovery After Surgery Society Recommendations (Part 2). *Am. J. Obstet. Gynecol.*, 219 : 533–544.
- Cawich, S.O., Mohammed, F., Spence, R., FaSiOen, P., & Naraynsingh, V. (2019). Surgeons' attitudes toward mechanical bowel preparation in the 21st century: A survey of the Caribbean College of Surgeons. *Curr. Med. Res. Pract.*, 9: 53 - 57.
- Çevik, S.A., & Başer, M. (2016). Effect of bed exercises and gum chewing on abdominal sounds, flatulence and early discharge in the early period after caesarean section. *J. Clin. Nurs.*, 25: 1416 - 1425.
- Chen, J., Zheng, D., Cui, H., Liu, S., Zhang, L., & Liu, C. (2018). Roles and mechanisms of TRPC3 and the PLCγ/PKC/CPI-17 signaling pathway in regulating parturition. *Mol. Med. Rep.*, 17: 898–910.

- Chen, P.S., & Lin, M.H. (2017). Development of simulation optimization methods for solving patient referral problems in the hospital-collaboration environment. *J. Biomed. Inform.*, 73: 148 - 158.
- Chhabra, P. (2014). Maternal near miss: An indicator for maternal health and maternal care. *Indian J. Community Med.* 39: 132 - 137.
- Ciardulli, A., Saccone, G., Di Mascio, D., Caissutti, C., & Berghella, V. (2018). Chewing gum improves postoperative recovery of gastrointestinal function after cesarean delivery: a systematic review and meta-analysis of randomized trials. *J. Matern. Neonatal Med.*, 31: 1924–1932.
- Cisse, B., Moore, L., Kuimi, B.L.B., Porgo, T.V., Boutin, A., Lavoie, A., *et al.* (2016). Impact of socio-economic status on unplanned readmission following injury: A multicenter cohort study. *Injury*, 47: 1083 - 1090.
- Cloninger, C.R., & Zohar, A.H. (2011). Personality and the perception of health and happiness. *J. Affect. Disord.*, 128: 24 - 32.
- Coates, E., Fuller, G., Hind, D., Wrench, I.J., Wilson, M.J., & Stephens, T. (2016). Enhanced recovery pathway for elective caesarean section. *Int. J. Obstet. Anesth.*, 27: 94 - 95.
- Cohen, R., & Gooberman-Hill, R. (2019). Staff experiences of enhanced recovery after surgery: Systematic review of qualitative studies. *BMJ Open*, 9: e022259
- Corniola, M. V., Meling, T.R., & Tessitore, E. (2020). Enhanced recovery after spine surgery—a multinational survey assessing surgeons’ perspectives. *Acta Neurochir.*, 162: 1371 - 1377
- Corso, E., Hind, D., Beever, D., Fuller, G., Wilson, M.J., Wrench, I.J., *et al.* (2017). Enhanced recovery after elective caesarean: A rapid review of clinical protocols, and an umbrella review of systematic reviews. *BMC Pregnancy Childbirth*, 17: 1-10
- Crano, W.D., & Prislin, R. (2011). Attitudes and attitude change, *Psychology Press*.
- Cusack, L., Digenis, C., Schultz, T., Klaer, B., & Hobbs, M. (2020). Women’s experiences with enhanced recovery after elective caesarean section with next day discharge: A qualitative study. *Midwifery*, 83: 102632.
- Dadlez, N.M., Brubaker, S.G., Simpson, L.L., Yilmaz, B., & Williams, I.A. (2014). Impact of change in delivery practice on neonatal and maternal outcomes in cases of significant congenital heart disease. *Congenit. Heart Dis.*, 9 : 368–372.
- Dahabreh, I.J., Steele, D.W., Shah, N., & Trikalinos, T.A. (2015). Oral mechanical bowel preparation for colorectal surgery: Systematic review and meta-analysis. *Dis. Colon Rectum*, 58: 698 - 707.
- Deniau, B., Bouhadjari, N., Faitot, V., Mortazavi, A., Kayem, G., Mandelbrot, L., *et al.* (2016). Evaluation of a continuous improvement programme of enhanced recovery after caesarean delivery under neuraxial anaesthesia. *Anaesth. Crit. Care Pain*

Med., 35 : 395–399.

- Dexter, F., Epstein, R.H., Dexter, E.U., Lubarsky, D.A., & Sun, E.C. (2017). Hospitals with briefer than average lengths of stays for common surgical procedures do not have greater odds of either re-admission or use of short-term care facilities. *Anaesth. Intensive Care*, 45 : 210–219.
- Didden, A.G.M., Punt, I.M., Feczko, P.Z., & Lenssen, A.F. (2019). Enhanced recovery in usual health care improves functional recovery after total knee arthroplasty. *Int. J. Orthop. Trauma Nurs.*, 34 : 9–15.
- Dodd, J.M., Anderson, E.R., Gates, S., & Grivell, R.M. (2014). Surgical techniques for uterine incision and uterine closure at the time of caesarean section. *Cochrane Database Syst. Rev.*, 7.
- Drury, C., & Hunter, J. (2016). The Hole in Holistic Patient Care. *Open J. Nurs.*, 6: 776 - 792.
- Echeverry-Marín, P.C., Rincón-Valenzuela, D.A., Monroy-Charry, A.G., Ruiz-Villa, J.O., Higuera-Redondo, G., & Rubio-Elorza, J.H. (2016). Survey about attitudes on perioperative temperature monitoring and thermal protection in Colombia. *Colomb. J. Anesthesiol.*, 44: 282 - 291.
- Edwards, D.A., Hedrick, T.L., Jayaram, J., Argoff, C., Gulur, P., Holubar, S.D., *et al.* (2019). American Society for Enhanced Recovery and Perioperative Quality Initiative Joint Consensus Statement on Perioperative Management of Patients on Preoperative Opioid Therapy. *Anesth. Analg.*, 129 : 553–566.
- Eignor, D.R. (2013). The standards for educational and psychological testing. Accessed from: psycnet.apa.org
- El-Achi, V., Wan, K.M., Brown, J., Marshall, D., & McGee, T. (2018). Readmissions for surgical site infections following caesarean section. *Aust. New Zeal. J. Obstet. Gynaecol.*, 58: 582 - 585.
- El-Kefraoui, C., Olleik, G., Chay, M.A., Kouyoumdjian, A., Nguyen-Powanda, P., Rajabiyazdi, F., *et al.* (2020). Opioid versus opioid-free analgesia after surgical discharge: Protocol for a systematic review and meta-analysis. *BMJ Open*, 10: e035443.
- Fawcett, W.J., & Ljungqvist, O. (2017). Starvation, carbohydrate loading, and outcome after major surgery. *BJA Educ.*, 17: 312 - 316.
- Fay, E.E., Hitti, J.E., Delgado, C.M., Savitsky, L.M., Mills, E.B., Slater, J.A.L., *et al.* (2019). An enhanced recovery after surgery pathway for cesarean delivery decreases hospital stay and cost. *American Journal of Obstetrics and Gynecology*, 221: 349.e1-349.e9.
- Fecher-Jones, I., & Taylor, C. (2015). Lived experience, enhanced recovery and laparoscopic colonic resection. *Br. J. Nurs.*, 24 : 223–228.
- Feizpour, C.A., Patel, M.S., Syed, M.A., Carrasco, A., Shah, J., Hanish, S., *et al.* (2021).

Enhanced recovery in liver transplantation: A value-based approach to complex surgical care. *Surgery*, 170: 1830 - 1837.

Fernández, R., del Sol Calderón, P., Izquierdo de la Puente, Á., Blanco Fernández, R., & Martín García, M. (2023). Psychiatric Comorbidity and Length of Stay in a general hospital. *Eur. Psychiatry*, 66: S588 - S589.

Fiore, J.F., Castelino, T., Pecorelli, N., Niculiseanu, P., Balvardi, S., Hershorn, O., *et al.* (2017). Ensuring Early Mobilization Within an Enhanced Recovery Program for Colorectal Surgery. *Ann. Surg.*, 4: 223 - 231.

Fischer, C., & Wick, E. (2020). An AHRQ national quality improvement project for implementation of enhanced recovery after surgery. *Semin. Colon Rectal Surg.*, 31: 100778.

Francis, N.K., Walker, T., Carter, F., Hübner, M., Balfour, A., Jakobsen, D.H., *et al.* (2018). Consensus on Training and Implementation of Enhanced Recovery After Surgery: A Delphi Study. *World J. Surg.*, 42: 1919 - 1928.

Gamez, B.H., & Habib, A.S. (2018). Predicting severity of acute pain after cesarean delivery: A narrative review. *Anesth. Analg.*, 126: 1606 - 1614.

Ganer Herman, H., Kleiner, I., Tairy, D., Gonen, N., Ben Zvi, M., Kovo, M., *et al.* (2020). Effect of Digital Step Counter Feedback on Mobility After Cesarean Delivery: A Randomized Controlled Trial. *Obstet. Gynecol.*, 135 : 1345–1352.

Giacomelli, E., Dorigo, W., Campolmi, M., Casini, A., Fargion, A., Bush, R.L., *et al.* (2021). A pilot study of the enhanced recovery after surgery protocol in aortic surgery. *J. Vasc. Surg.*, 74 : 90-96.

Gibbons, L., Belizán, J.M., Lauer, J. a, Betrán, A.P., Merialdi, M., & Althabe, F. (2010). The Global Numbers and Costs of Additionally Needed and Unnecessary Caesarean Sections Performed per Year: Overuse as a Barrier to Universal Coverage. *World Heal. Rep. Backgr. Pap.*, 30: 1 - 31.

Glanz, K., Rimer, B.K., & Viswanath, K. (2015). Health Behavior: Theory, Research, and Practice. *John Wiley & Sons*.

Gonçalves-Bradley, D.C., Lannin, N.A., Clemson, L.M., Cameron, I.D., & Shepperd, S. (2016). Discharge planning from hospital. *Cochrane Database Syst. Rev.*, 2.

Gotlib Conn, L., McKenzie, M., Pearsall, E.A., & McLeod, R.S. (2015). Successful implementation of an enhanced recovery after surgery programme for elective colorectal surgery: A process evaluation of champions' experiences. *Implement. Sci.*, 10: 1 - 11.

Grant, M.C., Hanna, A., Benson, A., Hobson, D., Wu, C.L., Yuan, C.T., *et al.* (2018). Dedicated Operating Room Teams and Clinical Outcomes in an Enhanced Recovery after Surgery Pathway for Colorectal Surgery. *J. Am. Coll. Surg.*, 226 : 267–276.

Grasch, J.L., Rojas, J.C., Sharifi, M., McLaughlin, M.M., Bhamidipalli, S.S., & Haas,

- D.M. (2023). Impact of Enhanced Recovery After Surgery pathway for cesarean delivery on postoperative pain. *AJOG Glob. Reports*, 3: 100169.
- Green, A., Heinink, T., & Banks, A. (2016). Enhanced recovery after caesarean section benefits patients and hospitals. *BJOG An Int. J. Obstet. Gynaecol.*, 123: 147 - 148.
- Guo, J. guo, Li, H. ling, Pei, Q. qing, & Feng, Z. ying (2018). The analgesic efficacy of subcostal transversus abdominis plane block with Mercedes incision. *BMC Anesthesiol.*, 18: 1 - 11.
- Gur, R., Duggal, S.D., Rongpharpi, S.R., Srivastava, R., Kumar, A., Gupta, V., *et al.* (2015). Post caesarean surgical site infections. *Arch. Clin. Microbiol.*, 6: 1 - 4.
- Guyton, A.C., & Hall, J.E. (2014). Guyton dan Hall Buku Ajar Fisiologi Kedokteran. Edisi 12., *Elsevier Jakarta : EGC*.
- Habib, A.S., & Ituk, U. (2018). Enhanced recovery after cesarean delivery [version 1; referees: 2 approved]. *F1000Research*.
- Hair, J.F., Black, W.C., Babin, B.J., & Anderson, R.E. (2010). Multivariate Data Analysis. *Vectors*.
- Harzif, A.K., Wicaksono, M.D., Kallista, A., Emeraldi, M., & Pratama, G. (2020). Overview of risk factor and bacterial pattern in patient with surgical site infection after caesarean section in Ciptomangunkusumo Hospital from 2016 to 2018. *Infect. Prev. Pract.*, 2: 100090.
- Hedderson, M., Lee, D., Hunt, E., Lee, K., Xu, F., Mustille, A., *et al.* (2019). Enhanced Recovery after Surgery to Change Process Measures and Reduce Opioid Use after Cesarean Delivery: A Quality Improvement Initiative. *Obstet. Gynecol.*. 134: 511–519.
- Heng, H., Jazayeri, D., Shaw, L., Kiegaldie, D., Hill, A.M., & Morris, M.E. (2020). Hospital falls prevention with patient education: A scoping review. *BMC Geriatr.*, 20: 1 - 12.
- Herbert, G., Sutton, E., Burden, S., Lewis, S., Thomas, S., Ness, A., *et al.* (2017). Healthcare professionals' views of the enhanced recovery after surgery programme: A qualitative investigation. *BMC Health Serv. Res.*, 17: 1-14.
- Herbert, K.A., Yurashevich, M., Fuller, M., Pedro, C.D., & Habib, A.S. (2020). Impact of a multimodal analgesic protocol modification on opioid consumption after cesarean delivery: a retrospective cohort study. *J. Matern. Neonatal Med.*, 35: 4743 - 4749.
- Ho, Y.M., Smith, S.R., Pockney, P., Lim, P., & Attia, J. (2014). A meta-analysis on the effect of sham feeding following colectomy: Should gum chewing be included in enhanced recovery after surgery protocols? *Dis. Colon Rectum*, 57: 115 - 126.
- Hoe, C.Y.W., Ahmad, B., & Watterson, J. (2023). The use of videos for diabetes patient education: A systematic review. *Diabetes. Metab. Res. Rev.*, 40: e3722.

- Huang, J., Cao, C., Nelson, G., & Wilson, R.D. (2019). A Review of Enhanced Recovery After Surgery Principles Used for Scheduled Caesarean Delivery. *J. Obstet. Gynaecol. Canada*, 41: 1775 - 1788.
- Hughes, M., Coolsen, M.M.E., Aahlin, E.K., Harrison, E.M., McNally, S.J., Dejong, C.H.C., *et al.* (2015). Attitudes of patients and care providers to enhanced recovery after surgery programs after major abdominal surgery. *J. Surg. Res.*, 193: 102 - 110.
- Iancu, G., Botezatu, R., Gică, N., Ciobanu, A., Gică, C., Mustață, L., *et al.* (2019). Enhanced recovery after surgery (ERAS) – preoperative and intraoperative recommendations for caesarean delivery. *Obstet. Ginecol.*, 67: 4.
- Ilyas, S., Simmons, S., & Bampoe, S. (2019). Systematic review of enhanced recovery protocols for elective caesarean section versus conventional care. *Aust. New Zeal. J. Obstet. Gynaecol.*, 59 : 767–776.
- Irwin, R., & Tan, T. (2020). Evaluation of the impact of enhanced recovery after surgery protocol implementation on maternal outcomes following elective caesarean delivery. *Int. J. Obstet. Anesth.*, 43: 17.
- Jacques, V., Vial, F., Lerintiu, M., Thilly, N., Mc Nelis, U., Raft, J., *et al.* (2013). Enhanced recovery following uncomplicated elective caesarean section in France: A survey of national practice. *Ann. Fr. Anesth. Reanim.*, 32: 142 - 148.
- Jarraya, A., Boujelbene, M., Zghal, J., Laabidi, S., Smaoui, M., & Kolsi, K. (2016). Réhabilitation précoce post césarienne: Enquête de pratique auprès des maternités de la région de sfax en Tunisie. *Pan Afr. Med., J.* 23: 12 - 15.
- Jasim, H.H., Sulaiman, S.A.B.S., Khan, A.H., & Rajah, U.A.S. (2017). Factors affecting post caesarean pain intensity among women in the northern peninsular of Malaysia. *J. Clin. Diagnostic Res.*, 11 : IC07-IC11.
- Jeff, A., & Taylor, C. (2014). Ward nurses' experience of enhanced recovery after surgery: A grounded theory approach. *Gastrointest. Nurs.*, 12: 23 - 31.
- Jeong, O., & Kim, H.G. (2019). Implementation of enhanced recovery after surgery (ERAS) program in perioperative management of gastric cancer surgery: A nationwide survey in korea. *J. Gastric Cancer.*, 19: 72 - 76.
- Jia, W.D. (2018). Multidisciplinary team building in enhanced recovery after surgery. *Zhonghua Wai Ke Za Zhi*, 56: 14 - 17.
- Jin, J., Peng, L., Chen, Q., Zhang, D., Ren, L., Qin, P., *et al.* (2016). Prevalence and risk factors for chronic pain following cesarean section: A prospective study. *BMC Anesthesiol.*, 16: 1 - 11.
- Joewono, H.T., Sulistyono, A., Haslinda, Y., & Aditiawarman (2020). Maternal and perinatal outcome of women with obesity in pregnancy. *Syst. Rev. Pharm.*, 11: 695 - 698.
- Joshi, R., Thomas, M., Patkar, S., Gupta, A.M., Panhale, K., Rane, P.P., *et al.* (2021).

- Impact of enhanced recovery pathway in 408 gallbladder cancer resections. *HPB*, 24: 47 - 56.
- Juanamasta, I.G., Aungsuroch, Y., Gunawan, J., Suniyadewi, N.W., & Nopita Wati, N.M. (2021). Holistic Care Management of Diabetes Mellitus: An Integrative Review. *Int. J. Prev. Med.*, 12: 69 - 74.
- Kadzielski, J., McCormick, F., Herndon, J.H., Rubash, H., & Ring, D. (2015). Surgeons' Attitudes Are Associated With Reoperation and Readmission Rates. *Clin. Orthop. Relat. Res.*, 473: 1544 - 1551.
- Kahokehr, A., Sammour, T., Zargar-Shoshtari, K., Thompson, L., & Hill, A.G. (2009). Implementation of ERAS and how to overcome the barriers. *Int. J. Surg.*, 473: 1544 - 1551.
- Kalogera, E., & Dowdy, S.C. (2016). Enhanced Recovery Pathway in Gynecologic Surgery: Improving Outcomes Through Evidence-Based Medicine. *Obstet. Gynecol. Clin. North Am.*, 43: 551 - 573.
- Kalsbeek, A., La Fleur, S., & Fliers, E. (2014). Circadian control of glucose metabolism. *Mol. Metab.*, 3: 372 - 383.
- Kaur, H., kaur, S., & sikka, P. (2015). A Quasi-Experimental Study To Assess The Effectiveness Of Early Ambulation In Post-Operative Recovery Among Post-Caesarean Mothers Admitted In Selected Areas Of Nehru Hospital,PGIMER,Chandigarh . *Nurs. Midwifery Res. Journal*, 11: 33 - 44.
- Kementerian Kesehatan RI Badan Penelitian dan Pengembangan (2018). RISKESDAS 2018. *Kementrian Kesehat. Republik Indones.*
- Kim, B.-Y., Park, K.-J., & Ryoo, S.-B. (2018). Effects of a Mobile Educational Program for Colorectal Cancer Patients Undergoing the Enhanced Recovery After Surgery. *Open Nurs. J.*, 12 : 142–154.
- Kinshella, M.L.W., Salimu, S., Chiwaya, B., Chikoti, F., Chirambo, L., Mwaungulu, E., *et al.* (2020). “So sometimes, it looks like it’s a neglected ward”: Health worker perspectives on implementing kangaroo mother care in southern Malawi. *PLoS One*, 15: e0243770.
- Kleiman, A.M., Chisholm, C.A., Dixon, A.J., Sariosek, B.M., Thiele, R.H., Hedrick, T.L., *et al.* (2020). Evaluation of the impact of enhanced recovery after surgery protocol implementation on maternal outcomes following elective cesarean delivery. *Int. J. Obstet. Anesth.*, 43 : 39–46.
- Koch, F., Green, M., Dietrich, M., Moikow, L., Pontau, F., Ulmer, S., *et al.* (2021). Perioperative management-tasks and significance of specially trained nursing staff. *Zentralblatt fur Chir. - Zeitschrift fur Allg. Visz. und Gefasschirurgie*, 146 : 260–268.
- Kripalani, S., Theobald, C.N., Anctil, B., & Vasilevskis, E.E. (2014). Reducing hospital readmission rates: Current strategies and future directions. *Annu. Rev. Med.*, 65:

471 - 485.

- Kruse, A.R., Lauszus, F.F., Forman, A., Kesmodel, U.S., Rugaard, M.B., Knudsen, R.K., *et al.* (2021). Effect of early discharge after planned cesarean section on recovery and parental sense of security. A randomized clinical trial. *Acta Obstet. Gynecol. Scand.*, 100: 955–963.
- Lambrechts, M.J., Boeyer, M.E., Tweedy, N.M., Gupta, S.K., Kimchi, E.T., & Hoernschemeyer, D.G. (2022). Team Integrated Enhanced Recovery (TIGER) Protocol after Adolescent Idiopathic Scoliosis Correction Lowers Direct Cost and Length of Stay While Increasing Daily Contribution Margins. *Mo. Med.*, 119: 152 - 159.
- Laronche, A., Popescu, L., & Benhamou, D. (2017). An enhanced recovery programme after caesarean delivery increases maternal satisfaction and improves maternal-neonatal bonding: A case control study. *Eur. J. Obstet. Gynecol. Reprod. Biol.*, 210: 212 - 216.
- Larson, D.W., Lovely, J.K., Welsh, J., Annaberdyev, S., Coffey, C., Corning, C., *et al.* (2018). A Collaborative for Implementation of an Evidence-Based Clinical Pathway for Enhanced Recovery in Colon and Rectal Surgery in an Affiliated Network of Healthcare Organizations. *Jt. Comm. J. Qual. Patient Saf.*, 44 : 204–211.
- Lavand’homme, P. (2018). Postoperative cesarean pain: real but is it preventable? *Curr. Opin. Anaesthesiol.*, 31 : 262–267.
- Ledari, F.M., Barat, S., & Delavar, M.A. (2012). Chewing gums has stimulatory effects on bowel function in patients undergoing cesarean section: A randomized controlled trial. *Bosn. J. Basic Med. Sci.*, 12: 265 - 171.
- Leffert, L., Butwick, A., Carvalho, B., Arendt, K., Bates, S.M., Friedman, A., *et al.* (2018). The Society for Obstetric Anesthesia and Perinatology consensus statement on the anesthetic management of pregnant and postpartum women receiving thromboprophylaxis or higher dose anticoagulants. *Anesth. Analg.*, 126: 928 - 944.
- Leibo, J.Z., & Poggio, T. (2018). Perception. *Living Machines: A Handbook of Research in Biomimetic and Biohybrid Systems*.
- Leslie, M.S., & Briggs, L.A. (2016). Preeclampsia and the Risk of Future Vascular Disease and Mortality: A Review. *J. Midwifery Women’s Heal.*, 61: 315 - 324.
- Lester, S.A., Kim, B., Tubinis, M.D., Morgan, C.J., & Powell, M.F. (2020). Impact of an enhanced recovery program for cesarean delivery on postoperative opioid use. *Int. J. Obstet. Anesth.*, 43 : 47–55.
- Li, X., Tang, K., Cui, C., & Huang, P. (2022). Nutrition protocol implemented in ERAS of hypopharyngeal cancer: a single center nutrition protocol in China. *Int. J. Food Prop.*, 25: 1343 - 1350.
- Liabsuetrakul, T., Chongsuvivatwong, V., Lumbiganon, P., & Lindmark, G. (2003).

Obstetricians' attitudes, subjective norms, perceived controls, and intentions on antibiotic prophylaxis in caesarean section. *Soc. Sci. Med.*, 57: 1665 - 1674.

- Lisby, M., Klingenberg, M., Ahrensberg, J.M., Hoeyem, P.H., & Kirkegaard, H. (2019). Clinical impact of a comprehensive nurse-led discharge intervention on patients being discharged home from an acute medical unit: Randomised controlled trial. *Int. J. Nurs. Stud.*, 100: 330 - 339.
- Liu, B., Liu, S., Wang, Y., Zhao, B., Zhao, T., Zhao, L., *et al.* (2019). Neurosurgical enhanced recovery after surgery (ERAS) programme for elective craniotomies: Are patients satisfied with their experiences? A quantitative and qualitative analysis. *BMJ Open*, 9: e028706.
- Liu, Z.Q., Du, W.J., & Yao, S.L. (2020). Enhanced recovery after cesarean delivery: a challenge for anesthesiologists. *Chin. Med. J. (Engl.)*, 133 : 590–596.
- Lof, S., Benedetti Cacciaguerra, A., Aljarrah, R., Okorocha, C., Jaber, B., Shamali, A., *et al.* (2020). Implementation of enhanced recovery after surgery for pancreatoduodenectomy increases the proportion of patients achieving textbook outcome: A retrospective cohort study. *Pancreatology*, 20 : 976–983.
- Lovely, J.K., Hyland, S.J., Smith, A.N., Nelson, G., Ljungqvist, O., & Parrish, R.H. (2019). Clinical pharmacist perspectives for optimizing pharmacotherapy within Enhanced Recovery After Surgery (ERAS®) programs. *Int. J. Surg.*, 63 : 58–62.
- Lyon, A., Solomon, M.J., & Harrison, J.D. (2014). A qualitative study assessing the barriers to implementation of enhanced recovery after surgery. *World J. Surg.*, 38: 1374 - 1380
- Macones, G.A., Caughey, A.B., Wood, S.L., Wrench, I.J., Huang, J., Norman, M., *et al.* (2019). Guidelines for postoperative care in cesarean delivery: Enhanced Recovery After Surgery (ERAS) Society recommendations (part 3). *Am. J. Obstet. Gynecol.*, 221: 247.e1-247.e9.
- Mai, L.J. and J.L.J. (2015). PONV Attitudes, Knowledge and Antiemetic Prescribing Practices Amongst Surgeons in a Paediatric Hospital. *JMHH*, 3 : 9–25.
- Malbrain, M.L.N.G., Langer, T., Annane, D., Gattinoni, L., Elbers, P., Hahn, R.G., *et al.* (2020). Intravenous fluid therapy in the perioperative and critical care setting: Executive summary of the International Fluid Academy (IFA). *Ann. Intensive Care*, 10: 1 - 19.
- Mangesi, L., & Hofmeyr, G.J. (2002). Early compared with delayed oral fluids and food after caesarean section. *Cochrane Database Syst. Rev.*, 1.
- Mann, J., & Rosenthal, T.L. (1969). Vicarious and direct counterconditioning of test anxiety through individual and group desensitization. *Behav. Res. Ther.*, &: 359 - 367.
- Marroquin, B., Feng, C., Balofsky, A., Edwards, K., Iqbal, A., Kanel, J., *et al.* (2017). Neuraxial opioids for post-cesarean delivery analgesia: can hydromorphone

replace morphine? A retrospective study Hydromorphone and cesarean section. *Int. J. Obstet. Anesth.*, 30: 16 - 22.

- Marsanto, A.E., Kuntaman, K., Paraton, H., & Prasetyo, B. (2019). Pattern of disease and type of operation of Surgical Site Infection in obstetrics and gynecology at Dr Soetomo Hospital, Surabaya, Indonesia. *Maj. Obstet. Ginekol.*, 27: 49 - 55.
- Martin, D., Roulin, D., Grass, F., Addor, V., Ljungqvist, O., Demartines, N., *et al.* (2018). A multicentre qualitative study assessing implementation of an Enhanced Recovery After Surgery program. *Clin. Nutr.*, 37: 2172 - 2177.
- Masini, L., De Luca, C., Noia, G., Caruso, A., Lanzone, A., Rendeli, C., *et al.* (2019). Prenatal diagnosis, natural history, postnatal treatment and outcome of 222 cases of spina bifida: experience of a tertiary center. *Ultrasound Obstet. Gynecol.*, 53 : 302–308.
- Mathiesen, O., Wetterslev, J., Kontinen, V.K., Pommergaard, H.C., Nikolajsen, L., Rosenberg, J., *et al.* (2014). Adverse effects of perioperative paracetamol, NSAIDs, glucocorticoids, gabapentinoids and their combinations: A topical review. *Acta Anaesthesiol. Scand.*, 58: 1182 - 1198.
- Matthews, K.C., White, R.S., Ewing, J., Abramovitz, S.E., & Kalish, R.B. (2020). Enhanced recovery after surgery for cesarean delivery. *Am. J. Obstet. Gynecol.*, 222: S396–S397.
- Mboya, I.B., Mahande, M.J., Obure, J., & Mwambi, H.G. (2020). Predictors of perinatal death in the presence of missing data: A birth registry-based study in northern Tanzania. *PLoS One*, 15: e0231636.
- Mc Loughlin, S., Terrasa, S.A., Ljungqvist, O., Sanchez, G., Garcia Fornari, G., & Alvarez, A.O. (2019). Nausea and vomiting in a colorectal ERAS program: Impact on nutritional recovery and the length of hospital stay. *Clin. Nutr., ESPEN* 34: 73–80.
- McDonald, S., & Bott, A. (2020). Surgical site infections. *Surg. (United Kingdom)*, 38: 150 - 154.
- McGinn, B., & Zhou, J. (2015). Anesthesia for cesarean delivery. *Surg. (United Kingdom)*, 40: 180 - 186.
- McLemore, E.C., Paige, J.T., Bergman, S., Hori, Y., Schwarz, E., & Farrell, T.M. (2015). Ongoing evolution of practice gaps in gastrointestinal and endoscopic surgery: 2014 report from the Society of American Gastrointestinal and Endoscopic Surgeons (SAGES) Continuing Education Committee. *Surg. Endosc.*, 29: 3017–3029.
- McLeod, R.S., Aarts, M.A., Chung, F., Eskicioglu, C., Forbes, S., Conn, L.G., *et al.* (2015). Development of an enhanced recovery after surgery guideline and implementation strategy based on the knowledge-to-action cycle. *Ann. Surg.*, 262: 1016 - 1025.

- Mei, B., Wang, W., Cui, F., Wen, Z., & Shen, M. (2017). Chewing Gum for Intestinal Function Recovery after Colorectal Cancer Surgery: A Systematic Review and Meta-Analysis. *Gastroenterol. Res. Pract.*, 84: 212 - 220.
- Mendes, D.I.A., Ferrito, C.R. de A.C., & Gonçalves, M.I.R. (2018). Nursing Interventions in the Enhanced Recovery After Surgery®: Scoping Review. *Rev. Bras. Enferm.*, 71: 2824 - 2832.
- Michard, F., Gan, T.J., & Kehlet, H. (2017). Digital innovations and emerging technologies for enhanced recovery programmes. *Br. J. Anaesth.*, 119: 31 - 39.
- Miller, T.E., Roche, A.M., & Mythen, M. (2016). Fluid management and goal-directed therapy as an adjunct to Enhanced Recovery After Surgery (ERAS). *Can. J. Anesth.*, 62: 158 - 163.
- Mohammed, S.A., Khalis Jawad, A., & Jamal Murad, S. (2020). Enhanced Recovery Program after Cesarean Section in Sulaimani Maternity Teaching Hospital. *Kurdistan J. Appl. Res.*, 5: 287–293.
- Moussa, H., Hosseini Nasab, S., Fournie, D., Ontiveros, A., Alkawas, R., Chauhan, S., et al. (2019). The impact of time of delivery on gestations complicated by preterm premature rupture of membranes: daytime versus nighttime. *J. Matern. Neonatal Med.*, 32: 3319–3324.
- Narava, S., Pokhriyal, S.C., Singh, S.B., Barpanda, S., & Bricker, L. (2020). Outcome of multiple cesarean sections in a tertiary maternity hospital in the United Arab Emirates: A retrospective analysis. *Eur. J. Obstet. Gynecol. Reprod. Biol.*, 247: 143 - 148.
- Narendra, S., Singh, D., Prakash, J., Priye, S., Kumar, S., & Bharati, B. (2022). Comparison of ramosetron, ondansetron, and granisetron in attenuating maternal hypotension after spinal anesthesia in patients undergoing lower segment caesarean section: A prospective, randomized, double-blinded study. *Bali J. Anesthesiol.*, 6: 32 - 37.
- Nelson, G., & Macones, G. (2021). Despite barriers, Enhanced Recovery After Surgery for cesarean delivery should be prioritized in developing countries. *Am. J. Obstet. Gynecol.*, 224: 554–555.
- Nelson, Gregg, & Wilson, R.D. (2020). Enhanced recovery after cesarean delivery: is protocol compliance the missing link? *Am. J. Obstet. Gynecol.*, 223: 608 - 615.
- Nevalainen, J., Skarp, S., Savolainen, E.R., Ryyänen, M., & Järvenpää, J. (2017). Intrauterine growth restriction and placental gene expression in severe preeclampsia, comparing early-onset and late-onset forms. *J. Perinat. Med.*, 45: 869–877.
- Noh, J.J., Kim, M.-S., & Lee, Y.-Y. (2021). The implementation of enhanced recovery after surgery protocols in ovarian malignancy surgery. *Gland Surg.*, 10: 1182–1194.

- Notoatmodjo, S. (2012). *Promosi Kesehatan dan Perilaku Kesehatan (edisi revisi 2012)*, Jakarta: Rineka Cipta.
- O'Neill, B.J., Dwyer, T., Reid-Searl, K., & Parkinson, L. (2017). Managing the deteriorating nursing home resident after the introduction of a hospital avoidance programme: a nursing perspective. *Scand. J. Caring Sci.*, 31: 312–322.
- Olson, K.A., Fleming, R.Y.D., Fox, A.W., Grimes, A.E., Mohiuddin, S.S., Robertson, H.T., *et al.* (2021). The Enhanced Recovery After Surgery (ERAS) Elements that Most Greatly Impact Length of Stay and Readmission. *Am. Surg.*, 87: 473 - 479.
- Östman, L., Näsman, Y., Eriksson, K., & Nyström, L. (2019). Ethos: The heart of ethics and health. *Nurs. Ethics.*, 26: 26 - 36.
- Pan, J., Hei, Z., Li, L., Zhu, D., Hou, H., Wu, H., *et al.* (2020). The advantage of implementation of enhanced recovery after surgery (ERAS) in acute pain management during elective cesarean delivery: A prospective randomized controlled trial. *Ther. Clin. Risk Manag.*, 16: 369–378.
- Panda, S., Begley, C., & Daly, D. (2016). Readmission following caesarean section: Outcomes for women in an Irish maternity hospital. *Br. J. Midwifery.*, 24: 322 - 328.
- Pandey, D., Mehta, S., Grover, A., & Goel, N. (2015). Indwelling catheterization in caesarean section: Time to retire it! *J. Clin. Diagnostic Res.*, 9: QC01 - QC04.
- Patel, K., & Zakowski, M. (2021). Enhanced Recovery After Cesarean: Current and Emerging Trends. *Curr. Anesthesiol. Rep.*, 11: 136–144.
- Peahl, A.F., Smith, R., Johnson, T.R.B., Morgan, D.M., & Pearlman, M.D. (2019). Better late than never: why obstetricians must implement enhanced recovery after cesarean. *Am. J. Obstet. Gynecol.*, 221: 117.e1-117.e7.
- Pereira Gomes Morais, E., Riera, R., Porfírio, G.J.M., Macedo, C.R., Sarmiento Vasconcelos, V., de Souza Pedrosa, A., *et al.* (2016). Chewing gum for enhancing early recovery of bowel function after caesarean section. *Cochrane Database Syst. Rev.*, 10.
- Pereira, S.L., da Silva, T.P.R., Moreira, A.D., Novaes, T.G., Pessoa, M.C., Matozinhos, I.P., *et al.* (2019). Factors associated with the length of hospital stay of women undergoing cesarean section. *Rev. Saude Publica.*, 53: 65 - 71.
- Persico, M., Miller, D., Way, C., Williamson, M., O'Keefe, K., Strnatko, D., *et al.* (2019). Implementation of Enhanced Recovery After Surgery in a Community Hospital: An Evidence-Based Approach. *J. Perianesthesia Nurs.*, 34: 188–197.
- Pervez, S., Sharawi, N., Blake, L., Habib, A.S., Brookfield, K.F., & Carvalho, B. (2021). Impact of enhanced recovery after caesarean delivery on maternal outcomes: A systematic review and meta-analysis. *Anaesth. Crit. Care Pain Med.*, 40: 100935.
- Petty, R.E. (2012). *Attitude Change*. Encyclopedia of Human Behavior: Second Edition. Academic Press.

- Philp, S., Carter, J., Barnett, C., D'Abrew, N., Pather, S., & White, K. (2015). Patients' Perspectives of Fast-Track Surgery and the Role of the Fast-Track Clinical Nurse Consultant in Gynecological Oncology. *Holist. Nurs. Pract.*, 29: 158–166.
- Pujic, B., Kendrisic, M., Shotwell, M., Shi, Y., & Baysinger, C.L. (2018). Survey of enhanced recovery after surgery protocols for cesarean delivery in Serbia. *Front. Med.*, 5: 100 - 117.
- Pujić, B., Vejnović, T., Jovanović, L., Andjelić, N., Vejnović, A., & Palmer, C. (2022). Implementation Of ERAS Protocol for Cesarean Section. *Acta Clin. Croat.*, 61: 151 - 159.
- Qiang, H., & Yuanshui, S. (2020). Enhanced Recovery after Surgery (ERAS) at cesarean delivery to reduce postoperative length of stay: some details need to be improved. *Am. J. Obstet. Gynecol.*, 222: 512 - 518.
- Rajabpour, S., Rayyani, M., & Mangolian Shahrabaki, P. (2019). The relationship between Iranian patients' perception of holistic care and satisfaction with nursing care. *BMC Nurs.*, 18: 1 - 7.
- Richard, B., Reilly, A., Lyden, E., & Garvin, K. (2018). Is obesity a risk factor for extended length of stay and readmission after total hip arthroplasty? *Ann. Jt.*, 3: 87 - 93.
- Ricotta, C., Cintorino, D., Pagano, D., Bonsignore, P., Piazza, S., di Francesco, F., *et al.* (2019). Enhanced Recovery after Implementation of Surgery Protocol in Living Kidney Donors: The ISMETT Experience. *Transplant. Proc.*, 51: 2910–2913.
- Rifa'i, A.S., Sari, G.M., & Budipramana, V.S. (2020). Pre-Operative Enhanced Recovery After Surgery (ERAS) Protocol Compliance Towards Major Surgery Patients at Dr. Soetomo General Hospital, Surabaya. *Biomol. Heal. Sci. J.*, 3: 28 - 32
- Roberts, V. (2018). *Impact of birth experience on bonding and perceived attachment: A qualitative study of the subjective experience of birth*. California: Aliant Internal University.
- Rosyidah, R., Dewanto, A., Hapsari, E.D., & Widyastuti, Y. (2022). Health Professionals Perception of Enhanced Recovery After Surgery: A Scoping Review. *J. Perianesthesia Nurs.*, 37: 956 - 960.
- Rosyidah, R., Widyastuti, Y., Dewanto, A., Hapsari, E.D., & Wicaksana, A.L. (2021). The Attitude of Health Care Workers on Enhanced Recovery After Surgery for Cesarean Delivery: A Scoping Review., *J. Obstet. Gynaecol. Canada*. 43: 856 - 863.
- Rousseau, A., Sadoun, M., Aimé, I., Leguen, M., Carbonnel, M., & Ayoubi, J.M. (2017). Comparative study about enhanced recovery after cesarean section: What benefits, what risks? *Gynecol. Obstet. Fertil. Senol.*, 45: 387–392.
- Rubin, D.J., Maliakkal, N., Zhao, H., & Miller, E.E. (2023). Hospital Readmission Risk

and Risk Factors of People with a Primary or Secondary Discharge Diagnosis of Diabetes. *J. Clin. Med.*, 12: 1274 - 1284.

Rubio, D.M., Berg-weger, M., Tebb, S.S., Lee, E.S., & Rauch, S. (2003). Objectifying Content Validity: In Social Work Research. *Soc. Work Res.*, 27: 94 - 104.

Saad, A.F., Rahman, M., Costantine, M.M., & Saade, G.R. (2014). Blunt versus sharp uterine incision expansion during low transverse cesarean delivery: A metaanalysis. *Am. J. Obstet. Gynecol.*, 211: 684.e1 - 684.e11.

Salenger, R., Holmes, S.D., Rea, A., Yeh, J., Knott, K., Born, R., *et al.* (2021). Cardiac Enhanced Recovery After Surgery: Early Outcomes in a Community Setting. *Ann. Thorac. Surg.*, 113: 2008 - 2017.

Sales, I., Babelghaith, S.D., Wajid, S., Mahmoud, M.A., Alsaleh, S.S., Alfadly, S., *et al.* (2018). Impact of diabetes continuing education on health care professionals' attitudes towards diabetes care in a Yemeni city. *Trop. J. Pharm. Res.*, 17: 143 - 149.

Sato, H., Carvalho, G., Sato, T., Lattermann, R., Matsukawa, T., & Schricker, T. (2010). The association of preoperative glycemic control, intraoperative insulin sensitivity, and outcomes after cardiac surgery. *J. Clin. Endocrinol. Metab.*, 95: 4338 - 4334.

Schricker, T., & Lattermann, R. (2015). Perioperative catabolism. *Can. J. Anesth.*, 62: 182 - 190.

Seah, X.Y., Tham, X.C., Kamaruzaman, N.R., & Yobas, P. (Klainin) (2018). Nurses' perception of knowledge, attitudes and reported practice towards patients with eating disorders: A concurrent mixed-methods study. *Arch. Psychiatr. Nurs.*, 32: 133-151.

Sepúlveda, F.A.L., Jiménez, L.M.M., Castro, H.D.A., Castro, J.A.P., Cañas, E.M.A., Hidalgo, J.E.O., *et al.* (2021). Predictors for intraoperative requirement of inotropic and/or vasopresor support in miocardial revascularization surgery. *Rev. Chil. Anest.*, 50: 344 - 351.

Seraphim, I.B., Ravi, V., & Rajagopal, A. (2020). Prediction of diabetes readmission using machine learning. *Int. J. Adv. Sci. Technol.*, 29: 9 - 16.

Shinnick, J.K., Ruhotina, M., Has, P., Kelly, B.J., Brousseau, E.C., O'Brien, J., *et al.* (2020). Enhanced Recovery after Surgery for Cesarean Delivery Decreases Length of Hospital Stay and Opioid Consumption: A Quality Improvement Initiative. *Am. J. Perinatol.*, 38: e215 - e223.

Short, H.L., Taylor, N., Thakore, M., Piper, K., Baxter, K., Heiss, K.F., *et al.* (2018). A survey of pediatric surgeons' practices with enhanced recovery after children's surgery. *J. Pediatr. Surg.*, 53: 418 - 430.

Simsek, T., Uzelli Simsek, H., & Canturk, N.Z. (2014). Response to trauma and metabolic changes: posttraumatic metabolism. *Turkish J. Surg.*, 30: 153 - 160.

Slade, M., & Longden, E. (2015). Empirical evidence about recovery and mental health.

BMC Psychiatry, 15: 1 - 14.

Spandonaro, F., Mancusi, R.L., Terranova, L., Giannarelli, D., Grossi, P., Imberti, D., *et al.* (2013). Prevention of VTE in orthopedic surgery patients. *Farmeconomia. Heal. Econ. Ther. pathways.*, 14: 1 - 36.

Squeo, G.C., Sarosiek, B., & McConnell, G. (2021). Patient education is an essential component of an Enhanced Recovery Pathway in colon and rectal surgery. *Semin. Colon Rectal Surg.*, 100829.

Stephen, G., Mgongo, M., Hussein Hashim, T., Katanga, J., Stray-Pedersen, B., & Msuya, S.E. (2018). Anaemia in Pregnancy: Prevalence, Risk Factors, and Adverse Perinatal Outcomes in Northern Tanzania. *Anemia*, 2018: 1846280

Suharwardy, S., & Carvalho, B. (2020). Enhanced recovery after surgery for cesarean delivery. *Curr. Opin. Obstet. Gynecol.*, 32: 113 - 120.

Sultan, P., Sharawi, N., Blake, L., & Carvalho, B. (2020). Enhanced recovery after caesarean delivery versus standard care studies: a systematic review of interventions and outcomes. *Int. J. Obstet. Anesth.*, 43: 72 - 78.

Symonds, N.E., Vidler, M., Wiens, M.O., Omar, S., English, L.L., Ukah, U.V., *et al.* (2023). Risk factors for postpartum maternal mortality and hospital readmission in low- and middle-income countries: a systematic review. *BMC Pregnancy Childbirth*, 23: 303 - 311.

Tamang, T., Wangchuk, T., Zangmo, C., Wangmo, T., & Tshomo, K. (2021). The successful implementation of the Enhanced Recovery After Surgery (ERAS) program among caesarean deliveries in Bhutan to reduce the postoperative length of hospital stay. *BMC Pregnancy Childbirth*, 21: 1 - 7.

Taylor, C.R., Dominguez, J.E., & Habib, A.S. (2019). Obesity and obstetric anesthesia: Current insights. *Local Reg. Anesth*, 28: 111 - 124.

Tazreean, R., Nelson, G., & Twomey, R. (2022). Early mobilization in enhanced recovery after surgery pathways: Current evidence and recent advancements. *J. Comp. Eff. Res*, 11: 121 - 129.

Teigen, N.C., Sahasrabudhe, N., Doulaveris, G., Xie, X., Negassa, A., Bernstein, J., *et al.* (2020). Enhanced recovery after surgery at cesarean delivery to reduce postoperative length of stay: a randomized controlled trial. *Am. J. Obstet. Gynecol.*, 372: e1-372.e10.

Thiele, R.H., Rea, K.M., Turrentine, F.E., Friel, C.M., Hassinger, T.E., Goudreau, B.J., *et al.* (2015). Standardization of care: Impact of an enhanced recovery protocol on length of stay, complications, and direct costs after colorectal surgery. *Journal of the American College of Surgeons*, 220: 430 - 443.

Thorneloe, B., Carvalho, J.C.A., Downey, K., & Balki, M. (2019). Uterotonic drug usage in Canada: a snapshot of the practice in obstetric units of university-affiliated hospitals. *Int. J. Obstet. Anesth.*, 37: 45 - 51.

- Tran, H.T., Nippita, T., Nguyen, P.T.K., Nguyen, P.T.T., Huynh, T.T.D., Hua, O.T. Le, *et al.* (2018). Knowledge, experience and attitudes towards skin-to-skin contact following Caesarean sections among health professionals in Vietnam. *Acta Paediatr. Int. J. Paediatr.*, 107: 1917 - 1923.
- Vallejo, M.C., Attaallah, A.F., Elzamzamy, O.M., Cifarelli, D.T., Phelps, A.L., Hobbs, G.R., *et al.* (2017). An open-label randomized controlled clinical trial for comparison of continuous phenylephrine versus norepinephrine infusion in prevention of spinal hypotension during cesarean delivery. *Int. J. Obstet. Anesth.*, 29: 18 - 25.
- Wen, T., Yu, V.X., Wright, J.D., Goffman, D., Attenello, F., Mack, W.J., *et al.* (2020). Postpartum length of stay and risk for readmission among women with preeclampsia. *J. Matern. Neonatal Med.*, 33: 1086 - 1094.
- Weng, X. sheng, Liu, J., & Wu, D. (2020). Chinese Survey on Enhanced Recovery after Surgery and Thromboprophylaxis Following Arthroplasty. *Orthop. Surg.*, 12: 900–906.
- Wickenbergh, E., Nilsson, L., Bladh, M., Kjølhed, P., & Wodlin, N.B. (2020). Agreements on perceived use of principles for Enhanced Recovery After Surgery between patients and nursing staff in a gynecological ward. *Eur. J. Obstet. Gynecol. Reprod. Biol.*, 250: 216–223.
- Williams, J.B., McConnell, G., Allender, J.E., Woltz, P., Kane, K., Smith, P.K., *et al.* (2019). One-year results from the first US-based enhanced recovery after cardiac surgery (ERAS Cardiac) program. *J. Thorac. Cardiovasc. Surg.*, 157: 1881–1888.
- Wilson, R. Douglas, Caughey, A.B., Wood, S.L., Macones, G.A., Wrench, I.J., Huang, J., *et al.* (2018). Guidelines for Antenatal and Preoperative care in Cesarean Delivery: Enhanced Recovery After Surgery Society Recommendations (Part 1). *Am. J. Obstet. Gynecol.*, 219: 523.e1-523.e15.
- Witcher, A., Axley, J., Novak, Z., Laygo-Prickett, M., Guthrie, M., Xhaja, A., *et al.* (2021). Implementation of an enhanced recovery program for lower extremity bypass. *J. Vasc. Surg.*, 73 : 554–563.
- Wrench, I.J., Allison, A., Galimberti, A., Radley, S., & Wilson, M.J. (2015). Introduction of enhanced recovery for elective caesarean section enabling next day discharge: A tertiary centre experience. *Int. J. Obstet. Anesth.*, 24: 124–130.
- Wulandari, E.S., Ernawati, E., & Nuswantoro, D. (2021). Risk Factors of Preeclampsia With Severe Features and Its Complications. *Indones. Midwifery Heal. Sci., J.* 5: 29–37.
- Wyse, R., Smith, S., Zucca, A., Fakes, K., Mansfield, E., Johnston, S.A., *et al.* (2023). Effectiveness and cost-effectiveness of a digital health intervention to support patients with colorectal cancer prepare for and recover from surgery: study protocol of the RecoverEsupport randomised controlled trial. *BMJ Open*, 13: e067150.
- Yurashevich, M., Chow, A., Kowalczyk, J.J., Traynor, A.J., & Carvalho, B. (2019).

Preoperative fasting times for patients undergoing caesarean delivery: Before and after a patient educational initiative. *Turkish J. Anaesthesiol. Reanim.*, 47 : 282–286.

Zorrilla-Vaca, A., Stone, A.B., Ripolles-Melchor, J., Abad-Motos, A., Ramirez-Rodriguez, J.M., Galan-Menendez, P., *et al.* (2021). Institutional factors associated with adherence to enhanced recovery protocols for colorectal surgery: Secondary analysis of a multicenter study. *J. Clin. Anesth.*, 74: 110378.

Zuckerman, R.B., Sheingold, S.H., Orav, E.J., Ruhter, J., & Epstein, A.M. (2016). Readmissions, Observation, and the Hospital Readmissions Reduction Program. *N. Engl. J. Med.*, 374: 1543 - 1551.