

- Ahmadian, E., Zununi Vahed, S., Mammadova, S., & Abediazar, S. (2021). Immunosuppressant Management in Renal Transplant Patients with COVID-19. *BioMed Research International*, 2021, 9318725. <https://doi.org/10.1155/2021/9318725>
- Çankaya, E., Bilen, Y., Keles, M., Uyanik, A., Bilen, N., & Aydınli, B. (2015). Neutrophil-Lymphocyte Ratio Is Significantly Decreased in Preemptive Renal Transplant Patients. *Transplantation Proceedings*, 47(5), 1364–1368. <https://doi.org/10.1016/j.transproceed.2015.04.052>
- Dragun, D., Hoff, U., Park, J. K., Qun, Y., Schneider, W., Luft, F. C., & Haller, H. (2001). Prolonged cold preservation augments vascular injury independent of renal transplant immunogenicity and function. *Kidney International*, 60(3), 1173–1181. <https://doi.org/10.1046/j.1523-1755.2001.0600031173.x>
- Ergin, G., Değer, M., Köprü, B., Derici, Ü., & Arınsoy, T. (2019). High neutrophil to lymphocyte ratio predicts acute allograft rejection in kidney transplantation; a retrospective study. *Turkish Journal of Medical Sciences*, 49(2), 525–530. <https://doi.org/10.3906/sag-1811-41>
- Gallego Valcarce, E., Ortega Cerrato, A., Llamas Fuentes, F., Masiá Mondéjar, J., Martínez Fernández, G., López Rubio, E., López Montes, A., Pérez Martínez, J., Martínez Villaescusa, M., & Gómez Roldán, C. (2009). El tiempo de isquemia fría corto optimiza los resultados de los trasplantes renales efectuados con donantes con criterios expandidos. *Nefrología (Madrid)*, 29(5), 456–463.
- Garcia, G. G., Harden, P., & Chapman, J. (2013). The global role of kidney transplantation. *Nephrology, Dialysis, Transplantation : Official Publication of the European Dialysis and Transplant Association - European Renal Association*, 28(8), e1-5. <https://doi.org/10.1093/ndt/gfs013>
- Halazun, K. J., Marangoni, G., Hakeem, A., Fraser, S. M., Farid, S. G., & Ahmad, N. (2013). Elevated preoperative recipient neutrophil-lymphocyte ratio is associated with delayed graft function following kidney transplantation. *Transplantation Proceedings*, 45(9), 3254–3257. <https://doi.org/10.1016/j.transproceed.2013.07.065>
- Jha, V. (2004). End-stage renal care in developing countries: The India experience. *Renal Failure*, 26(3), 201–208. <https://doi.org/10.1081/jdi-120039516>

Shorter Cold Ischemia Time in Deceased Donor Kidney Transplantation Reduces the Incidence of Delayed Graft Function Especially Among Highly Sensitized Patients and Kidneys From Older Donors. *Transplantation Proceedings*, 52(1), 42–49. <https://doi.org/10.1016/j.transproceed.2019.11.025>

Naik, R. H., & Shawar, S. H. (2022). Renal Transplantation Rejection. In *StatPearls [Internet]*. StatPearls Publishing.

Naranjo, M., Agrawal, A., Goyal, A., & Rangaswami, J. (2018). Neutrophil-to-Lymphocyte Ratio and Platelet-to-Lymphocyte Ratio Predict Acute Cellular Rejection in the Kidney Allograft. *Annals of Transplantation*, 23, 467–474. <https://doi.org/10.12659/AOT.909251>

Nath, J., Hodson, J., Canbilien, S. W., Al Shakarchi, J., Inston, N. G., Sharif, A., & Ready, A. R. (2016). Effect of cold ischaemia time on outcome after living donor renal transplantation. *The British Journal of Surgery*, 103(9), 1230–1236. <https://doi.org/10.1002/bjs.10165>

Novianti, P. A., Duarsa, G. W. K., Aditya, G. A., Oka, A. A. G., Santosa, K. B., Yudiana, I. W., Tirtayasa, P. M. W., Pramana, I. B. P., Kandarini, Y., Sudana, W., Sidartha, D., & Widiana, R. (2020). Correlation of total ischemic time to creatinine serum level and resistive index value in kidney transplant. *Neurologico Spinale Medico Chirurgico*, 3(2).

O’Connell, P. J., Brown, M., Chan, T. M., Claire-Del Granado, R., Davies, S. J., Eiam-Ong, S., Hassan, M. H., Kalantar-Zadeh, K., Levin, A., Martin, D. E., Muller, E., Ossareh, S., Tchokhanelidze, I., Trask, M., Twahir, A., Were, A. J. O., Yang, C.-W., Zemchenkov, A., & Harden, P. N. (2020). The role of kidney transplantation as a component of integrated care for chronic kidney disease. *Kidney International Supplements*, 10(1), e78–e85. <https://doi.org/10.1016/j.kisu.2019.11.006>

Ponticelli, C. E. (2015). The impact of cold ischemia time on renal transplant outcome. *Kidney International*, 87(2), 272–275. <https://doi.org/10.1038/ki.2014.359>

Redfield, R. R., Scalea, J. R., Zens, T. J., Muth, B., Kaufman, D. B., Djamali, A., Astor, B. C., & Mohamed, M. (2016). Predictors and outcomes of delayed graft function after living-donor kidney transplantation. *Transplant International: Official Journal of the*

<https://doi.org/10.1111/tri.12696>

- van de Laar, S. C., Lafranca, J. A., Minnee, R. C., Papalois, V., & Dor, F. J. M. F. (2022). The Impact of Cold Ischaemia Time on Outcomes of Living Donor Kidney Transplantation: A Systematic Review and Meta-Analysis. *Journal of Clinical Medicine*, 11(6). <https://doi.org/10.3390/jcm11061620>
- Wang, X.-W., Guo, R.-D., Ma, J.-G., Wang, Y.-W., & Zou, X.-F. (2022). Effects of prolonged cold ischemia on the DCD kidney function and Inflammasome expression in rat kidney transplants. *Transplant Immunology*, 74, 101511. <https://doi.org/10.1016/j.trim.2021.101511>
- Zhang Y, Liu R, Zhao X, Ou Z, Wang S, Wang D, Huang K, Pan S, Wu Y. Dynamic changes of neutrophil-to-lymphocyte ratio in brain-dead donors and delayed graft function in kidney transplant recipients. *Ren Fail.* 2022 Dec;44(1):1897-1903. doi: 10.1080/0886022X.2022.2141646. PMID: 36346017; PMCID: PMC9648373.