



DAFTAR PUSTAKA

- Ahmed, D. M., Chen, J. M., & Sanders, D. A. R. (2022). Pyrazole and Triazole Derivatives as *Mycobacterium Tuberculosis* UDP-Galactopyranose Inhibitors. In *Pharmaceuticals*. <https://doi.org/10.3390/ph15020197>
- Ahmed, M., Ramadan, W., Rambhu, D., & Shakeel, F. (2008). Potential of nanoemulsions for intravenous delivery of rifampicin. *Die Pharmazie*, 63(11), 806–811
- Akhtar, N., Ahad, A., Khar, R. K., Jaggi, M., Aqil, M., Iqbal, Z., Ahmad, F. J., & Talegaonkar, S. (2011). The emerging role of P-glycoprotein inhibitors in drug delivery: A patent review. In *Expert Opinion on Therapeutic Patents* (Vol. 21, Issue 4, pp. 561–576). <https://doi.org/10.1517/13543776.2011.561784>
- Al-Ali, A. A. A., Nielsen, R. B., Steffansen, B., Holm, R., & Nielsen, C. U. (2019). Nonionic surfactants modulate the transport activity of ATP-binding cassette (ABC) transporters and solute carriers (SLC): Relevance to oral drug absorption. *International Journal of Pharmaceutics*, 566, 410-433.
- Annisa, R. (2023). The Development Formulation of Eleutherine Palmifolia Extract-Loaded Self Nanoemulsifying Drug Delivery System (Snedds) Using D-Optimal Mixture Design Approach. *International Journal of Applied Pharmaceutics*, 269–276. <https://doi.org/10.22159/ijap.2023v15i5.47645>
- Anwer, M. K., Iqbal, M., Aldawsari, M. F., Alalaiwe, A., Ahmed, M. M., Muharram, M. M., Ezzeldin, E., Mahmoud, M. A., Imam, F., & Ali, R. (2021). Improved antimicrobial activity and oral bioavailability of delafloxacin by self-nanoemulsifying drug delivery sistem (SNEDDS). *Journal of Drug Delivery Science and Technology*, 64, 102572. <https://doi.org/10.1016/J.JDDST.2021.102572>
- Astuti, I.Y., Marchaban, Martien, R., & Nugroho, A.E., 2017, Validation of UV-VIS-Vis Spectrophotometric Method for Soluility Study of Pentagamavunon-0 in Self Nanoemulsifying Drug Delivery Sistem Vehicles, *Pharmacy*, 14:127-136.
- Azeem, A., Rizwan, M., Ahmad, F. J., Iqbal, Z., Khar, R. K., Aqil, M., & Talegaonkar, S. (2009). Nanoemulsion components screening and selection: A technical note. *AAPS PharmSciTech*, 10(1), 69–76. <https://doi.org/10.1208/s12249-008-9178-x>
- Badran M. Formulation and in vitro evaluation of flufenamic acid loaded deformable liposomes for improved skin delivery. *Dig. J. Nanomater. Biostruct.* 2014;9:83–91.
- Bali, V., Ali, M., & Ali, J. (2010). Study of surfactant combinations and development of a novel nanoemulsion for minimising variations in bioavailability of ezetimibe. *Colloids and Surfaces. B, Biointerfaces*, 76(2), 410–420. doi.org/10.1016/j.colsurfb.2009.11.021
- Bansal, T., Akhtar, N., Jaggi, M., et al. (2009). Novel formulation approaches for optimising delivery of anticancer drugs based on p-glycoprotein modulation. *Drug Discov Today*, 14, 21-32.



- Becher, P. (1965). Emulsions: Theory and Practice, 3rd Edition. Industrial & Engineering Chemistry, 57(11), 37–38.
- Benetton, S. A., Kedor-Hackmann, E. R. M., Santoro, M. I. R. M., & Borges, V. M. (1998). Visible spectrophotometric and first-derivative UV spectrophotometric determination of rifampicin and isoniazid in pharmaceutical preparations. In *Talanta* (Vol. 47).
- Bhagwat, D. A., Swami, P. A., Nadaf, S. J., Choudhari, P. B., Kumbar, V. M., More, H. N., Killedar, S. G., & Kawtikwar, P. S. (2021). Capsaicin Loaded Solid SNEDDS for Enhanced Bioavailability and Anticancer Activity: In-Vitro, In-Silico, and In-Vivo Characterization. *Journal of Pharmaceutical Sciences*, 110(1), 280–291. <https://doi.org/10.1016/j.xphs.2020.10.020>
- Bhatt, S., Gour, A., Singh, G., Nandi, U. (2020). Tuberculosis. In: Rayees, S., Din, I., Singh, G., Malik, F. (eds) *Chronic Lung Diseases*. Springer, Singapore. https://doi.org/10.1007/978-981-15-3734-9_5
- Boehme, C. C., Nicol, M. P., Nabeta, P., Michael, J. S., Gotuzzo, E., Tahirli, R., Gler, M. T., Blakemore, R., Worodria, W., Gray, C., Huang, L., Caceres, T., Mehdiyev, R., Raymond, L., Whitelaw, A., Sagadevan, K., ... Perkins, M. D. (2011). Feasibility, diagnostic accuracy, and effectiveness of decentralised use of the Xpert MTB/RIF test for diagnosis of tuberculosis and multidrug resistance: a multicentre implementation study. *Lancet*, 377, 1495–1505. <https://doi.org/10.1016/S0140>
- Borah Slater, K., Kim, D., Chand, P., Xu, Y., Shaikh, H., & Undale, V. (2023). A Current Perspective on the Potential of Nanomedicine for Anti-Tuberculosis Therapy. In *Tropical Medicine and Infectious Disease* (Vol. 8, Issue 2). MDPI. <https://doi.org/10.3390/tropicalmed8020100>
- Borenstein, M., Hedges, L. V., Higgins, J. P. T., & Rothstein, H. R. (2021). Introduction to meta-analysis (2nd ed.). Chichester, UK: John Wiley and sons. [Chapter 17]
- Buya, M. G. (2020). Self-nanoemulsifying drug delivery sistem (SNEDDS) for enhancing oral bioavailability of poorly soluble drugs. *Journal of Drug Delivery Science and Technology*, 55, 101373.
- Campanerut, P.A.Z., Ghiraldi, L.D., Sposito, F.L.E., Sato, D.N., Leite, C.Q.F., Hirata, M.H., Hirata, R.D.C., Cardoso, R.F. (2011). Rapid detection of resistance to pyrazinamide in *Mycobacterium tuberculosis* using the resazurin microtitre assay. *J Antimicrob Chemother*, 66, 1044-1046.
- Cerpnjak, K., Zvonar, A., Gašperlin, M., & Vrečer, F. (2013). Lipid-Based Systems as a Promising Approach for Enhancing the Bioavailability of Poorly Water-Soluble Drugs. *Acta Pharmaceutica*, 63, 427–445.
- Chan, E. D., & Iseman, M. D. (2008). Multidrug-resistant and extensively drug-resistant tuberculosis: A review. In *Current Opinion in Infectious Diseases* (Vol. 21, Issue 6, pp. 587–595). <https://doi.org/10.1097/QCO.0b013e328319bce6>
- Chen, M. L. (2008). Lipid Excipients and Delivery Systems for Pharmaceutical Development: A Regulatory Perspective. *Advanced Drug Delivery Reviews*, 60, 768–777.



- Chiolero, A., Santschi, V., Burnand, B., Platt, R. W., & Paradis, G. (2012). Meta-analyses: With confidence or prediction intervals? In *European Journal of Epidemiology* (Vol. 27, Issue 10, pp. 823–825). <https://doi.org/10.1007/s10654-012-9738-y>
- Cole, S. T., Brosch, R., Parkhill, J., Garnier, T., Churcher, C., Harris, D., Gordon, S. V., Eiglmeier, K., Gas, S., Barry, C. E., 3rd, Tekaiia, F., Badcock, K., Basham, D., Brown, D., Chillingworth, T., Connor, R., Davies, R., Devlin, K., Feltwell, T., Gentles, S., ... Barrell, B. G. (1998). Deciphering the biology of *Mycobacterium tuberculosis* from the complete genome sequence. *Nature*, 393(6685), 537–544. <https://doi.org/10.1038/31159>
- Danaei, M., Dehghankhold, M., Ataei, S., Hasanzadeh Davarani, F., Javanmard, R., Dokhani, A., Khorasani, S., & Mozafari, M. R. (2018). Impact of particle size and polydispersity index on the clinical applications of lipidic nanocarrier systems. In *Pharmaceutics* (Vol. 10, Issue 2). MDPI AG. <https://doi.org/10.3390/pharmaceutics10020057>
- Danimayostu, A. A., Martien, R., Lukitaningsih, E., Danarti, R., & Bulaksumur, G. M. (2022). Formulation Of Topical Self-Nanoemulsifying Drug Delivery Systems (SNEDDS) Of Vitamin D3. *Jurnal Ilmu Kesehatan*, 11(2). <https://doi.org/10.37341/interest.v0i0.532>
- Desavathu, M., & Prameela Rani, A. (n.d.). Formulation and Evaluation of Self-Emulsifying Drug Delivery Systems of Rosuvastatin Calcium. In *Asian Journal of Pharmaceutics* (Vol. 15, Issue 1).
- Djekic, L., & Primorac, M. (2008). The influence of cosurfactants and oils on the formation of pharmaceutical microemulsions based on PEG-8 caprylic/capric glycerides. *International Journal of Pharmaceutics*, 352(1-2), 231-239.
- Eid, A.M.M., Elmarzugi, N.A., dan El-Enshasy, H.A., 2013. Preparation and Evaluation Of Olive Oil Nanoemulsion Using Sucrose Monoester. *International Journal of Pharmacy and Pharmaceutical Sciences*, 5: 434–440.
- El-Badry, M., Haq, N., Fetih, G., & Shakeel, F. (2014). Measurement and Correlation of Tadalafil Solubility in Five Pure Solvents at (298.15 to 333.15) K. *Journal of Chemical & Engineering Data*, 59(3), 839–843. <https://doi.org/10.1021/je400982r>
- Elmeliegy, M., Vourvahis, M., Guo, C., & Wang, D. D. (2020). Effect of P-glycoprotein (P-gp) Inducers on Exposure of P-gp Substrates: Review of Clinical Drug–Drug Interaction Studies. In *Clinical Pharmacokinetics* (Vol. 59, Issue 6, pp. 699–714). Adis. <https://doi.org/10.1007/s40262-020-00867-1>
- Gallo, G. G., & Radaelli, P. (1976). Anal. Profiles of Drug Subs. In K. Florey (Ed.), *Anal. Profiles of Drug Subs.* (Vol. 5, p. 467). Academic Press.
- Gohel, M., Purohit, A., Patel, A., & Hingorani, L. (2016). Optimization of bacoside a loaded snedds using d-optimal mixture design for enhancement insolubility and bioavailability. *International Journal of Pharmacy and Pharmaceutical Sciences*, 8(12), 213–220. <https://doi.org/10.22159/ijpps.2016v8i12.13488>
- Grumezescu, A., 2016. *Nanobiomaterials in Cancer Therapy - 1st Edition*. Elsevier, Amsterdam.



- Gupta, S., Kesarla, R., & Omri, A. (2013). Formulation Strategies to Improve the Bioavailability of Poorly Absorbed Drugs with Special Emphasis on Self-Emulsifying Systems. *ISRN Pharm*, 2013.
- Gupta, P., Garg, T., Tanmay, M., & Arora, S. (2015). Polymeric Drug-Delivery Systems: Role in P-gp Efflux System Inhibition. In *Critical ReviewsTM in Therapeutic Drug Carrier Systems* (Vol. 32, Issue 3). www.begellhouse.com
- Gupta, A., Eral, H. B., Hatton, T. A., & Doyle, P. S. (2016). Nanoemulsions: Formation, properties and applications. In *Soft Matter* (Vol. 12, Issue 11, pp. 2826–2841). Royal Society of Chemistry. <https://doi.org/10.1039/c5sm02958a>
- Gursoy, R. N., Benita, S., & Levy, R. H. (2003). Enhanced oral absorption of paclitaxel in a novel self-microemulsifying drug delivery system with or without concomitant use of P-glycoprotein inhibitors. *Pharmaceutical Research*, 20(8), 1284-1289.
- Hauss, D. J. (2007). Oral lipid-based formulations. *Advanced Drug Delivery Reviews*, 59(7), 667-676.
- Holm, R., Porter, C. J., & Edwards, G. A. (2002). Destabilization of colloidal lipid particles by digestion of triglycerides: implications for drug delivery and bioavailability. *Pharmaceutical Research*, 19(11), 1671-1676.
- Hugger, E. D., Novak, B. L., Burton, P. S., et al. (2002). A comparison of commonly used polyethoxylated pharmaceutical excipients on their ability to inhibit P-glycoprotein activity in vitro. *Journal of Pharmaceutical Sciences*, 91, 1991-2002.
- Hussain, A., Singh, S. K., Singh, N., & Prasad Verma, P. R. (2016). In vitro - In vivo - In silico simulation studies of anti-tubercular drugs doped with a self nanoemulsifying drug delivery system. *RSC Advances*, 6(95), 93147–93161. <https://doi.org/10.1039/c6ra14122f>
- Jaworska, M., Sikora, E., & Ogonowski, J. (2014). The influence of glicerides oil phase on O/W nanoemulsion formation by pic method, *Periodica Polytechnica Chemical Engineering*, vol. 58, Supplement, pp. 43–48.
- Juliantoni, Y., Hajrin, W., dan Subaidah, W.A., 2020. Nanoparticle Formula Optimization of Juwet Seeds Extract (*Syzygium cumini*) using Simplex Lattice Design Method. *Jurnal Biologi Tropis*, 20: 416–422.
- Kalepu, S., Manthina, M. & Padavala, V. (2013). Oral lipid-based drug delivery systems - an overview, *Acta Pharmaceutica Sinica B*, vol. 3, no. 6, pp. 361–372.
- Kasim, N. A., Whitehouse, M., Ramachandran, C., Bermejo, M., Lennernäs, H., Hussain, A. S., Junginger, H. E., Stavchansky, S. A., Midha, K. K., Shah, V. P., & Amidon, G. L. (2004). Molecular properties of WHO essential drugs and provisional biopharmaceutical classification. *Molecular Pharmaceutics*, 1(1), 85–96. <https://doi.org/10.1021/mp034006h>
- Kassem, M. G. A., Ahmed, A.-M. M., Abdel-Rahman, H. H. & Moustafa, A. H. E. (2019). Use of Span 80 and Tween 80 for blending gasoline and alcohol in spark ignition engines. *Energy Reports*, vol. 5, pp. 221–230.
- Katalinić-Janković, V., Furci, L., & Cirillo, D. M. (2012). Microbiology of mycobacterium tuberculosis and a new diagnostic test for TB. *European*



- Respiratory Monograph, 58, 8–13.
<https://doi.org/10.1183/1025448x.10022311>
- Katawera, V., Siedner, M., Ii, Y.B. (2014). Evaluation of the modified colorimetric resazurin microtiter plate-based antibacterial assay for rapid and reliable tuberculosis drug susceptibility testing. *BMC Microbiol*, 14(259), 1-4.
- Kawamura, K., Yamasaki, T., whoYui, J., et al. (2009). In vivo evaluation of p-glycoprotein and breast cancer resistance protein modulation in the brain using [11C] gefitinib. *Nucl Med Biol*, 36, 239-246.
- Kemenkes. (2020). *Pedoman Nasional Pelayanan Kedokteran : Tata Laksana Tuberkulosis*.
- Khandelia, R., Jaiswal, A., Ghosh, S. S., & Chattopadhyay, A. (2013). Gold nanoparticle-protein agglomerates as versatile nanocarriers for drug delivery. *Small*, 9(20), 3494–3505. <https://doi.org/10.1002/sml.201203095>
- Khoo, S. M., Ramachandran, G., & Lee, V. H. (2003). Lipid-based formulations for oral delivery of lipophilic drugs: clinical implications. *Expert Opinion on Drug Delivery*, 1(1), 99-114.
- Kusumorini, N., Nugroho, A., Pramono, S., & Martien, R. (2022). Spray-dried self-nanoemulsifying drug delivery systems as carriers for the oral delivery of piperine: Characterization and in vitro evaluation. *Journal of Applied Pharmaceutical Science*, 12(9), 89-96.
<https://doi.org/10.7324/japs.2022.120906>
- Li, L., Zhou, C. H., & Xu, Z. P. (2019). Self-Nanoemulsifying Drug-Delivery Sistem. In *Nanocarriers for Drug Delivery* (pp. 421-449). Elsevier.
- Listyorini, N.M.D., Wijayanti, N.L.P.D., dan Astuti, K.W., 2018. Optimasi Pembuatan Nanoemulsi Virgin Coconut Oil. *Jurnal Kimia*, 8–12.
- Luna, J.A.C. (2004). *A Tuberculosis Guide for Specialist Physicians*. International Union Against Tuberculosis and Lung Disease (IUATLD).
- Mahdi, E. S., Sakeena, M. H. F., Abdulkarim, M. F., Abdullah, G. Z., Sattar, M. A., & Noor, A. M. (2011). Effect of surfactant and surfactant blends on *pseudoternary* phase diagram behavior of newly synthesized palm kernel oil esters. *Drug Design, Development and Therapy*, 5, 311-323.
- Mahdi, L., Sudibyoy, R. S., & Martien, R. (2020). Self-Nano Emulsifying Drug Delivery System (SNEDDS) of Curcuma mangga Val. Essential Oil and The Stability Study. In *RESEARCH ARTICLE Indonesian Journal of Pharmacy Indonesian J Pharm* (Vol. 31, Issue 4).
- Maher, S., & Brayden, D. J. (2012). Overcoming Poor Permeability: Translating Permeation Enhancers for Oral Peptide Delivery. *Drug Discovery Today: Technologies*, 9, 113–119.
- Maison, D. P. (2022). Tuberculosis pathophysiology and anti-VEGF intervention. *Journal of Clinical Tuberculosis and Other Mycobacterial Diseases* (Vol. 27). Elsevier Ltd. <https://doi.org/10.1016/j.jctube.2022.100300>
- Majani, A.V., & Paradkar, M.U., 2018, Development and Validation of UV-VIS Method for the Estimation of Ceftriaxone Sodium in Nanoparticles, *Der Pharma Chemica*, 10: 151-157.



- Makadia, H. A., Bhatt, A. Y., Parmar, R. B., Paun, J. S., & Tank, H. M. (2013). Self-nano emulsifying drug delivery system (SNEDDS): future aspects. *Asian J. Pharm. Res*, 3(1), 21–24.
- Martin, A., Paasch, F., Docx, S., Fissette, K., Imperiale, B., Ribon, W., Gonzalez, L.A., Werngren, J., Engstrom, A., Skenders, G., Jureen, P., Hoffner, S., Portillo, P., Morcillo, N., Palomino, J.C. (2011). Multicentre laboratory validation of the colorimetric redox indicator (CRI) assay for the rapid detection of extensively drug-resistant (XDR) *Mycobacterium tuberculosis*. *J Antimicrob Chemother*, 66, 827-833.
- Mehta, S. K., Kaur, G., & Bhasin, K. K. (2007). Analysis of Tween based microemulsion in the presence of TB drug rifampicin. *Colloids and Surfaces B: Biointerfaces*, 60(1), 95–104. <https://doi.org/10.1016/j.colsurfb.2007.06.012>
- Mehta, S. K., Kaur, G., & Bhasin, K. K. (2008). Incorporation of antitubercular drug isoniazid in pharmaceutically accepted microemulsion: Effect on microstructure and physical parameters. *Pharmaceutical Research*, 25(1), 227–236. <https://doi.org/10.1007/s11095-007-9355-8>
- Mehta, S. K., Kaur, G., & Bhasin, K. K. (2010). Tween-embedded microemulsions-physicochemical and spectroscopic analysis for antitubercular drugs. *AAPS PharmSciTech*, 11(1), 143–153. <https://doi.org/10.1208/s12249-009-9356-5>
- Memvanga, P. B., Coco, R., & Pr at, V. (2013). An Oral Malaria Therapy: Curcumin-Loaded Lipid-Based Drug Delivery Systems Combined with β -Arteeether. *Journal of Controlled Release*, 172, 904–913.
- Moffat, A. C., et al. (Eds.). (1986). Clarke’s Isolation and Identification of Drugs (2nd ed., p. 60). The Pharmaceutical Press.
- Mountfield, R. J., Senepin, S., Schleimer, M., Walter, I., & Bittner, B. (2000). Potential Inhibitory Effects of Formulation Ingredients on Intestinal Cytochrome P450. *International Journal of Pharmaceutics*, 211, 89-92.
- Munita, J. M., & Arias, C. A. (2016). Mechanisms of Antibiotic Resistance. *Microbiology Spectrum*, 4(2). <https://doi.org/10.1128/microbiolspec.VMBF-0016-2015>
- Nanjwade, B. K., Patel, D. J., Udhani, R. A., Manvi, F. V., & Magadum, C. S. (2011). Self-nanoemulsifying drug delivery sistem of ezetimibe: formulation development and in-vitro characterization. *Current Drug Delivery*, 8(4), 438-446.
- Nasiruddin, M., Neyaz, Md. K., & Das, S. (2017). Nanotechnology-Based Approach in Tuberculosis Treatment. *Tuberculosis Research and Treatment*, 2017, 1–12. <https://doi.org/10.1155/2017/4920209>
- Nasr, A. M., Gardouh, A., & Ghorab, M. M. (2016). Novel Solid Self-Nanoemulsifying Drug Delivery System (S-Snedds) for Oral Delivery of Olmesartan Medoxomil: Design, Formulation, Pharmacokinetic and Bioavailability Evaluation. *Pharmaceutics*, 8(3), 20. <https://doi.org/10.3390/pharmaceutics8030020>
- Nguyen, T. T. L., Duong, V. A., & Maeng, H. J. (2021). Pharmaceutical formulations with p-glycoprotein inhibitory effect as promising approaches for enhancing oral drug absorption and bioavailability. In *Pharmaceutics*



- (Vol. 13, Issue 7). MDPI AG.
<https://doi.org/10.3390/pharmaceutics13071103>
- Palomino, J. C., Martin, A., Camacho, M., Guerra, H., Swings, J., & Portaels, F. (2002). Resazurin microtiter assay plate: Simple and inexpensive method for detection of drug resistance in *Mycobacterium tuberculosis*. *Antimicrobial Agents and Chemotherapy*, 46(8), 2720–2722. <https://doi.org/10.1128/AAC.46.8.2720-2722.2002>
- Patel, J., Kevin, G., Patel, A., Raval, M., & Sheth, N. (2011). Design and development of a self-nanoemulsifying drug delivery system for telmisartan for oral drug delivery. *International Journal of Pharmaceutical Investigation*, 1(2), 112–118. <https://doi.org/10.4103/2230-973X.82431>
- Poinard, B., Kamaluddin, S., Tan, A.Q.Q., Neoh, K.G., Kah, J.C.Y., 2019. Polydopamine coating enhances mucopenetration and cell uptake of nanoparticles. *ACS Appl. Mater. Interfaces* 11, 4777–4789.
- Pouton, C. W. (1997). Formulation of self-emulsifying drug delivery systems. *Advanced Drug Delivery Reviews*, 25(1), 47-58.
- Pouton, C. W., & Porter, C. J. H. (2008). Formulation of Lipid-Based Delivery Systems for Oral Administration: Materials, Methods and Strategies. *Advanced Drug Delivery Reviews*, 60, 625–637.
- Pudyastuti, B., & Wijaya, T. H. (2022). The effect of surfactant on the solubility of kencur rhizome ethanol extract in self-nanoemulsifying drug delivery system. *Pharmaciana*, 12(3), 319. <https://doi.org/10.12928/pharmaciana.v12i3.18881>
- Qian, J., Meng, H., Xin, L., Xia, M., Shen, H., Li, G., & Xie, Y. (2017). Self-Nanoemulsifying Drug Delivery Systems of Myricetin: Formulation Development, Characterization, and in Vitro and in Vivo Evaluation. *Colloids and Surfaces B: Biointerfaces*, 160, 101–109.
- Rahman, M. A., Hussain, A., Hussain, M. S., Mirza, M. A., & Iqbal, Z. (2013). Role of excipients in successful development of self-emulsifying/microemulsifying drug delivery system (SEDDS/SMEDDS). In *Drug Development and Industrial Pharmacy* (Vol. 39, Issue 1, pp. 1–19). <https://doi.org/10.3109/03639045.2012.660949>
- Rahman, M. A., Khoo, S. M., & Billa, N. (2013). Formulation and optimization of self-nanoemulsifying drug delivery sistem of all-trans-retinol acetate using Box-Behnken design. *International Journal of Nanomedicine*, 8, 103-113.
- Rakhmawatie, M. D., Wibawa, T., Lisdiyanti, P., Pratiwi, W. R., & Mustofa. (2019). Evaluation of crystal violet decolorization assay and resazurin microplate assay for antimycobacterial screening. *Heliyon*, 5(8). <https://doi.org/10.1016/j.heliyon.2019.e02263>
- Rancoita, P. M., Cugnata, F., Cruz, A. L. G., Borroni, E., Hoosdally, S., Walker, T. M., Grazian, C., Davies, T., Peto, T. E. A., Crook, D. W., Fowler, P., & Cirillo, D. M. (2018). Validating a 14-Drug Microtitre Plate Containing Bedaquiline and Delamanid for Large-Scale Research Susceptibility Testing Of *Mycobacterium Tuberculosis*. <https://doi.org/10.1101/244731>
- Rathod, S., Desai, H., Patil, R., & Sarolia, J. (2022). Non-ionic Surfactants as a P-Glycoprotein(P-gp) Efflux Inhibitor for Optimal Drug Delivery—A Concise Outlook. In *AAPS PharmSciTech* (Vol. 23, Issue 1). Springer Science and



- Business Media Deutschland GmbH. <https://doi.org/10.1208/s12249-022-02211-1>
- Rege, B. D., Kao, J. P. Y., & Polli, J. E. (2002). Effects of Nonionic Surfactants on Membrane Transporters in Caco-2 Cell Monolayers. *European Journal of Pharmaceutical Sciences*, 16, 237-246.
- Reynolds, James E.F. (1996). *Martindale The Extra Pharmacopoeia* (31st ed., p. 268). Royal Pharmaceutical Society.
- Russo, E., Selmin, F., Baldassari, S., Gennari, C.G.M., Caviglioli, G., Cilurzo, F., Minghetti, P., Parodi, B., 2016. A focus on mucoadhesive polymers and their application in buccal dosage forms. *J. Drug Deliv. Sci. Technol.* 32, 113–125.
- Sailor, G. U. (2021). Self-Nanoemulsifying Drug Delivery Systems (SNEDDS): An Innovative Approach to Improve Oral Bioavailability. In N. Shah (Ed.), *Nanocarriers: Drug Delivery System: An Evidence Based Approach* (pp. 255–280). Springer Singapore. https://doi.org/10.1007/978-981-33-4497-6_10
- Sandri, G., Rossi, S., Ferrari, F., Bonferoni, M.C., Zerrouk, N., Caramella, C., 2010. Mucoadhesive and penetration enhancement properties of three grades of hyaluronic acid using porcine buccal and vaginal tissue, Caco-2 cell lines, and rat jejunum. *J. Pharm. Pharmacol.* 56, 1083–1090.
- Schon, T., Miotto, P., Koser, C., Viveiros, M., Bottger, E., Cambau, E. (2017). *Mycobacterium tuberculosis* drug-resistance testing: challenges, recent developments, and perspectives. *Clin Microbiol Infect*, 23, 154-160.
- Sek L, Porter CJ, KaukonenAM et al (2002) Evaluation of the in-vitro digestion profiles of long and medium chain glycerides and the phase behaviour of their lipolytic products. *J Pharm Pharmacol* 54(1):29–4.
- Shakeel, F., Haq, N., & Alanazi, F. K. (2014). Measurement and Correlation of Solubility of Olmesartan Medoxomil in Six Green Solvents at 295.15–330.15 K. *Industrial & Engineering Chemistry Research*, 53(7), 2846–2849. <https://doi.org/10.1021/ie404373n>
- Singh, B., Bandopadhyay, S., Kapil, R., Singh, R., & Katare, O. P. (2009). Self-Emulsifying Drug Delivery Systems (SEDDS): Formulation Development, Characterization, and Applications. *Critical Reviews in Therapeutic Drug Carrier Systems*, 26, 427–521.
- Sonawale, P., Patil, A., Kamble, A., & Bhutkar, M. (2016). Solubility enhancement of lipophilic drugs-solid self micro-emulsifying drug delivery system. *Asian Journal of Pharmacy and Technology*, 6(3), 155.
- Syukri, Y., Nugroho, B. H., & Istanti, I. (2020). Penggunaan D-Optimal Mixture Design untuk Optimasi dan Formulasi Self-Nano Emulsifying Drug Delivery System (SNEEDS) Asam Mefenamat. *Jurnal Sains Farmasi & Klinis*, 7(3), 180. <https://doi.org/10.25077/jsfk.7.3.180-187.2020>
- Thakkar, H. P., Khunt, A., Dhande, R. D., & Patel, A. A. (2015). Formulation and evaluation of Itraconazole nanoemulsion for enhanced oral bioavailability. *Journal of Microencapsulation*, 32(6), 559–569. <https://doi.org/10.3109/02652048.2015.1065917>
- Wang, G., Tang, J., Feng, J., Wang, D., Huo, X., Lü, H., Wang, C., Lu, W., Wang, X., Chen, H., & Tan, C. (2021). Activity of Oritavancin and Its Synergy With Other Antibiotics Against *Mycobacterium Abscessus* Infection in Vitro and



- in Vivo. In International Journal of Molecular Sciences. <https://doi.org/10.3390/ijms22126346>
- Wang, X., Jiang, Y., Wang, Y., Huang, M., & Ye, Z. (2009). A self-nanoemulsifying drug delivery system for oral delivery of Lactoferrin: formulation optimization, microstructure, and cellular uptake in Caco-2 cells. *International Journal of Nanomedicine*, 4, 313–322.
- Webster, D., Lee, T.D.G., Moore, J., Manning, T., Kunimoto, D., Leblanc, D., Johnson, J.A., Gray, C.A. (2010). Antimycobacterial screening of traditional medicinal plants using the microplate resazurin assay. *Can J Microbiol*, 56, 487-494.
- WHO. (2022). Global Tuberculosis report 2022. <http://apps.who.int/bookorders>.
- Yin, Y. M., Cui, F. D., Mu, C. F., Choi, M. K., Kim, J. S., Chung, S. J., Kim, D. D. (2009). Docetaxel Microemulsion for Enhanced Oral Bioavailability: Preparation and in Vitro and in Vivo Evaluation. *Journal of Controlled Release*, 140, 86-94.
- Yunita, E., Yulianto, D., Fatimah, S., & Firanita, T., 2020, Validation of UV-Vis Spectrophotometric Method of Quercetin in Ethanol Extract of Tamarind Leaf, *Journal of Fundamental and Applied Pharmaceutical Science*, 1(1).
- Zhang, Z., Lei, Y., Yang, X., Shi, N., Geng, L., Wang, S., Zhang, J., & Shi, S. (2019). High drug-loading system of hollow carbon dots-doxorubicin: Preparation, in vitro release and pH-targeted research. *Journal of Materials Chemistry B*, 7(13), 2130–2137. <https://doi.org/10.1039/c9tb00032a>
- Zupancic, O., Grieinger, J. A., Rohrer, J., Pereira de Sousa, I., Danninger, L., Partenhauser, A., Bernkop-Schnürch, A. (2016). Development, in Vitro and in Vivo Evaluation of a Self-Emulsifying Drug Delivery System (SEDDS) for Oral Enoxaparin Administration. *European Journal of Pharmaceutics and Biopharmaceutics*, 109, 113-121.