

REFERENCES

- Akter, J., Hassan, J., Rahman, M. M., Biswas, M. S., Khan, H. I., Rahman Rajib, M. M., Ahmed, M. R., Noor-E-Azam Khan, M., & Ahamed Hasan, M. F. (2024). Colour, nutritional composition and antioxidant properties of dehydrated carrot (*Daucus carota* var. *Sativus*) using solar drying techniques and pretreatments. *Heliyon*, *10*(2), e24165. <https://doi.org/10.1016/j.heliyon.2024.e24165>
- AOAC. (2016). *Appendix F: Guidelines for Standard Method Performance Requirements*. AOAC International.
- Ardiansyah, A., Naufalin, R., Arsil, P., Latifasari, N., Wicaksono, R., Aliim, M. S., Kartiko, C., & Waluyo, S. (2022). Machine Learning Model for Quality Parameters Prediction and Control System Design in the Kecombrang Flower (*Etlingera elatior*) Extraction Process. *Processes*, *10*(7). Scopus. <https://doi.org/10.3390/pr10071341>
- Arévalo-Rodrigues, G., Cardoso-Gustavson, P., & Barros, F. (2022). Floral gland diversity in Pleurothallidinae (Epidendroideae–Orchidaceae). *Flora*, *297*, 152179. <https://doi.org/10.1016/j.flora.2022.152179>
- Aziman, N., Abdullah, N., Noor, Z. M., Kamarudin, W. S. S. W., & Zulkifli, K. S. (2014). Phytochemical Profiles and Antimicrobial Activity of Aromatic Malaysian Herb Extracts against Food-Borne Pathogenic and Food Spoilage Microorganisms. *Journal of Food Science*, *79*(4), M583–M592. Scopus. <https://doi.org/10.1111/1750-3841.12419>

- Baş, D., & Boyacı, İ. H. (2007). Modeling and optimization I: Usability of response surface methodology. *Journal of Food Engineering*, 78(3), 836–845.
<https://doi.org/10.1016/j.jfoodeng.2005.11.024>
- Baysal, T., Icier, F., Ersus, S., & Yıldız, H. (2003). Effects of microwave and infrared drying on the quality of carrot and garlic. *European Food Research and Technology*, 218(1), 68–73. <https://doi.org/10.1007/s00217-003-0791-3>
- Bender, O., & Atalay, A. (2021). Chapter 28—Polyphenol chlorogenic acid, antioxidant profile, and breast cancer. In V. R. Preedy & V. B. Patel (Eds.), *Cancer (Second Edition)* (pp. 311–321). Academic Press.
<https://doi.org/10.1016/B978-0-12-819547-5.00028-6>
- Berger, P. D., Maurer, R. E., & Celli, G. B. (2017). *Experimental Design: With Application in Management, Engineering, and the Sciences*. Springer.
- Bezerra, M. A., Santelli, R. E., Oliveira, E. P., Villar, L. S., & Escaleira, L. A. (2008). Response surface methodology (RSM) as a tool for optimization in analytical chemistry. *Talanta*, 76(5), 965–977.
<https://doi.org/10.1016/j.talanta.2008.05.019>
- Boonpeng, S., Siripongvutikorn, S., Sae-wong, C., & Sutthirak, P. (2014). The antioxidant and anti-cadmium toxicity properties of garlic extracts. *Food Science & Nutrition*, 2(6), 792–801. <https://doi.org/10.1002/fsn3.164>
- Butt, W., & Ahmed, D. (2023). Modeling and optimization of sustainable extraction of valuable bioactive compounds from *Foeniculum vulgare* by glycerol-based deep eutectic solvent and ultrasound and heat treatment. *Food*

Chemistry Advances, 3, 100535.

<https://doi.org/10.1016/j.focha.2023.100535>

Cares, M. G., Vargas, Y., Gaete, L., Sainz, J., & Alarcón, J. (2010). Ultrasonically assisted extraction of bioactive principles from *Quillaja Saponaria* Molina.

Physics Procedia, 3(1), 169–178.

<https://doi.org/10.1016/j.phpro.2010.01.024>

Carrasco-Sandoval, J., Falcó, I., Sánchez, G., Fabra, M. J., López-Rubio, A., Rodríguez, A., Henríquez-Aedo, K., & Aranda, M. (2022). Multivariable optimization of ultrasound-assisted extraction for the determination of phenolic and antioxidants compounds from arrayan (*Luma apiculata* (DC.) Burret) leaves by microplate-based methods and mass spectrometry.

Journal of Applied Research on Medicinal and Aromatic Plants, 28, 100356. <https://doi.org/10.1016/j.jarmap.2021.100356>

Chan, E. W. C., Lim, Y. Y., Ling, S. K., Tan, S. P., Lim, K. K., & Khoo, M. G. H. (2009). Caffeoylquinic acids from leaves of *Etilingera* species (Zingiberaceae). *LWT - Food Science and Technology*, 42(5), 1026–1030.

<https://doi.org/10.1016/j.lwt.2009.01.003>

Chan, E. W. C., Lim, Y. Y., Wong, S. K., Lim, K. K., Tan, S. P., Lianto, F. S., & Yong, M. Y. (2009a). Effects of different drying methods on the antioxidant properties of leaves and tea of ginger species. *Food Chemistry*, 113(1), 166–172. <https://doi.org/10.1016/j.foodchem.2008.07.090>

Chan, E. W. C., Lim, Y. Y., Wong, S. K., Lim, K. K., Tan, S. P., Lianto, F. S., & Yong, M. Y. (2009b). Effects of different drying methods on the antioxidant

properties of leaves and tea of ginger species. *Food Chemistry*, 113(1), 166–172. <https://doi.org/10.1016/j.foodchem.2008.07.090>

Choon, S. Y., & Ding, P. (2016). Growth Stages of Torch Ginger (*Etilingera elatior*) Plant. *Sains Malaysiana*, 45(4), 507–515.

Dawidowicz, A. L., & Typek, R. (2017). Transformation of chlorogenic acids during the coffee beans roasting process. *European Food Research and Technology*, 243(3), 379–390. <https://doi.org/10.1007/s00217-016-2751-8>

Ernilasari, Walil, K., Fitmawati, Roslim, D. E. W. I. I., Zumaidar, Saudah, & Rayhannisa. (2021). Antibacterial activity of leaves, flowers, and fruits extract of *etlingera elatior* from nagan raya district, indonesia against *escherichia coli* and *staphylococcus aureus*. *Biodiversitas*, 22(10), 4457–4464. Scopus. <https://doi.org/10.13057/biodiv/d221039>

Ferreira, S. L. C., Bruns, R. E., Ferreira, H. S., Matos, G. D., David, J. M., Brandão, G. C., da Silva, E. G. P., Portugal, L. A., dos Reis, P. S., Souza, A. S., & dos Santos, W. N. L. (2007). Box-Behnken design: An alternative for the optimization of analytical methods. *Analytica Chimica Acta*, 597(2), 179–186. <https://doi.org/10.1016/j.aca.2007.07.011>

Fraser, C. M., & Chapple, C. (2011). The phenylpropanoid pathway in *Arabidopsis*. *The Arabidopsis Book*, 9, e0152. <https://doi.org/10.1199/tab.0152>

Fudholi, A., Othman, M. Y., Ruslan, M. H., & Sopian, K. (2013). Drying of Malaysian *Capsicum annuum* L. (Red Chili) Dried by Open and Solar Drying. *International Journal of Photoenergy*, 2013, e167895. <https://doi.org/10.1155/2013/167895>

- Gąsecka, M., Siwulski, M., Magdziak, Z., Budzyńska, S., Stuper-Szablewska, K.,
Niedzielski, P., & Mleczek, M. (2020). The effect of drying temperature on
bioactive compounds and antioxidant activity of *Leccinum scabrum* (Bull.)
Gray and *Hericium erinaceus* (Bull.) Pers. *Journal of Food Science and
Technology*, 57(2), 513–525. <https://doi.org/10.1007/s13197-019-04081-1>
- Ghasemzadeh, A., Jaafar, H. Z. E., Rahmat, A., & Ashkani, S. (2015). Secondary
metabolites constituents and antioxidant, anticancer and antibacterial
activities of *Etlingera elatior* (Jack) R.M.Sm grown in different locations of
Malaysia. *BMC Complementary and Alternative Medicine*, 15(1), 335.
<https://doi.org/10.1186/s12906-015-0838-6>
- Ghorbannezhad, P., Bay, A., Yolmeh, M., Yadollahi, R., & Moghadam, J. Y.
(2016). Optimization of coagulation–flocculation process for medium
density fiberboard (MDF) wastewater through response surface
methodology. *Desalination and Water Treatment*, 57(56), 26916–26931.
<https://doi.org/10.1080/19443994.2016.1170636>
- Gray, J., Caparrós-Ruiz, D., & Grotewold, E. (2012). Grass phenylpropanoids:
Regulate before using! *Plant Science*, 184, 112–120.
<https://doi.org/10.1016/j.plantsci.2011.12.008>
- Gupta, A., Atanasov, A. G., Li, Y., Kumar, N., & Bishayee, A. (2022). Chlorogenic
acid for cancer prevention and therapy: Current status on efficacy and
mechanisms of action. *Pharmacological Research*, 186, 106505.
<https://doi.org/10.1016/j.phrs.2022.106505>

- Hapsari, S., Yohed, I., Kristianita, R. A., Jadid, N., Aparamarta, H. W., & Gunawan, S. (2022). Phenolic and flavonoid compounds extraction from *Calophyllum inophyllum* leaves. *Arabian Journal of Chemistry*, *15*(3), 103666. <https://doi.org/10.1016/j.arabjc.2021.103666>
- Harmita. (2004). PETUNJUK PELAKSANAAN VALIDASI METODE DAN CARA PERHITUNGANNYA. *Pharmaceutical Sciences and Research (PSR)*. <http://journal.ui.ac.id/index.php/mik/article/view/1136>
- Hu, Y., Yang, L., Liang, Z., Chen, J., Zhao, M., & Tang, Q. (2023). Comparative analysis of flavonoids extracted from *Dendrobium chrysotoxum* flowers by supercritical fluid extraction and ultrasonic cold extraction. *Sustainable Chemistry and Pharmacy*, *36*, 101267. <https://doi.org/10.1016/j.scp.2023.101267>
- ICH. (2005). *Q2(R1) Validation of Analytical Procedures: Text and Methodology Guidance for Industry*.
- Isleroglu, H., & Turker, I. (2022). Ultrasonic-assisted extraction and thermal stability of phytochemicals from fenugreek leaves. *Journal of Applied Research on Medicinal and Aromatic Plants*, *30*, 100390. <https://doi.org/10.1016/j.jarmap.2022.100390>
- Jackie, T., Haleagrahara, N., & Chakravarthi, S. (2011). Antioxidant effects of *Etlingera elatior* flower extract against lead acetate—Induced perturbations in free radical scavenging enzymes and lipid peroxidation in rats. *BMC Research Notes*, *4*(1), 67. <https://doi.org/10.1186/1756-0500-4-67>

- Jaisankar, A. I., & Arivarasu, L. (2020). Free Radical Scavenging and Anti-Inflammatory Activity of Chlorogenic Acid Mediated Silver Nanoparticle. *Journal of Pharmaceutical Research International*, 106–112. <https://doi.org/10.9734/jpri/2020/v32i1930715>
- Juwita, T., Melyani Pu, I., & Levita, J. (2018). Torch Ginger (*Etlingera elatior*): A Review on its Botanical Aspects, Phytoconstituents and Pharmacological Activities. *Pakistan Journal of Biological Sciences*, 21(4), 151–165. <https://doi.org/10.3923/pjbs.2018.151.165>
- Klein, B. D., & Rossin, D. F. (1999). Data quality in neural network models: Effect of error rate and magnitude of error on predictive accuracy. *Omega*, 27(5), 569–582. [https://doi.org/10.1016/S0305-0483\(99\)00019-5](https://doi.org/10.1016/S0305-0483(99)00019-5)
- Kusuma, H. S., Izzah, D. N., & Linggajati, I. W. L. (2023). Microwave-assisted drying of *Ocimum sanctum* leaves: Analysis of moisture content, drying kinetic model, and techno-economics. *Applied Food Research*, 3(2), 100337. <https://doi.org/10.1016/j.afres.2023.100337>
- Lachumy, S. J. T., Sasidharan, S., Sumathy, V., & Zuraini, Z. (2010). Pharmacological activity, phytochemical analysis and toxicity of methanol extract of *Etlingera elatior* (torch ginger) flowers. *Asian Pacific Journal of Tropical Medicine*, 3(10), 769–774. [https://doi.org/10.1016/S1995-7645\(10\)60185-X](https://doi.org/10.1016/S1995-7645(10)60185-X)
- Leardi, R. (2009). Experimental design in chemistry: A tutorial. *Analytica Chimica Acta*, 652(1), 161–172. <https://doi.org/10.1016/j.aca.2009.06.015>

Li, H., He, W., Xu, S., Wang, R., Ge, S., Xu, H., Shan, Y., & Ding, S. (2024).

Grafting chlorogenic acid enhanced the antioxidant activity of curdlan oligosaccharides and modulated gut microbiota. *Food Chemistry: X*, *21*, 101075. <https://doi.org/10.1016/j.fochx.2023.101075>

Liang, N., & Kitts, D. (2015). Role of Chlorogenic Acids in Controlling Oxidative and Inflammatory Stress Conditions. *Nutrients*, *8*(1), 16. <https://doi.org/10.3390/nu8010016>

Liu, W., Zhao, Y., Sun, J., Li, G., Shan, Y., & Chen, P. (2017). Study the effects of drying processes on chemical compositions in daylily flowers using flow injection mass spectrometric fingerprinting method and chemometrics. *Food Research International*, *102*, 493–503. <https://doi.org/10.1016/j.foodres.2017.09.015>

Liu, Y., Liu, H.-Y., Xia, Y., Guo, H., He, X.-Q., Li, H., Wu, D.-T., Geng, F., Lin, F.-J., Li, H.-B., Zhuang, Q.-G., & Gan, R.-Y. (2021). Screening and process optimization of ultrasound-assisted extraction of main antioxidants from sweet tea (*Lithocarpus litseifolius* [Hance] Chun). *Food Bioscience*, *43*, 101277. <https://doi.org/10.1016/j.fbio.2021.101277>

Lou, Z., Wang, H., Zhu, S., Ma, C., & Wang, Z. (2011). Antibacterial activity and mechanism of action of chlorogenic acid. *Journal of Food Science*, *76*(6), M398-403. <https://doi.org/10.1111/j.1750-3841.2011.02213.x>

Lu, J., Wang, Z., Qin, L., Shen, J., He, Z., Shao, Q., & Lin, D. (2021). Drying methods affect bioactive compound contents and antioxidant capacity of

Bletilla striata (Thunb.) Reichb.f. Flower. *Industrial Crops and Products*, 164, 113388. <https://doi.org/10.1016/j.indcrop.2021.113388>

Luo, B., Wen, Y., Ye, F., Wu, Y., Li, N., Farid, M. S., Chen, Z., El-Seedi, H. R., & Zhao, C. (2023). Bioactive phytochemicals and their potential roles in modulating gut microbiota. *Journal of Agriculture and Food Research*, 100583. <https://doi.org/10.1016/j.jafr.2023.100583>

M, B., C, H., M, P.-C., & W, B. (2003). Lignin: Genetic engineering and impact on pulping. *Critical Reviews in Biochemistry and Molecular Biology*, 38(4). <https://doi.org/10.1080/10409230391036757>

Mahdavi, B., Yaacob, W., & Din, L. (2017). Chemical composition, antioxidant, and antibacterial activity of essential oils from *Etlingera sayapensis* A.D. Poulsen & Ibrahim. *Asian Pacific Journal of Tropical Medicine*, 10. <https://doi.org/10.1016/j.apjtm.2017.08.006>

Mbondo, N. N., Owino, W. O., Ambuko, J., & Sila, D. N. (2018). Effect of drying methods on the retention of bioactive compounds in African eggplant. *Food Science & Nutrition*, 6(4), 814–823. <https://doi.org/10.1002/fsn3.623>

Miao, J., Liu, J., Gao, X., Lu, F., & Yang, X. (2022). Effects of different drying methods on chemical compositions, antioxidant activity and anti- α -glucosidase activity of *Coreopsis tinctoria* flower tea. *Heliyon*, 8(11), e11784. <https://doi.org/10.1016/j.heliyon.2022.e11784>

Myo, H., Yaowiwat, N., Pongkorpsakol, P., Aonbangkhen, C., & Khat-Udomkiri, N. (2023). Butylene Glycol Used as a Sustainable Solvent for Extracting Bioactive Compounds from *Camellia sinensis* Flowers with Ultrasound-

Assisted Extraction. *ACS Omega*, 8(5), 4976–4987. Scopus.

<https://doi.org/10.1021/acsomega.2c07481>

Naufalin, R., Sutrisna, E., & Wicaksono, R. (2021). *Antioxidant potential ingredient of kecombrang plants (Etlingera elatior)*. 653(1). Scopus.

<https://doi.org/10.1088/1755-1315/653/1/012130>

Naveed, M., Hejazi, V., Abbas, M., Kamboh, A. A., Khan, G. J., Shumzaid, M., Ahmad, F., Babazadeh, D., FangFang, X., Modarresi-Ghazani, F., WenHua, L., & XiaoHui, Z. (2018). Chlorogenic acid (CGA): A pharmacological review and call for further research. *Biomedicine & Pharmacotherapy = Biomedecine & Pharmacotherapie*, 97, 67–74.

<https://doi.org/10.1016/j.biopha.2017.10.064>

Nguyen, Q. V., & Chuyen, H. V. (2020). Processing of Herbal Tea from Roselle (*Hibiscus sabdariffa* L.): Effects of Drying Temperature and Brewing Conditions on Total Soluble Solid, Phenolic Content, Antioxidant Capacity and Sensory Quality. *Beverages*, 6(1), Article 1.

<https://doi.org/10.3390/beverages6010002>

Noordin, L., Ahmad, W. A. N. W., Nor, N. A. M., Bakar, N. H. A., & Ugusman, A. (2022). *Etlingera elatior* Flower Aqueous Extract Protects against Oxidative Stress-Induced Nephropathy in a Rat Model of Type 2 Diabetes. *Evidence-Based Complementary and Alternative Medicine*, 2022. Scopus.

<https://doi.org/10.1155/2022/2814196>

Nuryanti, S., Latifasari, N., Naufalin, R., Wicaksono, R., & Erminawati. (2021). Antioxidant activity and total phenol of extract of kecombrang flower, stem

and leaves with different types of solutions. *Molekul*, 16(2), 110–116.

Scopus. <https://doi.org/10.20884/1.jm.2021.16.2.631>

Oehlert, G. W. (2010). *A First Course in Design and Analysis of Experiments*.

<http://conservancy.umn.edu/handle/11299/168002>

Ozdemir, M., Gungor, V., Melikoglu, M., & Aydiner, C. (2024). Solvent selection and effect of extraction conditions on ultrasound-assisted extraction of phenolic compounds from galangal (*Alpinia officinarum*). *Journal of Applied Research on Medicinal and Aromatic Plants*, 38, 100525.

<https://doi.org/10.1016/j.jarmap.2023.100525>

Pai, M.-S., Wang, K.-C., Yeh, K.-C., & Wang, S.-J. (2023). Stabilization of mitochondrial function by chlorogenic acid protects against kainic acid-induced seizures and neuronal cell death in rats. *European Journal of Pharmacology*, 961, 176197. <https://doi.org/10.1016/j.ejphar.2023.176197>

Patras, A., Brunton, Nigel. P., O'Donnell, C., & Tiwari, B. K. (2010). Effect of thermal processing on anthocyanin stability in foods; mechanisms and kinetics of degradation. *Trends in Food Science & Technology*, 21(1), 3–11. <https://doi.org/10.1016/j.tifs.2009.07.004>

Phumthum, M., Balslev, H., & Barfod, A. S. (2019). Important Medicinal Plant Families in Thailand. *Frontiers in Pharmacology*, 10, 1125. <https://doi.org/10.3389/fphar.2019.01125>

Pinar, H., Çetin, N., Ciftci, B., Karaman, K., & Kaplan, M. (2021). Biochemical composition, drying kinetics and chromatic parameters of red pepper as

affected by cultivars and drying methods. *Journal of Food Composition and Analysis*, 102, 103976. <https://doi.org/10.1016/j.jfca.2021.103976>

Prakash Maran, J., Sivakumar, V., Sridhar, R., & Prince Immanuel, V. (2013).

Development of model for mechanical properties of tapioca starch based edible films. *Industrial Crops and Products*, 42, 159–168. <https://doi.org/10.1016/j.indcrop.2012.05.011>

Rachkeeree, A., Kantadoung, K., Suksathan, R., Puangpradab, R., Page, P. A., &

Sommano, S. R. (2018). Nutritional Compositions and Phytochemical Properties of the Edible Flowers from Selected Zingiberaceae Found in Thailand. *Frontiers in Nutrition*, 5, 3. <https://doi.org/10.3389/fnut.2018.00003>

Saleh, I. A., Vinatoru, M., Mason, T. J., Abdel-Azim, N. S., Aboutabl, E. A., &

Hammouda, F. M. (2016). A possible general mechanism for ultrasound-assisted extraction (UAE) suggested from the results of UAE of chlorogenic acid from *Cynara scolymus* L. (artichoke) leaves. *Ultrasonics Sonochemistry*, 31, 330–336. <https://doi.org/10.1016/j.ultsonch.2016.01.002>

Santanatoglia, A., Angeloni, S., Fiorito, M., Fioretti, L., Ricciutelli, M., Sagratini,

G., Vittori, S., & Caprioli, G. (2024). Development of new analytical methods for the quantification of organic acids, chlorogenic acids and caffeine in espresso coffee by using solid-phase extraction (SPE) and high-performance liquid chromatography-diode array detector (HPLC-DAD).

Journal of Food Composition and Analysis, 125, 105732.

<https://doi.org/10.1016/j.jfca.2023.105732>

Sari, K. R. P., Ikawati, Z., Danarti, R., & Hertiani, T. (2023). Micro-titer plate assay for measurement of total phenolic and total flavonoid contents in medicinal plant extracts. *Arabian Journal of Chemistry*, 16(9), 105003.

<https://doi.org/10.1016/j.arabjc.2023.105003>

Saudah, S., Zumaidar, Z., Darusman, D., Roslim, D. I., & Ernilasari, E. (2022). Ethnobotanical knowledge of *Etlingera elatior* for medicinal and food uses among ethnic groups in Aceh Province, Indonesia. *Biodiversitas Journal of Biological Diversity*, 23(8), Article 8.

<https://doi.org/10.13057/biodiv/d230862>

Sereshti, H., Samadi, S., & Jalali-Heravi, M. (2013). Determination of volatile components of green, black, oolong and white tea by optimized ultrasound-assisted extraction-dispersive liquid-liquid microextraction coupled with gas chromatography. *Journal of Chromatography A*, 1280, 1–8.

<https://doi.org/10.1016/j.chroma.2013.01.029>

Sheng, X., Zan, J., Jiang, Y., Shen, S., Li, L., & Yuan, H. (2023). Data fusion strategy for rapid prediction of moisture content during drying of black tea based on micro-NIR spectroscopy and machine vision. *Optik*, 276, 170645.

<https://doi.org/10.1016/j.ijleo.2023.170645>

Shi, X.-F., Chu, J.-Z., Zhang, Y.-F., Liu, C.-Q., & Yao, X.-Q. (2017). Nutritional and active ingredients of medicinal chrysanthemum flower heads affected

by different drying methods. *Industrial Crops and Products*, 104, 45–51.

<https://doi.org/10.1016/j.indcrop.2017.04.021>

Shivanna, S. K., Naik N, L., & Rao, P. S. (2024). Impact of drying techniques on the composition, physicochemical attributes, antioxidant capacity, antidiabetic potential, and ACE inhibition properties of *Moringa oleifera* pod pulp powder. *South African Journal of Botany*, 165, 405–416.

<https://doi.org/10.1016/j.sajb.2024.01.008>

Simanjuntak, M. E., Ristiarini, S., & Widyawati, P. S. (2022). The effect of rotary drying temperature on drying characteristic and antioxidant activity of *Etlingera elatior* Jack. *Food Research*, 6(3), 196–202. Scopus.

[https://doi.org/10.26656/fr.2017.6\(3\).333](https://doi.org/10.26656/fr.2017.6(3).333)

Singh, N., & Yadav, S. S. (2022). A review on health benefits of phenolics derived from dietary spices. *Current Research in Food Science*, 5, 1508–1523.

<https://doi.org/10.1016/j.crfs.2022.09.009>

Sulaiman, S. F., Sajak, A. A. B., Ooi, K. L., Supriatno, & Seow, E. M. (2011). Effect of solvents in extracting polyphenols and antioxidants of selected raw vegetables. *Journal of Food Composition and Analysis*, 24(4), 506–515.

<https://doi.org/10.1016/j.jfca.2011.01.020>

Suryani, N., Nurjanah, D., & Indriatmoko, D. (2019). Aktivitas Antibakteri Ekstrak Batang Kecombrang (*Etlingera elatior* (Jack) R.M.Sm.) Terhadap Bakteri Plak Gigi *Streptococcus mutans*. *Jurnal Kartika Kimia*, 2.

<https://doi.org/10.26874/jkk.v2i1.19>

- Syafriana, V., Purba, R. N., & Djuhariah, Y. S. (2021). Antibacterial Activity of Kecombrang Flower (*Etlingera elatior* (Jack) R.M. Sm) Extract against *Staphylococcus epidermidis* and *Propionibacterium acnes*. *Journal of Tropical Biodiversity and Biotechnology*, 6(1). Scopus. <https://doi.org/10.22146/jtbb.58528>
- Tacherfiout, M., Kherbachi, S., Kheniche, M., Mattonai, M., Degano, I., Ribechini, E., & Khettal, B. (2022). HPLC-DAD and HPLC-ESI-MS-MS profiles of hydroalcoholic extracts of *Chamaemelum nobile* and *Mentha pulegium*, and study of their antihemolytic activity against AAPH-induced hemolysis. *South African Journal of Botany*, 150, 678–690. <https://doi.org/10.1016/j.sajb.2022.08.001>
- Tajik, N., Tajik, M., Mack, I., & Enck, P. (2017). The potential effects of chlorogenic acid, the main phenolic components in coffee, on health: A comprehensive review of the literature. *European Journal of Nutrition*, 56(7), 2215–2244. <https://doi.org/10.1007/s00394-017-1379-1>
- Takahashi, J. A., Rezende, F. A. G. G., Moura, M. A. F., Dominguet, L. C. B., & Sande, D. (2020). Edible flowers: Bioactive profile and its potential to be used in food development. *Food Research International*, 129, 108868. <https://doi.org/10.1016/j.foodres.2019.108868>
- Tomac, I., Šeruga, M., & Labuda, J. (2020). Evaluation of antioxidant activity of chlorogenic acids and coffee extracts by an electrochemical DNA-based biosensor. *Food Chemistry*, 325, 126787. <https://doi.org/10.1016/j.foodchem.2020.126787>

- Ucella-Filho, J. G. M., Freire, A. da S. M., Carréra, J. C., Lucas, F. M. F., Zucolotto, S. M., Dias Júnior, A. F., & Mori, F. A. (2022). Tannin-rich bark extract of plants as a source of antimicrobial bioactive compounds: A bibliometric analysis. *South African Journal of Botany*, *150*, 1038–1050. <https://doi.org/10.1016/j.sajb.2022.09.018>
- Vo, T. P., Ho, M. T., Nguyen Nguyen, P. U., Pham, N. D., Truong, K. V., Yen Nguyen, T. H., Nguyen, D. Q., & Huong Vo, T. T. (2024). Extracting phenolics, flavonoids, and terpenoids from *Codonopsis pilosula* using green solvents. *Sustainable Chemistry and Pharmacy*, *37*, 101395. <https://doi.org/10.1016/j.scp.2023.101395>
- Wale, K., Tolessa, K., Atlabachew, M., Mehari, B., Alemayehu, M., Mengistu, D. A., & Kerisew, B. (2024). Level of caffeine, trigonelline and chlorogenic acids in green coffee (*Coffea arabica* L.) beans from Amhara region, Ethiopia. *Journal of Agriculture and Food Research*, *16*, 101082. <https://doi.org/10.1016/j.jafr.2024.101082>
- Wanderley, R. de O. S., Figueirêdo, R. M. F. de, Queiroz, A. J. de M., Santos, F. S. dos, Silva, A. P. de F., Paiva, Y. F., Moura, H. V., Silva, E. T. de V., Carvalho, A. J. de B. A., Lima, M. dos S., Campos, A. R. N., Gregório, M. G., & Lima, A. G. B. de. (2023). Effect of drying temperature on antioxidant activity, phenolic compound profile and hygroscopic behavior of pomegranate peel and seed flours. *LWT*, *189*, 115514. <https://doi.org/10.1016/j.lwt.2023.115514>

- Wianowska, D., & Gil, M. (2019). Recent advances in extraction and analysis procedures of natural chlorogenic acids. *Phytochemistry Reviews*, 18. <https://doi.org/10.1007/s11101-018-9592-y>
- Wijekoon, M. M. J. O., Bhat, R., & Karim, A. A. (2011). Effect of extraction solvents on the phenolic compounds and antioxidant activities of bunga kantan (*Etlingera elatior* Jack.) inflorescence. *Journal of Food Composition and Analysis*, 24(4), 615–619. <https://doi.org/10.1016/j.jfca.2010.09.018>
- Wulandari, A., Nurlaelasari, A., Ari, H., Cahyadi, M., Kurniawan, W., & Hamid, P. (2023). Ethanolic extract of *Etlingera elatior* flower exhibits anthelmintic properties to *Fasciola gigantica* in vitro. *Open Veterinary Journal*, 13, 576. <https://doi.org/10.5455/OVJ.2023.v13.i5.10>
- Xiao, J., Zou, Y., Wen, X., Guo, Y., Hu, F., Chen, G., Wu, Z., Lin, Y., Wang, Z., Sun, L., Pan, Y., & Li, N. (2022). Functional contents and antioxidant potency of Chinese Wenguan flower tea. *Food Control*, 138, 109002. <https://doi.org/10.1016/j.foodcont.2022.109002>
- Yan, Y., Zhou, X., Guo, K., Zhou, F., & Yang, H. (2020). Chlorogenic Acid Protects Against Indomethacin-Induced Inflammation and Mucosa Damage by Decreasing Bacteroides-Derived LPS. *Frontiers in Immunology*, 11. <https://doi.org/10.3389/fimmu.2020.01125>
- Yasar, B., Kutlu, G., & Tornuk, F. (2022). Edible flowers as sources of bioactive compounds: Determination of phenolic extraction conditions. *International Journal of Gastronomy and Food Science*, 30, 100618. <https://doi.org/10.1016/j.ijgfs.2022.100618>

- Yuan, L., & Grotewold, E. (2020). Plant specialized metabolism. *Plant Science*, 298, 110579. <https://doi.org/10.1016/j.plantsci.2020.110579>
- Yunus, M. F., Meenakshi Sundram, T. C., Zainuddin, Z., Ismail, N. A., Mohd Rosli, N., & Mohamad Hamdan, M. S. (2022). Current status and biotechnological development of *Etlingera elatior*, a promising horticultural and medicinal plant. *Journal of Horticultural Science and Biotechnology*, 97(4), 429–436. Scopus. <https://doi.org/10.1080/14620316.2021.2021812>
- Zainol, N., Aziz, N. H., & Baharudin, A. S. (2023). Influence of agitation and solvent percentage on the extraction of phytochemical compound from *Asystasia gangetica*. *Food Chemistry Advances*, 3, 100538. <https://doi.org/10.1016/j.focha.2023.100538>